

FREE BOUNDARY REGULARITY FOR ALMOST MINIMIZERS OF THE PARABOLIC SIGNORINI PROBLEM

SEONGMIN JEON AND ARSHAK PETROSYAN

ABSTRACT. In this paper, we study the regularity of the “regular” part of the free boundary for almost minimizers in the parabolic Signorini problem with zero thin obstacle. This work is a continuation of our earlier research on the regularity of almost minimizers. We first establish the Weiss-type monotonicity formula by comparing almost minimizers with parabolically homogeneous replacements and utilizing conformal self-similar coordinates. Subsequently, by deriving the Almgren-type frequency formula and applying the epiperimetric inequality, we obtain the optimal growth near regular free boundary points and achieve the regularity of the regular set.

CONTENTS

1. Introduction	1
1.1. Solutions of parabolic Signorini problem	1
1.2. Almost minimizers	3
1.3. Main results	4
2. Notation and Preliminaries	6
2.1. Notation	6
2.2. Preliminaries	7
3. Weiss- and Almgren-type monotonicity formulas	7
4. Almgren rescalings and blowups	15
5. Growth estimates	22
6. 3/2-Homogeneous blowups	29
7. Regularity of the regular set	33
Appendix A. Existence of weak solutions	36
Appendix B. Examples of almost minimizers	37
References	45

1. INTRODUCTION

1.1. Solutions of parabolic Signorini problem. Let Ω be a domain in \mathbb{R}^n , $n \geq 2$, and \mathcal{M} be a smooth $(n - 1)$ -dimensional manifold that divides Ω into two parts: $\Omega \setminus \mathcal{M} = \Omega^+ \cup \Omega^-$. For $T > 0$, we set $\Omega_T := \Omega \times (0, T]$, $\mathcal{M}_T := \mathcal{M} \times (0, T]$ (*the thin space*), and $(\partial\Omega)_T := \partial\Omega \times (0, T]$. Let also $\varphi : \mathcal{M}_T \rightarrow \mathbb{R}$ (*the thin obstacle*),

2020 *Mathematics Subject Classification.* Primary 35R35.

Key words and phrases. Almost minimizers, parabolic thin obstacle (or Signorini) problem, regular set, Weiss-type monotonicity formula, Almgren-type frequency, epiperimetric inequality.

S. Jeon was supported by the Academy of Finland grant 347550 and the research fund of Hanyang University(HY-202400000003278).

$\varphi_0 : \Omega \times \{0\} \rightarrow \mathbb{R}$ (*the initial value*), and $g : (\partial\Omega)_T \rightarrow \mathbb{R}$ (*the boundary value*) be prescribed functions satisfying the compatibility conditions: $\varphi_0 \geq \varphi$ on $\mathcal{M} \times \{0\}$, $g \geq \varphi$ on $(\mathcal{M} \cap \partial\Omega) \times (0, T]$, and $g = \varphi_0$ on $\partial\Omega \times \{0\}$.

We then say that a function $u \in W_2^{1,0}(\Omega_T)$ (see Subsection 2.1 for notations) is a solution of the *parabolic thin obstacle* (or *Signorini*) *problem* in Ω_T , if it satisfies the variational inequality

$$\begin{aligned} \int_{\Omega_T} \nabla u \nabla (v - u) + \partial_t u (v - u) &\geq 0 \quad \text{for any } v \in \mathcal{K}, \\ u \in \mathcal{K}, \partial_t u &\in L^2(\Omega_T), u(\cdot, 0) = \varphi_0 \text{ on } \Omega, \end{aligned}$$

where $\mathcal{K} = \{v \in W_2^{1,0}(\Omega_T) : v \geq \varphi \text{ on } \mathcal{M}_T, v = g \text{ on } (\partial\Omega)_T\}$. It is known that the solution u satisfies

$$\begin{aligned} \Delta u - \partial_t u &= 0 \text{ in } \Omega_T \setminus \mathcal{M}_T, \\ u &\geq \varphi, \partial_{\nu^+} u + \partial_{\nu^-} u \geq 0, (u - \varphi)(\partial_{\nu^+} u + \partial_{\nu^-} u) = 0 \text{ on } \mathcal{M}_T, \\ u &= g \text{ on } (\partial\Omega)_T, \\ u(\cdot, 0) &= \varphi_0 \text{ on } \Omega \times \{0\}, \end{aligned}$$

to be understood in a certain weak sense, where ν^\pm are the outer unit normal to Ω^\pm on \mathcal{M} .

In the study of the above problem, the main interests are

- the regularity of the solution u ,
- the regularity and structure of the *free boundary*

$$\Gamma(u) = \partial_{\mathcal{M}_T} \{(x, t) \in \mathcal{M}_T : u(x, t) > \varphi(x, t)\}.$$

A comprehensive examination of both regularity of the solution and the properties of the free boundary was conducted in [DGPT17] by the second author, Danielli, Garofalo and To, under the condition that the thin manifold \mathcal{M} is flat (cf. refer to [AS24] for the variable coefficients parabolic Signorini problem). Specifically, they established a generalized frequency formula, and employed it to achieve the optimal $H^{3/2, 3/4}$ regularity of the solution and classified the free boundary points according to their frequency limits. [DGPT17] dealt with two special types of subsets of the free boundary: the *regular set* and the *singular set*.

The regular set is defined as the set of the free boundary points with minimal frequency $3/2$. Similarly to the elliptic Signorini problem [ACS08, CSS08, PSU12], [DGPT17] showed that there is a cone of spatial direction in which $u - \varphi$ is monotone. Combining this with the fact that the blowups at regular points are time independent, they obtained the Lipschitz regularity of the regular set in the space variables. Moreover, by applying the parabolic boundary Harnack principles with thin Lipschitz complement, they proved that the regular set is given locally as a graph with $H^{\alpha, \alpha/2}$ regular gradient.

The singular set corresponds to the free boundary points with frequency $2m$, $m \in \mathbb{N}$, which have the characterization that the coincidence set $\{u = \varphi\}$ has zero H^n -density in the thin manifold \mathcal{M}_T . Following the approach in [GP09] by the second author and Garofalo, [DGPT17] established Weiss- and Monneau-type monotonicity formulas and applied the parabolic version of the Whitney's extension theorem to achieve the C^1 structure of the singular set.

1.2. Almost minimizers. In this paper we investigate the almost minimizers concerning the parabolic Signori problem described above. It serves as a continuation of [JP23], where the authors previously explored the regularity of almost minimizers. For technical reasons, we consider two different notions of almost minimizers: “unweighted” almost minimizers and “weighted” almost minimizers. We first introduce unweighted almost minimizers, which correspond to those studied in [JP23].

We let φ be the thin obstacle on Q'_1 . Given $r_0 > 0$, we say that $\eta : [0, r_0) \rightarrow [0, \infty)$ is a *gauge function* or a modulus of continuity if η is monotone nondecreasing and $\eta(0+) = 0$. Here and henceforth we use notations from Subsection 2.1.

Definition 1.1 (unweighted version). Let $z_0 = (x_0, t_0) \in Q_1$. We say that a function $u \in W_2^{1,1}(Q_1)$ satisfies the *unweighted almost parabolic Signorini property* at z_0 if $u \geq \varphi$ on Q'_1 and for any parabolic cylinder $Q_r(z_0) \Subset Q_1$ with $0 < r < r_0$, we have

$$\int_{Q_r(z_0)} (1 - \eta(r)) |\nabla u|^2 + 2\partial_t u(u - v) \leq (1 + \eta(r)) \int_{Q_r(z_0)} |\nabla v|^2$$

for any $v \in W_2^{1,0}(Q_r(z_0))$ with $v \geq \varphi$ on $Q'_r(z_0)$ and $v - u \in L^2(t_0 - r^2, t_0; W_0^{1,2}(B_r(x_0)))$.

We say that $u \in W_2^{1,1}(Q_1)$ is an *unweighted almost minimizer for the parabolic Signorini problem* in Q_1 if $u \geq \varphi$ on Q'_1 and u satisfies the almost parabolic Signorini property at every $z_0 \in Q_1$.

Next, we define the weighted version of almost minimizers. To this aim, we observe that if u is a solution of the parabolic Signorini problem in the strip S_1 , then

$$(1.1) \quad \int_{S_1} [(-t)|\nabla u|^2 + (-x \cdot \nabla u - 2t\partial_t u)(u - w)] G \, dxdt \leq \int_{S_1} (-t)|\nabla w|G \, dxdt$$

for every proper competitor w . This motivates the following definition of weighted almost minimizers.

Definition 1.2 (weighted version). Let $z_0 = (x_0, t_0) \in Q'_1$. We say that a function $u \in \mathcal{F}_{z_0}$ satisfies the *weighted almost parabolic Signorini property* at z_0 if $u \geq \varphi$ on S'_1 and

$$(1.2) \quad \begin{aligned} & \int_{S_r(t_0) \setminus S_\rho(t_0)} [(1 - \eta(r))(t_0 - t)|\nabla u|^2 \\ & \quad + (-x - x_0) \cdot \nabla u - 2(t - t_0)\partial_t u](u - w)] G_{z_0} \, dxdt \\ & \leq (1 + \eta(r)) \int_{S_r(t_0) \setminus S_\rho(t_0)} (t_0 - t)|\nabla w|^2 G_{z_0} \, dxdt \\ & \quad + \eta(r) \int_{S_r(t_0) \setminus S_\rho(t_0)} (u - w)^2 G_{z_0} \, dxdt + \|u\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}}. \end{aligned}$$

for any $0 \leq \rho < r < r_0$ with $-1 < t_0 - r^2$, and $w \in L^2(t_0 - r^2, t_0 - \rho^2; W^{1,2}(\mathbb{R}^n, G_{z_0}))$ with $w \geq \varphi$ on $S'_r(t_0) \setminus S'_\rho(t_0)$ and $u - w \in L^2(t_0 - r^2, t_0 - \rho^2; W_0^{1,2}(\mathbb{R}^n, G_{z_0}))$.

We say that a function $u \in \mathcal{F}$ is a *weighted almost minimizer for the parabolic Signori problem* on Q'_1 if $u \geq \varphi$ on S'_1 and u satisfies the weighted almost parabolic Signorini property at every $z_0 \in Q'_1$.

The readers might be intrigued by the presence of the exponential term $\|u\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}}$ in (1.2). We incorporated this term since we discovered that solutions of some

perturbed parabolic Signorini problems exhibit characteristics of almost minimizers, with the inclusion of the exponential error, see Appendix B.

Definition 1.3. We say that a function $u \in \mathcal{F}$ is an *almost minimizer for the parabolic Signorini problem* in Q_1 if it is both an unweighted almost minimizer in Q_1 and a weighted almost minimizer on Q'_1 .

The notion of a weighted almost minimizer is crucial for establishing monotonicity formulas, which are significant ingredients in our analysis of the free boundary. We will verify in Appendix B that solutions of some perturbed parabolic Signorini problems, multiplied by a standard cutoff function, satisfy the weighted almost parabolic Signorini property.

For background information and relevant literature concerning almost minimizers, we refer to [JP23] and references therein.

The time-independent almost minimizers for the Signorini problem were comprehensively treated by the authors in [JP21]. This paper extends specific results from the elliptic to the parabolic setting by employing similar energy methods. However, the parabolic case presents significant new challenges compared to the elliptic setting, primarily because we have to work with energy functionals involving singular weights.

1.3. Main results. This paper focuses on the local regularity results for free boundaries. Thus we assume that the domain $\Omega_T \subset \mathbb{R}^n \times \mathbb{R}$ is the parabolic cylinder Q_1 . Given the technical nature of the problem, we specifically examine the scenario where the thin space \mathcal{M}_T is Q'_1 (flat thin space), the thin obstacle φ is identically zero (zero thin obstacle), and the gauge function $\eta(r) = r^\alpha$ for some $0 < \alpha < 1$ with $r_0 = 1$.

Our first central result of this paper concerns the Weiss-type monotonicity formula.

Theorem A. Fix $\kappa_0 > 2$, $0 < \delta < 2$ and $0 < \varepsilon \leq \alpha < 1$. For $z_0 \in \Gamma(u) \cap Q'_{1/2}$, let $u \in \mathcal{F}_{z_0}$ satisfy the weighted almost parabolic Signorini property at z_0 . For $0 < \kappa < \kappa_0$, we set

$$W_{\kappa, \alpha, \varepsilon, \delta}(r, u, z_0) := \frac{e^{ar^\alpha}}{r^{2\kappa+2}} \left(\int_{S_r(t_0)} (2(t_0 - t)|\nabla u|^2 - \kappa(1 - br^\varepsilon)u^2) G_{z_0} dxdt + \|u\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}r^{-\delta}} \right),$$

where $a = a(\kappa, \alpha) > 0$ and $b = b(\kappa, \varepsilon) > 0$ are as in Theorem 3.3. Then $W_{\kappa, \alpha, \varepsilon, \delta}(r, u, z_0)$ is nondecreasing in r for $0 < r < r_0 = r_0(\kappa_0, \varepsilon)$.

Since almost minimizers do not satisfy a partial differential equations, we prove Theorem A by comparing them with appropriate homogeneous replacements, as done in the elliptic counterpart [JP21]. However, in our parabolic case, we encounter new technical challenges, making the proof significantly more complicated. This is mainly due to the formulation of the Weiss-type energy, which is defined in the unbounded strip and involves the singular weight. It turns out that we have to employ conformal self-similar coordinates.

By making use of the above one-parameter family of Weiss-type monotonicity formulas, we derive the Almgren-type frequency formula. For caloric functions, the

monotonicity of the following frequency was established in [Poo96]:

$$r \mapsto N(r, u, z_0) = \frac{r^2 \int_{\mathbb{R}^n \times \{t_0 - r^2\}} |\nabla u|^2 G_{z_0} dx}{\int_{\mathbb{R}^n \times \{t_0 - r^2\}} u^2 G_{z_0} dx}.$$

Recently, its averaged version was considered in [DGPT17] for the study of the parabolic Signorini problem. Regarding almost minimizers, we show that a modification of those quantities is monotone. To describe it, we denote

$$N_\delta(r, u, z_0) := \frac{\int_{S_r(t_0)} 2(t_0 - t) |\nabla u|^2 G_{z_0} + \|u\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}} r^{-\delta}}{\int_{S_r(t_0)} u^2 G_{z_0}}.$$

Theorem B (Almgren-type monotonicity formula). *Let $\kappa_0, \delta, \varepsilon, \alpha, z_0, u$, and b be as in Theorem A. Then $\widehat{N}_{\kappa_0, \varepsilon, \delta}(r, u, z_0) := \min\{\frac{1}{1-br^\varepsilon} N_\delta(r, u, z_0), \kappa_0\}$ is nondecreasing in $0 < r < r_0 = r_0(\kappa_0, \varepsilon)$. Moreover, if u is even-symmetric in x_n -variable, then we have either*

$$\widehat{N}_{\kappa_0, \delta}(0+, u, z_0)^1 = 3/2 \quad \text{or} \quad \widehat{N}_{\kappa_0, \delta}(0+, u, z_0) \geq 2.$$

In contrast to the Almgren frequency utilized for solutions to the parabolic Signorini problem [DGPT17], the Almgren-type frequencies we work with for almost minimizers include the extra exponential term $\|u\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}} r^{-\delta}$. Yet, we will show that this term is unsubstantial (see Lemma 4.5) and derive the same minimal frequency and frequency gap as presented in [DGPT17] (see Lemma 4.6).

Next, we consider the subset of the free boundary

$$\mathcal{R}(u) = \{z_0 \in \Gamma(u) \cap Q'_{1/2} : \widehat{N}_{\kappa_0, \delta}(0+, u, z_0) = 3/2 \text{ for some } \kappa_0 > 2, 0 < \delta < 2\},$$

the set of all free boundary points with the minimal frequency 3/2, known as the *regular set*.

Theorem C (Optimal growth near regular free boundary). *Fix $\kappa_0 > 2$. Suppose that an even-symmetric function $u \in \mathcal{F}_{z_0}$ satisfies the almost parabolic Signorini property at $z_0 \in \mathcal{R}(u)$. Then,*

$$\int_{S_r(t_0)} u^2 G_{z_0} dx dt \leq C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_{z_0}}^2 r^5,$$

for $0 < r < r_0 = r_0(\kappa_0, n, \alpha)$.

In the elliptic counterpart [JP21], an analogous result was derived using the epiperimetric inequality. Regarding the parabolic Signorini problem, Shi [Shi20] obtained a similar result by introducing the parabolic epiperimetric inequality. In our case, we adopt similar approaches. It is worth noting that while the application of these inequalities is rather immediate or standard in [Shi20, JP21], it is considerably more complicated for the parabolic almost minimizers (see Lemmas 5.3-5.4).

Finally, the main result concerning the regularity of the regular set is as follows.

Theorem D (Regularity of the regular set). *Let $u \in \mathcal{F}$ be a symmetric almost minimizer for the parabolic Signorini problem in Q_1 . Then $\mathcal{R}(u)$ can be represented locally as an $(n-2)$ -dimensional graph of a function, which has Hölder continuous spatial derivatives.*

¹From the monotonicity of $\widehat{N}_{\kappa_0, \varepsilon, \delta}$ and $\lim_{r \rightarrow 0} (1 - br^\varepsilon) = 1$, we see that the limit $\widehat{N}_{\kappa_0, \delta}(0+, u, z_0) = \lim_{r \rightarrow 0} \widehat{N}_{\kappa_0, \varepsilon, \delta}(r, u, z_0)$ exists and its value is independent of ε .

1.3.1. *Proofs of Theorems A–D.* Although we do not provide formal proofs of Theorems A–D in the main body of the paper, they can be deduced from the combination of results there. To be more precise,

- Theorem A is contained in Theorem 3.3.
- Theorem B follows by combining Theorem 3.4 and Lemma 4.6.
- The statement of Theorem C is contained in that of Lemma 5.4.
- The statement of Theorem D is contained in that of Theorem 7.8.

2. NOTATION AND PRELIMINARIES

2.1. **Notation.** We use the following notations throughout the paper.

For a function u , a set $\Omega \subset \mathbb{R}^{n+1}$, a constant $\varepsilon \in (0, 1)$, and a point $z_0 = (x_0, t_0)$, we denote

$$\begin{aligned} Q_r(z_0) &= B_r(x_0) \times (t_0 - r^2, t_0] \\ Q_{r,\rho}^\varepsilon(z_0) &= B_{r^\varepsilon}(x_0) \times (t_0 - r^2, t_0 - \rho^2] \\ \partial_p Q_r(z_0) &= (\partial B_r(x_0) \times [t_0 - r^2, t_0]) \cup (B_r(x_0) \times \{t_0 - r^2\}) : \text{parabolic boundary} \\ S_\rho(t_0) &= \mathbb{R}^n \times (t_0 - \rho^2, t_0] \\ \Omega' &= \Omega \cap \{x_n = 0\} \\ u_\Omega &= \int_\Omega u \\ u_{z_0,r} &= u_{Q_r(z_0)} = \int_{Q_r(z_0)} u \\ \|z_0\| &= (|x_0|^2 + |t_0|)^{1/2} : \text{parabolic norm} \\ \Gamma(u) &= \partial_{Q_1'} \{(x', t) \in Q_1' : u(x', 0, t) = 0\} : \text{free boundary} \end{aligned}$$

Given $l = k + \gamma$ with $k \in \mathbb{N} \cup \{0\}$ and $0 < \gamma \leq 1$, we use standard notations for parabolic Hölder spaces of functions $H^{l,l/2}$. For $1 \leq q \leq \infty$, we denote $W_q^{1,0}$ and $W_q^{1,1}$ by standard parabolic Sobolev spaces of functions. We refer to [DGPT17, JP23] for detailed definition.

We denote the backward heat kernel by

$$G(x, t) = \begin{cases} (-4\pi t)^{-n/2} e^{-\frac{|x|^2}{4t}}, & t < 0 \\ 0, & t \geq 0, \end{cases}$$

and write its translations

$$G_{z_0} = G(\cdot - x_0, \cdot - t_0).$$

Given $z_0 = (x_0, t_0) \in Q_1'$ and $0 < r < 1$, we let

$$\begin{aligned} \|u\|_{W_2^{1,0}(S_r(t_0), G_{z_0})} &:= \left[\int_{S_r(t_0)} (u^2 + (t_0 - t)|\nabla u|^2) G_{z_0} dxdt \right]^{1/2}, \\ \|u\|_{W_2^{1,1}(S_r(t_0), G_{z_0})} &:= \left[\int_{S_r(t_0)} (u^2 + (t_0 - t)(|\nabla u|^2 + (\partial_t u)^2)) G_{z_0} dxdt \right]^{1/2}. \end{aligned}$$

We say that $u \in \mathcal{F}_{z_0}$ if $u \in W_2^{1,1}(\mathbb{R}^n \times (-1, t_0), G_{z_0}) \cap W_2^{1,1}(B_1 \times (-1, t_0)) \cap L^\infty(\mathbb{R}^n \times (-1, t_0))$. We define the associated norm by

$$\|u\|_{\mathcal{F}_{z_0}} := \|u\|_{W_2^{1,1}(\mathbb{R}^n \times (-1, t_0), G_{z_0})} + \|u\|_{W_2^{1,1}(B_1 \times (-1, t_0))} + \|u\|_{L^\infty(\mathbb{R}^n \times (-1, t_0))}.$$

In addition, we say that $u \in \mathcal{F}$ if $u \in \mathcal{F}_{z_0}$ for every $z_0 \in Q_1'$ and

$$\|u\|_{\mathcal{F}} := \sup_{z_0 \in Q_1'} \|u\|_{W_2^{1,1}(\mathbb{R}^n \times (-1, t_0), G_{z_0})} + \|u\|_{W_2^{1,1}(Q_1)} + \|u\|_{L^\infty(S_1)} < \infty.$$

2.2. Preliminaries. The following regularity result for unweighted almost minimizers was proved in [JP23].

Theorem 2.1. *Let u be an unweighted almost minimizer for the parabolic Signorini problem in Q_1 . Then*

- (1) $u \in H^{\sigma, \sigma/2}(Q_1)$ for every $0 < \sigma < 1$;
- (2) $\nabla u \in H^{\beta, \beta/2}(Q_1^\pm \cup Q_1')$ for some $\beta = \beta(n, \alpha) > 0$.

Moreover, the authors showed in [JP23] that if u is an almost caloric function, then a stronger result than (2) in Theorem 2.1 holds:

$$\nabla u \in H^{\alpha/2, \alpha/4}(Q_1).$$

Here, an almost caloric function essentially is an unweighted almost minimizer without the obstacle condition; we refer to [JP23, Definitions 2.1-2.2] for its precise definition and [JP23, Theorem 2.8] for its regularity result.

By using Theorem 2.1 and the above Hölder continuity of spatial gradients of almost caloric functions across the thin space Q_1' , we can follow the argument in [JP21, Lemma 4.7] to derive the following complementarity condition.

Lemma 2.2 (Complementarity condition). *Let u be an unweighted almost minimizer for the parabolic Signorini problem in Q_1 , even in x_n -variable. Then u satisfies the following complementarity condition*

$$u \partial_{x_n}^+ u = 0 \quad \text{on } Q_1'.$$

In addition, we define

$$\widehat{\nabla} u(x', x_n, t) := \begin{cases} \nabla u(x', x_n, t), & x_n \geq 0, \\ \nabla u(x', -x_n, t), & x_n < 0, \end{cases}$$

the even extension of ∇u from Q_1^+ to Q_1 . If $z_0 \in \Gamma(u)$, then

$$u(z_0) = 0 \quad \text{and} \quad |\widehat{\nabla} u(z_0)| = 0.$$

3. WEISS- AND ALMGREN-TYPE MONOTONICITY FORMULAS

The purpose of this section is to establish monotonicity formulas of Weiss- and Almgren-type. They will play a crucial role in the analysis of the free boundary.

We first prove the Weiss-type monotonicity formula, which represents one of the most technical aspects of this paper. In its elliptic counterpart [JP21], the authors derived the formula by comparing almost minimizers and homogeneous replacements, inspired by the approach in [Wei99]. In the current parabolic case, we compare almost minimizers and parabolically homogeneous replacements, and utilize conformal self-similar coordinates. For its proof, we need the following auxiliary results.

Lemma 3.1. *Fix $\kappa_0 > 2$ and $0 < \varepsilon \leq \alpha < 1$. For $0 < \kappa < \kappa_0$ and $0 \leq \rho < r$, let*

$$\Phi(r) := \Phi_{\rho, \kappa, \alpha}(r) = \frac{e^{ar^\alpha}}{r^{2\kappa+2} - \rho^{2\kappa+2}}, \quad \Psi(r) := \Psi_{\rho, \kappa, \kappa_0, \alpha, \varepsilon}(r) = \frac{(1 - br^\varepsilon)e^{ar^\alpha}}{r^{2\kappa+2} - \rho^{2\kappa+2}}$$

with

$$a = \frac{8(\kappa + 1)}{\alpha}, \quad b = \frac{128(\kappa_0 + 1)}{\varepsilon}.$$

Then, there is a small constant $r_0 = r_0(\kappa_0, \varepsilon) = \frac{r_0(\varepsilon)}{\kappa_0^{2/\varepsilon}} > 0$ such that for $0 \leq \rho < r < r_0$ with $\rho/r \leq 1/\sqrt{2}$,

$$(3.1) \quad \Phi'(r) \leq 0,$$

$$(3.2) \quad \frac{\Phi'(r)}{1 - r^\alpha} - \Psi'(r) \geq -\frac{(2\kappa + 2 - \varepsilon/4)b\Phi(r)r^{2\kappa+1+\varepsilon}}{r^{2\kappa+2} - \rho^{2\kappa+2}},$$

$$(3.3) \quad \frac{1 + r^\alpha}{1 - r^\alpha}\Phi'(r) + \frac{2(\kappa + 1)r^{2\kappa+1}}{r^{2\kappa+2} - \rho^{2\kappa+2}}\Phi(r) \geq 0,$$

$$(3.4) \quad -\frac{\Phi'(r)}{1 - r^\alpha} - \frac{2(\kappa + 1)r^{2\kappa+1}}{r^{2\kappa+2} - \rho^{2\kappa+2}}\Psi(r) \geq \frac{(2\kappa + 2 - \varepsilon/8)b\Phi(r)r^{2\kappa+1+\varepsilon}}{r^{2\kappa+2} - \rho^{2\kappa+2}}.$$

Proof. We first prove (3.1). By using $0 < \varepsilon < \alpha$, we simply compute

$$\begin{aligned} \Phi'(r) &= \left(a\alpha r^\alpha - \frac{(2\kappa + 2)r^{2\kappa+2}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \right) \frac{\Phi(r)}{r} \\ &\leq ((8\kappa + 8)r^\alpha - (2\kappa + 2)) \frac{\Phi(r)}{r} \leq 0, \quad r < r_0(\varepsilon). \end{aligned}$$

For (3.2), we note $r^{2\kappa+2} - \rho^{2\kappa+2} \geq (1 - (\rho/r)^2)r^{2\kappa+2} \geq \frac{1}{2}r^{2\kappa+2}$ and get

$$\begin{aligned} (r^{2\kappa+2} - \rho^{2\kappa+2})\Phi'(r) &= ((r^{2\kappa+2} - \rho^{2\kappa+2})a\alpha r^\alpha - (2\kappa + 2)r^{2\kappa+2}) \frac{\Phi(r)}{r} \\ &\geq (1/2a\alpha r^\alpha - (2\kappa + 2))\Phi(r)r^{2\kappa+1}. \end{aligned}$$

Moreover, using $r^{2\kappa+2} - \rho^{2\kappa+2} \geq \frac{1}{2}r^{2\kappa+2}$ again along with $b\varepsilon r^\varepsilon \geq a\alpha r^\alpha \geq (1 - br^\varepsilon)a\alpha r^\alpha$, we find

$$\begin{aligned} &(r^{2\kappa+2} - \rho^{2\kappa+2})\Psi'(r) \\ &= (r^{2\kappa+2} - \rho^{2\kappa+2})(-b\varepsilon r^{\varepsilon-1}\Phi(r) + (1 - br^\varepsilon)\Phi'(r)) \\ &= ((r^{2\kappa+2} - \rho^{2\kappa+2})(-b\varepsilon r^\varepsilon + (1 - br^\varepsilon)a\alpha r^\alpha) - (1 - br^\varepsilon)(2\kappa + 2)r^{2\kappa+2}) \frac{\Phi(r)}{r} \\ &\leq (1/2(-b\varepsilon r^\varepsilon + (1 - br^\varepsilon)a\alpha r^\alpha) - (1 - br^\varepsilon)(2\kappa + 2))\Phi(r)r^{2\kappa+1}. \end{aligned}$$

Thus, we have

$$\begin{aligned} &(r^{2\kappa+2} - \rho^{2\kappa+2}) \left(\frac{\Phi'(r)}{1 - r^\alpha} - \Psi'(r) \right) \\ &\geq \left(1/2a\alpha r^\alpha - (2\kappa + 2) + \frac{1/2a\alpha}{1 - r^\alpha} r^{2\alpha} - \frac{2\kappa + 2}{1 - r^\alpha} r^\alpha \right. \\ &\quad \left. + 1/2b\varepsilon r^\varepsilon - 1/2(1 - br^\varepsilon)a\alpha r^\alpha + (2\kappa + 2) - (2\kappa + 2)br^\varepsilon \right) \Phi(r)r^{2\kappa+1} \\ &\geq - \left((2\kappa + 2 - \varepsilon/2)br^\varepsilon + \frac{2\kappa + 2}{1 - r^\alpha} r^\alpha \right) \Phi(r)r^{2\kappa+1} \end{aligned}$$

$$\geq -(2\kappa + 2 - \varepsilon/4) br^\varepsilon \Phi(r) r^{2\kappa+1},$$

where the last inequality follows from $\frac{2\kappa+2}{1-r^\alpha} \leq 16(\kappa+1) \leq b\varepsilon/8$ and $r^\alpha \leq r^\varepsilon$. Regarding (3.3), we use $r^{2\kappa+2} - \rho^{2\kappa+2} \geq \frac{1}{2}r^{2\kappa+2}$ once again to obtain

$$\begin{aligned} & \frac{1+r^\alpha}{1-r^\alpha} \Phi'(r) + \frac{2(\kappa+1)r^{2\kappa+1}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \Phi(r) \\ &= \left(\frac{1+r^\alpha}{1-r^\alpha} \left(a\alpha r^\alpha - \frac{2(\kappa+1)r^{2\kappa+2}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \right) + \frac{2(\kappa+1)r^{2\kappa+2}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \right) \frac{\Phi(r)}{r} \\ &= \left((1+r^\alpha)a\alpha r^\alpha - \frac{4(\kappa+1)r^{2\kappa+2+\alpha}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \right) \frac{\Phi(r)}{(1-r^\alpha)r} \\ &\geq ((1+r^\alpha)a\alpha - 8(\kappa+1)) \frac{\Phi(r)}{(1-r^\alpha)r^{1-\alpha}} \geq 0. \end{aligned}$$

Finally, we prove (3.4).

$$\begin{aligned} & (r^{2\kappa+2} - \rho^{2\kappa+2}) \left(-\frac{\Phi'(r)}{1-r^\alpha} - \frac{2(\kappa+1)r^{2\kappa+1}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \Psi(r) \right) \\ &= \left(-\frac{a\alpha}{1-r^\alpha} (r^{2\kappa+2} - \rho^{2\kappa+2}) r^\alpha + \frac{2\kappa+2}{1-r^\alpha} r^{2\kappa+2} - (1-br^\varepsilon)(2\kappa+2)r^{2\kappa+2} \right) \frac{\Phi(r)}{r} \\ &\geq \left(-\frac{a\alpha}{1-r^\alpha} r^\alpha + (2\kappa+2)br^\varepsilon \right) \Phi(r) r^{2\kappa+1} \geq (2\kappa+2 - \varepsilon/8) br^\varepsilon \Phi(r) r^{2\kappa+1}, \end{aligned}$$

where the last step follows from $\frac{a\alpha}{1-r^\alpha} \leq 16(\kappa+1) \leq b\varepsilon/8$ and $r^\alpha \leq r^\varepsilon$. \square

As previously mentioned, we will make use of conformal self-similar coordinates. Given constants $0 < r < 1$ and $\kappa > 0$ and a function u defined in S_r , we define

$$(3.5) \quad \tilde{u}(y, \tau) = \tilde{u}_\kappa(y, \tau) := e^{\kappa\tau/2} u \left(2e^{-\tau/2} y, -e^{-\tau} \right), \quad (y, \tau) \in \mathbb{R}^n \times (-2 \ln r, \infty).$$

In addition, we let

$$(3.6) \quad w(x, t) := \left(\frac{\sqrt{-t}}{r} \right)^\kappa u \left(\frac{r}{\sqrt{-t}} x, -r^2 \right), \quad (x, t) \in S_r$$

be the parabolically κ -homogeneous replacement of u in S_r . From its construction, it is easily seen that w satisfies the homogeneity

$$(3.7) \quad \kappa w - x \cdot \nabla w - 2t \partial_t w = 0.$$

Then, $\tilde{w}(y, \tau) := e^{\kappa\tau/2} w(2e^{-\tau/2} y, -e^{-\tau})$ satisfies

$$(3.8) \quad \partial_\tau \tilde{w}(y, \tau) = 0 \quad \text{for } (y, \tau) \in \mathbb{R}^n \times (-2 \ln r, \infty),$$

which implies that $\tilde{w}(y) = \tilde{w}(y, \tau)$ is independent of τ -variable. This, along with the fact that $w(x, -r^2) = u(x, -r^2)$ for $x \in \mathbb{R}^n$, yields

$$(3.9) \quad \tilde{w}(y) = \tilde{u}(y, -2 \ln r), \quad y \in \mathbb{R}^n.$$

Lemma 3.2. *Let $u \in \mathcal{F}_0$. Then, for $\kappa > 0$ and $0 \leq \rho < r < 1$,*

$$(3.10) \quad \begin{aligned} & \int_{S_r \setminus S_\rho} (\kappa u - x \cdot \nabla u - 2t \partial_t u)(u - w) G \\ &= \frac{\rho^{2\kappa+2}}{\pi^{n/2}} \int_{\mathbb{R}^n} (\tilde{u}(y, -2 \ln \rho) - \tilde{u}(y, -2 \ln r))^2 e^{-|y|^2} dy + (\kappa + 1) \int_{S_r \setminus S_\rho} (u - w)^2 G, \end{aligned}$$

where \tilde{u} and w are as in (3.5) and (3.6), respectively. In particular,

$$(3.11) \quad \int_{S_r} (\kappa u - x \cdot \nabla u - 2t \partial_t u)(u - w)G = (\kappa + 1) \int_{S_r} (u - w)^2 G.$$

Proof. By using (3.7) and (3.9), we obtain (3.10):

$$\begin{aligned} & \int_{S_r \setminus S_\rho} (\kappa u - x \cdot \nabla u - 2t \partial_t u)(u - w)G \, dxdt \\ &= \int_{S_r \setminus S_\rho} (\kappa(u - w) - x \cdot \nabla(u - w) - 2t \partial_t(u - w))(u - w)G \, dxdt \\ &= \frac{1}{2} \int_{S_r \setminus S_\rho} (2\kappa(u - w)^2 - x \cdot \nabla((u - w)^2) - 2t \partial_t((u - w)^2))G \, dxdt \\ &= \frac{1}{\pi^{n/2}} \int_{\mathbb{R}^n \times (-2 \ln r, -2 \ln \rho)} \partial_\tau((\tilde{u} - \tilde{w})^2) e^{-|y|^2} e^{-(\kappa+1)\tau} \, dyd\tau \\ &= \frac{1}{\pi^{n/2}} \int_{\mathbb{R}^n \times \{-2 \ln \rho\}} (\tilde{u} - \tilde{w})^2 e^{-|y|^2} \rho^{2\kappa+2} \, dy \\ &\quad + \frac{\kappa + 1}{\pi^{n/2}} \int_{\mathbb{R}^n \times (-2 \ln r, -2 \ln \rho)} (\tilde{u} - \tilde{w})^2 e^{-|y|^2} e^{-(\kappa+1)\tau} \, dyd\tau \\ &= \frac{\rho^{2\kappa+2}}{\pi^{n/2}} \int_{\mathbb{R}^n} (\tilde{u}(y, -2 \ln \rho) - \tilde{u}(y, -2 \ln r))^2 e^{-|y|^2} \, dy + (\kappa + 1) \int_{S_r \setminus S_\rho} (u - w)^2 G. \end{aligned}$$

Moreover, (3.11) follows from (3.10) by taking $\rho \rightarrow 0$ with the observation

$$\lim_{\rho \rightarrow 0} \rho^{2\kappa+2} (\tilde{u}(y, -2 \ln \rho) - \tilde{u}(y, -2 \ln r))^2 \leq \lim_{\rho \rightarrow 0} \rho^{2\kappa+2} ((\rho^{-\kappa} + r^{-\kappa}) \|u\|_{L^\infty(S_r)})^2 = 0.$$

□

We now prove the Weiss-type monotonicity formula with the help of Lemmas 3.1 and 3.2. We note that for any $\kappa > 0$, the weighted almost parabolic Signorini property (1.2) is equivalent to

$$(3.12) \quad \begin{aligned} & \int_{S_r(t_0) \setminus S_\rho(t_0)} [2(1 - \eta(r))(t_0 - t)|\nabla u|^2 - \kappa u^2 \\ & \quad + 2(\kappa u - (x - x_0) \cdot \nabla u - 2(t - t_0) \partial_t u)(u - w)] G_{z_0} \, dxdt \\ & \leq \int_{S_r(t_0) \setminus S_\rho(t_0)} [2(1 + \eta(r))(t_0 - t)|\nabla w|^2 - \kappa w^2 \\ & \quad + (\kappa + 2\eta(r))(u - w)^2] G_{z_0} \, dxdt + 2\|u\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}}. \end{aligned}$$

Theorem 3.3 (Weiss-type monotonicity formula). *Fix $\kappa_0 > 2$, $0 < \delta < 2$ and $0 < \varepsilon \leq \alpha < 1$. Suppose that for $z_0 = (x_0, t_0) \in Q'_{1/2}$, $u \in \mathcal{F}_{z_0}$ satisfies the weighted almost parabolic Signorini property at z_0 . For $0 < \kappa < \kappa_0$, set*

$$\begin{aligned} & W_{\kappa, \alpha, \varepsilon, \delta, \rho}(r, u, z_0) \\ & := \frac{e^{ar^\alpha}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \left(\int_{S_r(t_0) \setminus S_\rho(t_0)} (2(t_0 - t)|\nabla u|^2 - \kappa(1 - br^\varepsilon)u^2) G_{z_0} \, dxdt \right. \\ & \quad \left. + \|u\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}} r^{-\delta} \right), \end{aligned}$$

where constants a, b are as in Lemma 3.1

$$a = \frac{8(\kappa + 1)}{\alpha}, \quad b = \frac{128(\kappa_0 + 1)}{\varepsilon}.$$

(i) For $0 < \rho < r < r_0 = r_0(\kappa_0, \varepsilon) = \frac{r_0(\varepsilon)}{\kappa_0^{2/\varepsilon}}$ with $\rho/r \leq 1/\sqrt{2}$,

$$(3.13) \quad \begin{aligned} & \frac{d}{dr} W_{\kappa, \alpha, \varepsilon, \delta, \rho}(r, u, z_0) \\ & \geq \frac{(4\kappa + 2)r^{2\kappa+1}\rho^{2\kappa+2}}{\pi^{n/2}(r^{2\kappa+2} - \rho^{2\kappa+2})^2} \int_{\mathbb{R}^n} (\tilde{u}(y, -2\ln \rho) - \tilde{u}(y, -2\ln r))^2 e^{-|y|^2} dy, \end{aligned}$$

where $\tilde{u} = \tilde{u}_\kappa$ is as in (3.5).

(ii) When $\rho = 0$, for $W_{\kappa, \alpha, \varepsilon, \delta} = W_{\kappa, \alpha, \varepsilon, \delta, 0}$ and $0 < r < r_0 = r_0(\kappa_0, \alpha, \varepsilon) = \frac{r_0(\alpha, \varepsilon)}{\kappa_0^{2/\varepsilon}}$,

$$(3.14) \quad \begin{aligned} & \frac{d}{dr} W_{\kappa, \alpha, \varepsilon, \delta}(r, u, z_0) \\ & \geq \frac{\kappa}{2r^{2\kappa+3-\varepsilon/2}} \left| \int_{S_r} (\kappa u - (x - x_0) \cdot \nabla u - 2(t - t_0)\partial_t u) u G_{z_0} \right|. \end{aligned}$$

Although we work with $W_{\kappa, \alpha, \varepsilon, \delta}$ throughout most of this paper, the monotonicity of $W_{\kappa, \alpha, \varepsilon, \delta, \rho}$ will be used when we establish the rotation estimate in Lemma 6.1.

Proof. The proof is divided into several steps.

Step 1. Without loss of generality, we assume $z_0 = 0$. We write for simplicity $W_{\kappa, \rho} = W_{\kappa, \alpha, \varepsilon, \delta, \rho}$. Let w be the homogeneous replacement as in (3.6). Note that we can write

$$W_{\kappa, \rho}(r, u) = \Phi(r) \int_{S_r \setminus S_\rho} (-2t)|\nabla u|^2 G - \Psi(r) \int_{S_r \setminus S_\rho} \kappa u^2 G + \Phi(r) \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}} r^{-\delta},$$

where $\Phi(r) = \frac{e^{ar\alpha}}{r^{2\kappa+2} - \rho^{2\kappa+2}}$ and $\Psi(r) = \frac{(1-br^\varepsilon)e^{ar\alpha}}{r^{2\kappa+2} - \rho^{2\kappa+2}}$ are as in Lemma 3.1. Then, by using (3.12) and (3.1), we deduce

$$\begin{aligned} & \frac{d}{dr} W_{\kappa, \rho}(r, u) \\ & = \Phi'(r) \int_{S_r \setminus S_\rho} (-2t)|\nabla u|^2 G - \Psi'(r) \int_{S_r \setminus S_\rho} \kappa u^2 G + 2r\Phi(r) \int_{\mathbb{R}^n \times \{-r^2\}} 2r^2 |\nabla u|^2 G \\ & \quad - 2r\Psi(r) \int_{\mathbb{R}^n \times \{-r^2\}} \kappa u^2 G + \|u\|_{\mathcal{F}_0}^2 \frac{d}{dr} \left(\Phi(r) e^{-\frac{1}{r}} r^{-\delta} \right) \\ & = \frac{\Phi'(r)}{1-r^\alpha} \int_{S_r \setminus S_\rho} ((1-r^\alpha)(-2t)|\nabla u|^2 - \kappa u^2) G + \left(\frac{\Phi'(r)}{1-r^\alpha} - \Psi'(r) \right) \int_{S_r \setminus S_\rho} \kappa u^2 G \\ & \quad + 2r\Phi(r) \int_{\mathbb{R}^n \times \{-r^2\}} 2r^2 |\nabla u|^2 G - 2r\Psi(r) \int_{\mathbb{R}^n \times \{-r^2\}} \kappa u^2 G \\ & \quad + \|u\|_{\mathcal{F}_0}^2 \frac{d}{dr} \left(\Phi(r) e^{-\frac{1}{r}} r^{-\delta} \right) \\ & \geq \frac{\Phi'(r)}{1-r^\alpha} \left(\int_{S_r \setminus S_\rho} [(1+r^\alpha)(-2t)|\nabla w|^2 - \kappa w^2 - 2(\kappa u - x \cdot \nabla u - 2t\partial_t u)(u-w) \right. \\ & \quad \left. + (\kappa + 2r^\alpha)(u-w)^2] G + 2\|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}} \right) \end{aligned}$$

$$\begin{aligned}
& + \left(\frac{\Phi'(r)}{1-r^\alpha} - \Psi'(r) \right) \int_{S_r \setminus S_\rho} \kappa u^2 G + 2r\Phi(r) \int_{\mathbb{R}^n \times \{-r^2\}} 2r^2 |\nabla u|^2 G \\
& - 2r\Psi(r) \int_{\mathbb{R}^n \times \{-r^2\}} \kappa u^2 G + \|u\|_{\mathcal{F}_0}^2 \frac{d}{dr} \left(\Phi(r) e^{-\frac{1}{r}} r^{-\delta} \right) \\
& = I + II + III + IV + V,
\end{aligned}$$

where

$$\begin{aligned}
I & = \frac{\Phi'(r)(1+r^\alpha)}{1-r^\alpha} \int_{S_r \setminus S_\rho} (-2t) |\nabla w|^2 G + 2r\Phi(r) \int_{\mathbb{R}^n \times \{-r^2\}} 2r^2 |\nabla u|^2 G, \\
II & = -\frac{\Phi'(r)}{1-r^\alpha} \int_{S_r \setminus S_\rho} \kappa w^2 G - 2r\Psi(r) \int_{\mathbb{R}^n \times \{-r^2\}} \kappa u^2 G, \\
III & = \frac{\Phi'(r)}{1-r^\alpha} \int_{S_r \setminus S_\rho} \left[-2(\kappa u - x \cdot \nabla u - 2t\partial_t u)(u-w) + (\kappa + 2r^\alpha)(u-w)^2 \right] G, \\
IV & = \left(\frac{\Phi'(r)}{1-r^\alpha} - \Psi'(r) \right) \int_{S_r \setminus S_\rho} \kappa u^2 G, \\
V & = \frac{2\Phi'(r)\|u\|^2 e^{-\frac{1}{r}}}{1-r^\alpha} + \|u\|_{\mathcal{F}_0}^2 \frac{d}{dr} \left(\Phi(r) e^{-\frac{1}{r}} r^{-\delta} \right).
\end{aligned}$$

Step 2. In this step, we estimate the terms I - V . We begin with I and II . By using the homogeneity of w , we can directly compute

$$\begin{aligned}
\int_{S_r \setminus S_\rho} (-2t) |\nabla w|^2 G & = \frac{r^{2\kappa+2} - \rho^{2\kappa+2}}{(\kappa+1)r^{2\kappa}} \int_{\mathbb{R}^n \times \{-r^2\}} 2r^2 |\nabla u|^2 G \, dx, \\
\int_{S_r \setminus S_\rho} w^2 G & = \frac{r^{2\kappa+2} - \rho^{2\kappa+2}}{(\kappa+1)r^{2\kappa}} \int_{\mathbb{R}^n \times \{-r^2\}} u^2 G \, dx.
\end{aligned}$$

Combining these equalities with (3.3) and (3.4), we obtain

$$I = \left(\frac{\Phi'(r)(1+r^\alpha)}{1-r^\alpha} + 2r\Phi(r) \frac{(\kappa+1)r^{2\kappa}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \right) \int_{S_r \setminus S_\rho} (-2t) |\nabla w|^2 G \geq 0,$$

and

$$\begin{aligned}
II & = \left(-\frac{\Phi'(r)}{1-r^\alpha} - 2r\Psi(r) \frac{(\kappa+1)r^{2\kappa}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \right) \int_{S_r \setminus S_\rho} \kappa w^2 G \\
& \geq \frac{(2\kappa+2-\varepsilon/8)b\Phi(r)r^{2\kappa+1+\varepsilon}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \int_{S_r \setminus S_\rho} \kappa w^2 G.
\end{aligned}$$

Next, we estimate III . Note that

$$\begin{aligned}
\Phi'(r) & = (a\alpha r^\alpha (r^{2\kappa+2} - \rho^{2\kappa+2}) - (2\kappa+2)r^{2\kappa+2}) \frac{\Phi(r)}{r(r^{2\kappa+2} - \rho^{2\kappa+2})} \\
& \leq (a\alpha r^\alpha - (2\kappa+2)) \frac{\Phi(r)r^{2\kappa+1}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \\
& \leq -(2\kappa+1) \frac{\Phi(r)r^{2\kappa+1}}{r^{2\kappa+2} - \rho^{2\kappa+2}}, \quad r < \frac{1}{(16\kappa_0)^{1/\alpha}}.
\end{aligned}$$

This, along with (3.10), produces

$$III = -\frac{\Phi'(r)}{1-r^\alpha} \left(\frac{2\rho^{2\kappa+2}}{\pi^{n/2}} \int_{\mathbb{R}^n} (\tilde{u}(y, -2\ln \rho) - \tilde{u}(y, -2\ln r))^2 e^{-|y|^2} dy \right)$$

$$\begin{aligned}
 & + (\kappa + 2 - 2r^\alpha) \int_{S_r \setminus S_\rho} (u - w)^2 G \\
 \geq & \frac{(4\kappa + 2)\Phi(r)r^{2\kappa+1}\rho^{2\kappa+2}}{\pi^{n/2}(r^{2\kappa+2} - \rho^{2\kappa+2})} \int_{\mathbb{R}^n} (\tilde{u}(y, -2\ln \rho) - \tilde{u}(y, -2\ln r))^2 e^{-|y|^2} dy \\
 & + \frac{(2\kappa + 1)(\kappa + 1)\Phi(r)r^{2\kappa+1}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \int_{S_r \setminus S_\rho} (u - w)^2 G.
 \end{aligned}$$

Regarding IV , we simply use (3.2) to get

$$IV \geq -\frac{(2\kappa + 2 - \varepsilon/4) b\Phi(r)r^{2\kappa+1+\varepsilon}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \int_{S_r \setminus S_\rho} \kappa u^2 G.$$

Finally, to deal with V , we recall the inequality $r^{2\kappa+2} - \rho^{2\kappa+2} \geq \frac{1}{2}r^{2\kappa+2}$ to get

$$\Phi'(r) = \left(a\alpha r^\alpha - \frac{(2\kappa + 2)r^{2\kappa+2}}{r^{2\kappa+2} - \rho^{2\kappa+2}} \right) \frac{\Phi(r)}{r} \geq (a\alpha r^\alpha - 4(\kappa + 1)) \frac{\Phi(r)}{r},$$

which yields

$$\begin{aligned}
 \frac{V}{\|u\|_{\mathcal{F}_0}^2} &= \frac{2e^{-\frac{1}{r}}}{1 - r^\alpha} \Phi'(r) + \Phi'(r)e^{-\frac{1}{r}r^{-\delta}} + \left(\frac{1}{r} - \delta \right) \frac{\Phi(r)}{r} e^{-\frac{1}{r}r^{-\delta}} \\
 &\geq \left(\left(\frac{2}{1 - r^\alpha} + r^{-\delta} \right) (a\alpha r^\alpha - 4(\kappa + 1)) + \left(\frac{1}{r} - \delta \right) r^{-\delta} \right) \frac{\Phi(r)}{r} e^{-\frac{1}{r}} \\
 &\geq 0, \quad 0 < r < \frac{r_0(\varepsilon)}{\kappa_0}.
 \end{aligned}$$

Step 3. By combining the results in Step 1 and Step 2, we get

$$\begin{aligned}
 & \frac{r^{2\kappa+2} - \rho^{2\kappa+2}}{\Phi(r)r^{2\kappa+1}} \frac{d}{dr} W_{\kappa,\rho}(r, u) \\
 & \geq (2\kappa + 2 - \varepsilon/8) \kappa b r^\varepsilon \int_{S_r \setminus S_\rho} w^2 G + (2\kappa + 1)(\kappa + 1) \int_{S_r \setminus S_\rho} (u - w)^2 G \\
 & \quad - (2\kappa + 2 - \varepsilon/4) \kappa b r^\varepsilon \int_{S_r \setminus S_\rho} u^2 G \\
 & \quad + \frac{(4\kappa + 2)\rho^{2\kappa+2}}{\pi^{n/2}} \int_{\mathbb{R}^n} (\tilde{u}(y, -2\ln \rho) - \tilde{u}(y, -2\ln r))^2 e^{-|y|^2} dy.
 \end{aligned}$$

On the other hand, we take $\mu = \frac{\varepsilon}{12(2\kappa+2-\varepsilon/4)}$, which is tailor-made to satisfy $1 + \mu = \frac{2\kappa+2-\varepsilon/6}{2\kappa+2-\varepsilon/4}$, and apply Young's inequality to have that for $0 \leq \rho < r < \frac{r_0(\alpha, \varepsilon)}{\kappa_0^{2/\varepsilon}}$,

$$\begin{aligned}
 & (2\kappa + 2 - \varepsilon/8) \kappa b r^\varepsilon \int_{S_r \setminus S_\rho} w^2 G + (2\kappa + 1)(\kappa + 1) \int_{S_r \setminus S_\rho} (u - w)^2 G \\
 & \quad - (2\kappa + 2 - \varepsilon/4) \kappa b r^\varepsilon \int_{S_r \setminus S_\rho} u^2 G \\
 & \geq (2\kappa + 2 - \varepsilon/8) \kappa b r^\varepsilon \int_{S_r \setminus S_\rho} w^2 G + (2\kappa + 1)(\kappa + 1) \int_{S_r \setminus S_\rho} (u - w)^2 G \\
 & \quad - (2\kappa + 2 - \varepsilon/4) \kappa b r^\varepsilon \left((1 + \mu) \int_{S_r \setminus S_\rho} w^2 G + (1 + 1/\mu) \int_{S_r \setminus S_\rho} (u - w)^2 G \right)
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{24} \varepsilon \kappa b r^\varepsilon \int_{S_r \setminus S_\rho} w^2 G \\
&\quad + ((2\kappa + 1)(\kappa + 1) - (2\kappa + 2 - \varepsilon/4) \kappa b r^\varepsilon (1 + 1/\mu)) \int_{S_r \setminus S_\rho} (u - w)^2 G \\
&\geq 5\kappa(\kappa + 1)r^\varepsilon \int_{S_r \setminus S_\rho} w^2 G + \kappa(\kappa + 1) \int_{S_r \setminus S_\rho} (u - w)^2 G.
\end{aligned}$$

By combining the precious two inequalities, we deduce

$$\begin{aligned}
(3.15) \quad &\frac{r^{2\kappa+2} - \rho^{2\kappa+2}}{\Phi(r)r^{2\kappa+1}} \frac{d}{dr} W_{\kappa,\rho}(r, u) \\
&\geq 5\kappa(\kappa + 1)r^\varepsilon \int_{S_r \setminus S_\rho} w^2 G + \kappa(\kappa + 1) \int_{S_r \setminus S_\rho} (u - w)^2 G \\
&\quad + \frac{(4\kappa + 2)\rho^{2\kappa+2}}{\pi^{n/2}} \int_{\mathbb{R}^n} (\tilde{u}(y, -2 \ln \rho) - \tilde{u}(y, -2 \ln r))^2 e^{-|y|^2} dy.
\end{aligned}$$

This gives (3.13).

Step 4. The purpose of this step is to obtain (3.14). To this aim, we let $\rho = 0$, and observe that $(\tilde{u} - \tilde{w})\tilde{u} = 0$ on $\mathbb{R}^n \times \{-2 \ln r\}$ and that for any $y \in \mathbb{R}^n$

$$\lim_{\tau \rightarrow \infty} \left| (\tilde{u}(y, \tau) - \tilde{w}(y, \tau))\tilde{u}(y, \tau) e^{-(\kappa+1)\tau} \right| \leq \lim_{\tau \rightarrow \infty} \left(2\|u\|_{L^\infty(S_1)}^2 e^{-\tau} \right) = 0.$$

It then follows that

$$\begin{aligned}
&\int_{S_r} (\kappa u - x \cdot \nabla u - 2t \partial_t u) u G \\
&= \frac{2}{\pi^{n/2}} \int_{\mathbb{R}^n \times (-2 \ln r, \infty)} (\partial_\tau (\tilde{u} - \tilde{w})) \tilde{u} e^{-|y|^2} e^{-(\kappa+1)\tau} dy d\tau \\
&= -\frac{2}{\pi^{n/2}} \int_{\mathbb{R}^n \times (-2 \ln r, \infty)} (\tilde{u} - \tilde{w}) (\partial_\tau \tilde{u}) e^{-|y|^2} e^{-(\kappa+1)\tau} dy d\tau \\
&\quad + \frac{2(\kappa + 1)}{\pi^{n/2}} \int_{\mathbb{R}^n \times (-2 \ln r, \infty)} (\tilde{u} - \tilde{w}) \tilde{u} e^{-|y|^2} e^{-(\kappa+1)\tau} dy d\tau \\
&= -\int_{S_r} (u - w) (\kappa u - x \cdot \nabla u - 2t \partial_t u) G dx dt + 2(\kappa + 1) \int_{S_r} (u - w) u G dx dt \\
&= (\kappa + 1) \int_{S_r} (u - w)^2 G + 2(\kappa + 1) \int_{S_r} (u - w) w G,
\end{aligned}$$

where we used (3.8) in the first step and (3.11) in the last equality. Thus

$$\left| \int_{S_r} (\kappa u - x \cdot \nabla u - 2t \partial_t u) u G \right| \leq 2(\kappa + 1)r^{-\varepsilon/2} \int_{S_r} (u - w)^2 G + (\kappa + 1)r^{\varepsilon/2} \int_{S_r} w^2 G.$$

Therefore, by combining this with (3.15), we conclude that

$$\begin{aligned}
\frac{\kappa}{2} r^{\varepsilon/2} \left| \int_{S_r} (\kappa u - x \cdot \nabla u - 2t \partial_t u) u G \right| &\leq \kappa(\kappa + 1) \int_{S_r} (u - w)^2 G + \frac{\kappa(\kappa + 1)}{2} r^\varepsilon \int_{S_r} w^2 G \\
&\leq \frac{r}{\Phi(r)} \frac{d}{dr} W_\kappa(r, u).
\end{aligned}$$

This implies (3.14). \square

Next, we deal with the Almgren-type frequency in the parabolic setting. Poon proved in [Poo96] that if u is a caloric function in S_1 (i.e., $\Delta u - \partial_t u = 0$ in S_1), then its caloric frequency

$$N(r, u, z_0) := \frac{r^2 \int_{\mathbb{R}^n \times \{t_0 - r^2\}} |\nabla u|^2 G_{z_0} dx}{\int_{\mathbb{R}^n \times \{t_0 - r^2\}} u^2 G_{z_0} dx}$$

is monotone nondecreasing in $r \in (0, 1)$. Concerning the parabolic Signorini problem, [DGPT17] considered its averaged version

$$N^0(r, u, z_0) := \frac{\int_{S_r(t_0)} 2(t_0 - t) |\nabla u|^2 G_{z_0}}{\int_{S_r(t_0)} u^2 G_{z_0}},$$

and proved the generalized frequency formula related to N^0 when z_0 is a free boundary point. For almost minimizers, we need some modifications on N^0 . Given free boundary point $z_0 \in \Gamma(u) \cap Q'_{1/2}$, we let

$$N_\delta(r, u, z_0) := \frac{\int_{S_r(t_0)} 2(t_0 - t) |\nabla u|^2 G_{z_0} + \|u\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}} r^{-\delta}}{\int_{S_r(t_0)} u^2 G_{z_0}}.$$

We then define the multiplicative modification of N_δ

$$\tilde{N}_{\kappa_0, \varepsilon, \delta}(r, u, z_0) := \frac{1}{1 - br^\varepsilon} N_\delta(r, u, z_0),$$

where b is as in Theorem 3.3 (or Lemma 3.1), as well as the truncation of $\tilde{N}_{\kappa_0, \varepsilon, \delta}$

$$\hat{N}_{\kappa_0, \varepsilon, \delta}(r, u, z_0) := \min\{\tilde{N}_{\kappa_0, \varepsilon, \delta}(r, u, z_0), \kappa_0\}, \quad 0 < r < r_0 = r_0(\kappa_0, \varepsilon) = \frac{r_0(\varepsilon)}{\kappa_0^{2/\varepsilon}}.$$

When $z_0 = 0$, we simply write $N^0(r, u)$, $N_\delta(r, u)$, etc.

As demonstrated in [JP21, Theorem 5.4], the monotonicity of $W_{\kappa, \alpha, \varepsilon, \delta}$ readily implies that of the truncated frequency $\hat{N}_{\kappa_0, \varepsilon, \delta}$.

Theorem 3.4 (Almgren-type monotonicity formula). *Let $u, z_0, \kappa_0, \delta, \varepsilon$ be as in Theorem 3.3. Then $\hat{N}_{\kappa_0, \varepsilon, \delta}(r, u, z_0)$ is nondecreasing in $0 < r < r_0 = r_0(\kappa_0, \varepsilon) = \frac{r_0(\varepsilon)}{\kappa_0^{2/\varepsilon}}$.*

Proof. We may assume without loss of generality $z_0 = 0$. Take $r_0 = r_0(\kappa_0, \varepsilon)$ small so that $1 - br^\varepsilon > 0$. If $\hat{N}_{\kappa_0, \varepsilon, \delta}(r, u) < \kappa$ for some $r \in (0, r_0)$ and $\kappa \in (0, \kappa_0)$, then

$$W_{\kappa, \alpha, \varepsilon, \delta}(r, u) = \frac{e^{ar^\alpha}}{r^{2\kappa+2}} (1 - br^\varepsilon) \left(\int_{S_r} u^2 G \right) (\tilde{N}_{\kappa_0, \varepsilon, \delta}(r, u) - \kappa) < 0.$$

For any $0 < s < r$, we have by Theorem 3.3 that $W_{\kappa, \alpha, \varepsilon, \delta}(s, u) \leq W_{\kappa, \alpha, \varepsilon, \delta}(r, u) < 0$, and hence $\hat{N}_{\kappa_0, \varepsilon, \delta}(s, u) < \kappa$, as desired. \square

4. ALMGREN RESCALINGS AND BLOWUPS

The main objective of this section is to derive the proper lower bound for the frequency for almost minimizers at free boundary points. For this purpose, we deal with so-called Almgren blowups, which become global solutions of the parabolic Signorini problem. It is known that even-symmetric (in x_n -variable) solutions possess the minimal frequency of $3/2$.

In the study of the Signorini problem (both in elliptic and parabolic settings), the even symmetry of the solution with respect to the thin space is imperative. The symmetry ensures that the growth rate of the solution over the “thick” strip $S_r(z_0)$ match that over the “thin” strip $S'_r(z_0)$. This allows us to extract the information about the behavior of solutions on the thin space using the Almgren-type monotonicity formula.

In the case of solutions of the parabolic Signorini problem, the symmetry assumption is not restrictive, because if u is a solution then its even symmetrization

$$u^*(x', x_n, t) = \frac{u(x', x_n, t) + u(x', -x_n, t)}{2}$$

is still a solution. However, this property is not available for almost minimizers, as the even symmetrization can disrupt the almost Signorini property, even in the time-independent case (see [JPS24, Example 6.1]).

Therefore, in the remainder of this paper, we assume that the almost minimizer u is even symmetric in x_n -variable.

Next, we introduce another type of competitor for $u \in \mathcal{F}_{z_0}$ aside from homogeneous replacement. We say that v is a *parabolic Signorini replacement* of u in $S_r(t_0)$ if v is the solution of a parabolic Signorini problem in $S_r(t_0)$ with $v = u$ on $\mathbb{R}^n \times \{t_0 - r^2\}$ and $v - u \in L^2(t_0 - r^2, t_0; W_0^{1,2}(\mathbb{R}^n, G_{z_0}))$.

We remark that the regularity assumption on $u \in \mathcal{F}_{z_0}$ is not sufficient to ensure the existence of its parabolic Signorini replacement. To rectify this issue, we consider convolutions with mollifiers. For a standard mollifier $\varphi = \varphi(x)$ in \mathbb{R}^n and a small constant $\mu > 0$, we let $\varphi_\mu(x) := (1/\mu)^n \varphi(x/\mu)$. We set

$$(4.1) \quad u_\mu(x, t) := u * \varphi_\mu(x, t), \quad (x, t) \in S_1.$$

Then $u_\mu(\cdot, -r^2) \in W_\infty^2(\mathbb{R}^n)$ for a.e. $r \in (0, 1)$ and $\|u_\mu - u\|_{\mathcal{F}_{z_0}} \rightarrow 0$ as $\mu \rightarrow 0$. By Theorem A.1, for such r , there exists a unique parabolic Signorini replacement of u_μ in S_r .

Remark 4.1. u_μ satisfies the almost parabolic Signorini property in S_r , $0 < r < 1$, with a gauge function $\eta(r) = r^{\alpha/2}$ and additional additive error $C(n, \alpha) \|u - u_\mu\|_{\mathcal{F}_{z_0}}^2$.

Indeed, we assume without loss of generality $z_0 = 0$. Since $v := u - u_\mu + v_\mu$ is a valid competitor of u in S_r , we have by (1.2) and Young’s inequality

$$\begin{aligned} & \int_{S_r} (1 - r^{\alpha/2})(-t) |\nabla u_\mu|^2 G + (x \cdot \nabla u_\mu - 2t \partial_t u_\mu)(u_\mu - v_\mu) G \\ &= \int_{S_r} (1 - r^{\alpha/2})(-t) |\nabla u + \nabla(u_\mu - u)|^2 G + (-x \cdot \nabla u - 2t \partial_t u)(u - v) G \\ & \quad + (-x \cdot \nabla(u_\mu - u) - 2t \partial_t(u_\mu - u))(u_\mu - v_\mu) G \\ &\leq \int_{S_r} (1 - r^\alpha)(-t) |\nabla u|^2 G + (-x \cdot \nabla u - 2t \partial_t u)(u - v) G \\ & \quad + C(r) \|u - u_\mu\|_{\mathcal{F}_0}^2 + r^\alpha \int_{S_r} (u_\mu - v_\mu)^2 G \\ &\leq (1 + r^\alpha) \int_{S_r} (-t) |\nabla v|^2 G + r^\alpha \int_{S_r} (u - v)^2 G + \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}} \\ & \quad + C(r) \|u - u_\mu\|_{\mathcal{F}_0}^2 + r^\alpha \int_{S_r} (u_\mu - v_\mu)^2 G \end{aligned}$$

$$\begin{aligned} &\leq (1 + r^{\alpha/2}) \int_{S_r} (-t) |\nabla v_\mu|^2 G + 2r^{\alpha/2} \int_{S_r} (u_\mu - v_\mu)^2 G + \|u_\mu\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}} \\ &\quad + C(r) \|u - u_\mu\|_{\mathcal{F}_0}^2. \end{aligned}$$

Lemma 4.2. *Let $u \in \mathcal{F}_{z_0}$ satisfy the almost parabolic Signorini property at $z_0 \in Q'_{1/2}$. Suppose that u has a parabolic Signorini replacement v in $S_r(t_0)$. Then there exist constants $r_0 > 0$ and $C > 0$, depending only on α , such that if $0 < r < r_0$,*

$$(4.2) \quad \int_{S_r(t_0)} (t_0 - t) |\nabla(u - v)|^2 G_{z_0} \leq Cr^\alpha \int_{S_r(t_0)} (t_0 - t) |\nabla u|^2 G_{z_0} + C \|u\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}},$$

$$(4.3) \quad \int_{S_r(t_0)} (u - v)^2 G_{z_0} \leq Cr^\alpha \int_{S_r(t_0)} (t_0 - t) |\nabla u|^2 G_{z_0} + C \|u\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}}.$$

Proof. We may assume without loss of generality $z_0 = 0$. By the variational inequality of v , we have

$$\int_{S_r} (-2t) \nabla v \nabla(v - u) G + (-x \cdot \nabla v - 2t \partial_t v)(v - u) G \leq 0.$$

This, combined with the almost Signorini property of u (equation (1.2)), gives

$$\begin{aligned} &\int_{S_r} (-t) |\nabla(u - v)|^2 G \\ &= \int_{S_r} (-t) |\nabla u|^2 G - \int_{S_r} (-t) |\nabla v|^2 G + 2 \int_{S_r} (-t) \nabla(v - u) \nabla v G \\ &\leq r^\alpha \int_{S_r} (-t) (|\nabla u|^2 + |\nabla v|^2) G + \int_{S_r} (-x \cdot \nabla u - 2t \partial_t u)(v - u) G \\ &\quad + r^\alpha \int_{S_r} (u - v)^2 G + \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}} - \int_{S_r} (-x \cdot \nabla v - 2t \partial_t v)(v - u) G \\ &= \int_{S_r} (-x \cdot \nabla(u - v) - 2t \partial_t(u - v))(v - u) G + r^\alpha \int_{S_r} (-t) (|\nabla u|^2 + |\nabla v|^2) G \\ &\quad + r^\alpha \int_{S_r} (u - v)^2 G + \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}}. \end{aligned}$$

To compute the first term in the last line, we consider $\tilde{u}(y, \tau) := u(2e^{-\frac{\tau}{2}} y, -e^{-\tau})$ and $\tilde{v}(y, \tau) := v(2e^{-\frac{\tau}{2}} y, -e^{-\tau})$, which correspond to (3.5) with $\kappa = 0$. Since $\tilde{u} - \tilde{v} = 0$ on $\mathbb{R}^n \times \{-2 \ln r\}$, we have by Integration by parts

$$\begin{aligned} &\int_{S_r} (-x \cdot \nabla(u - v) - 2t \partial_t(u - v))(v - u) G \\ &= \frac{2}{\pi^{n/2}} \int_{\mathbb{R}^n \times (-2 \ln r, \infty)} (\partial_\tau(\tilde{u} - \tilde{v}))(\tilde{v} - \tilde{u}) e^{-|y|^2} e^{-\tau} dy d\tau \\ (4.4) \quad &= -\frac{1}{\pi^{n/2}} \int_{\mathbb{R}^n \times (-2 \ln r, \infty)} \partial_\tau((\tilde{u} - \tilde{v})^2) e^{-|y|^2} e^{-\tau} dy d\tau \\ &\leq -\frac{1}{\pi^{n/2}} \int_{\mathbb{R}^n \times (-2 \ln r, \infty)} (\tilde{u} - \tilde{v})^2 e^{-|y|^2} e^{-\tau} dy d\tau \\ &= -\int_{S_r} (u - v)^2 G dx dt. \end{aligned}$$

It then follows that

$$(4.5) \quad \int_{S_r} (-t)|\nabla(u-v)|^2 \leq r^\alpha \int_{S_r} (-t)(|\nabla u|^2 + |\nabla v|^2)G + \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}}.$$

This gives

$$\begin{aligned} \int_{S_r} (-t)|\nabla v|^2 G &\leq 2 \int_{S_r} (-t)|\nabla u|^2 G + 2 \int_{S_r} (-t)|\nabla(u-v)|^2 G \\ &\leq 4 \int_{S_r} (-t)|\nabla u|^2 G + 2r^\alpha \int_{S_r} (-t)|\nabla v|^2 G + 2\|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}}, \end{aligned}$$

which implies

$$(4.6) \quad \int_{S_r} (-t)|\nabla v|^2 G \leq C \int_{S_r} (-t)|\nabla u|^2 + C\|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}}, \quad r < r(\alpha).$$

By combining this with (4.5), we obtain (4.2).

Regarding (4.3), we use the almost parabolic Signorini property of u and the parabolic Signorini property of v (i.e., equations (1.2) and (1.1)) to have

$$\begin{aligned} &\int_{S_r} (-t)|\nabla u|^2 G + \int_{S_r} (-x \cdot \nabla u - 2t\partial_t u)(u-v)G \\ &\leq \int_{S_r} (-t)|\nabla v|^2 G + r^\alpha \int_{S_r} (u-v)^2 G + \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}} + r^\alpha \int_{S_r} (-t)(|\nabla u|^2 + |\nabla v|^2)G \\ &\leq \int_{S_r} (-t)|\nabla u|^2 G - \int_{S_r} (-x \cdot \nabla v - 2t\partial_t v)(v-u)G \\ &\quad + r^\alpha \int_{S_r} (u-v)^2 G + \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}} + r^\alpha \int_{S_r} (-t)(|\nabla u|^2 + |\nabla v|^2)G, \end{aligned}$$

and thus

$$\begin{aligned} &\int_{S_r} (-x \cdot \nabla(u-v) - 2t\partial_t(u-v))(u-v)G \\ &\leq r^\alpha \int_{S_r} (u-v)^2 G + \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}} + r^\alpha \int_{S_r} (-t)(|\nabla u|^2 + |\nabla v|^2)G. \end{aligned}$$

This, together with (4.4), gives

$$\int_{S_r} (u-v)^2 G \leq Cr^\alpha \int_{S_r} (-t)(|\nabla u|^2 + |\nabla v|^2)G + C\|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}}.$$

Finally, by combining this and (4.6), we conclude (4.3). \square

Corollary 4.3. *Suppose that $u \in \mathcal{F}_{z_0}$ satisfies the almost Signorini property at $z_0 \in Q'_{1/2}$. For u_μ be as in (4.1), let v_μ be the parabolic Signorini replacement of u_μ in $S_r(t_0)$. Then there are $r_0 > 0$ and $C > 0$, depending only on α , such that for $0 < r < r_0$*

$$\begin{aligned} \int_{S_r(t_0)} (t_0 - t)|\nabla(u_\mu - v_\mu)|^2 G_{z_0} &\leq Cr^{\alpha/2} \int_{S_r(t_0)} |\nabla u_\mu|^2 G_{z_0} + C\|u_\mu\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}} \\ &\quad + C_0(n, \alpha)\|u - u_\mu\|_{\mathcal{F}_{z_0}}^2, \\ \int_{S_r(t_0)} (u_\mu - v_\mu)^2 G_{z_0} &\leq Cr^{\alpha/2} \int_{S_r(t_0)} |\nabla u_\mu|^2 G_{z_0} + C\|u_\mu\|_{\mathcal{F}_{z_0}}^2 e^{-\frac{1}{r}} \\ &\quad + C_0(n, \alpha)\|u - u_\mu\|_{\mathcal{F}_{z_0}}^2. \end{aligned}$$

Proof. For the proof, we can use the almost parabolic Signorini property of u_μ (Remark 4.1) and follow the argument in Lemma 4.2. \square

For $z_0 = (x_0, t_0) \in \Gamma(u) \cap Q'_{1/2}$ and $0 < r < 1/2$, consider the *Almgren rescaling* of u at z_0

$$u_{z_0, r}^A(x, t) := \frac{u(rx + x_0, r^2t + t_0)}{\left(\frac{1}{r^2} \int_{S_r(t_0)} u^2 G_{z_0}\right)^{1/2}}, \quad (x, t) \in Q_{1/(2r)}.$$

It satisfies the normalization and scaling properties

$$\begin{aligned} \int_{S_1} (u_{z_0, r}^A)^2 G &= 1, \\ N^0(\rho, u_{z_0, r}^A, 0) &= N^0(\rho r, u, z_0), \quad \rho < 1/(2r). \end{aligned}$$

We will call the limits of $u_{z_0, r}^A$ over any sequence $r = r_j \rightarrow 0+$ *Almgren blowups* of u at z_0 , denoted by $u_{z_0, 0}^A$. When $z_0 = 0$, we simply write $u_r^A = u_{0, r}^A$ and $u_0^A = u_{0, 0}^A$.

Proposition 4.4 (Existence of Almgren blowups). *Let $z_0 \in Q'_{1/2} \cap \Gamma(u)$ be such that $\widehat{N}_{\kappa_0, \delta}(0+, u, z_0) = \kappa < \kappa_0$ for some $0 < \delta < 2$ and $\kappa_0 > 2$. Then every sequence of Almgren rescalings u_{z_0, r_j}^A , with $r_j \rightarrow 0+$, contains a subsequence, still denoted by r_j , such that for a function $u_{x_0, 0}^A \in W_{2, \text{loc}}^{1,0}(S_1, G) \cap C_{\text{loc}}^{1,0}(Q_1^\pm \cup Q'_1)$*

$$\begin{aligned} u_{z_0, r_j}^A &\rightarrow u_{z_0, 0}^A \quad \text{in } W_{2, \text{loc}}^{1,0}(S_1, G), \\ u_{z_0, r_j}^A &\rightarrow u_{z_0, 0}^A \quad \text{in } C_{\text{loc}}^{1,0}(Q_1^\pm \cup Q'_1). \end{aligned}$$

Moreover, $u_{z_0, 0}^A$ is a nonzero solution of the parabolic Signorini problem in S_1 , even in x_n .

Proof. We may assume without loss of generality $z_0 = 0$. From $\widehat{N}_{\kappa_0, \delta}(0+, u) = \kappa < \kappa_0$, we have $N_\delta(0+, u) = \widehat{N}_{\kappa_0, \delta}(0+, u) = \kappa$, and thus $N_\delta(r_j, u) < \kappa_0$ for small r_j . For such r_j ,

$$(4.7) \quad \int_{S_1} (-2t) |\nabla u_{r_j}^A|^2 G = N^0(1, u_{r_j}^A) = N^0(r_j, u) \leq N_\delta(r_j, u) < \kappa_0.$$

Combining this with $\int_{S_1} (u_{r_j}^A)^2 G = 1$, we see that $u_{r_j}^A$ is bounded in $W_2^{1,0}(S_1, G)$. Thus, there is a function $u_0^A \in W_{2, \text{loc}}^{1,0}(S_1, G)$ such that up to a subsequence

$$u_{r_j}^A \rightarrow u_0^A \quad \text{weakly in } W_{2, \text{loc}}^{1,0}(S_1, G).$$

Moreover, it is easy to see that each $u_{r_j}^A$ is an unweighted almost minimizers in $Q_{1/(2r_j)}$ with gauge function $\eta_{r_j}(\rho) = (r_j \rho)^\alpha \leq \rho^\alpha$. Thus, for small $\varepsilon > 0$ and $K \Subset Q_1 \cap \{t \leq -\varepsilon\}$, we have by Theorem 2.1

$$\begin{aligned} \|u_{r_j}^A\|_{C^{\alpha, \alpha/2}(K)} + \|\nabla u_{r_j}^A\|_{C^{\beta, \beta/2}(K \pm \cap K')} &\leq C(n, K, \alpha, \varepsilon) \|u_{r_j}^A\|_{W_2^{1,0}(Q_1 \cap \{t \leq -\varepsilon\})} \\ &\leq C(n, K, \alpha, \varepsilon) \|u_{r_j}^A\|_{W_2^{1,0}(S_1, G)}. \end{aligned}$$

This, along with the boundedness of $\{u_{r_j}^A\}$ in $W_2^{1,0}(S_1, G)$, yields

$$u_{r_j}^A \rightarrow u_0^A \quad \text{in } C_{\text{loc}}^{1,0}((Q_1^\pm \cup Q'_1) \cap \{t \leq -\varepsilon\}).$$

Taking $\varepsilon \searrow 0$ and using Cantor's diagonal argument, we infer that over a subsequence $r = r_j \searrow 0$

$$u_{r_j}^A \rightarrow u_0^A \quad \text{in } C_{\text{loc}}^{1,0}(Q_1^\pm \cup Q_1').$$

Now, for each r_j , we take $\mu_j > 0$ small so that the convolution $(u_{r_j}^A)_{\mu_j} = u_{r_j}^A * \varphi_{\mu_j}$ as in (4.1) satisfies $C_0(r_j, \alpha) \|(u_{r_j}^A)_{\mu_j} - u_{r_j}^A\|_{\mathcal{F}_0}^2 \rightarrow 0$ as $r_j \rightarrow 0$, where $C_0(r_j, \alpha)$ is as in Corollary 4.3. We then apply Theorem A.1. By considering $S_{1-\varepsilon}$ with small $\varepsilon > 0$ if necessary, we may assume that there exists the parabolic Signorini replacement $(v_{r_j})_{\mu_j}$ of $(u_{r_j}^A)_{\mu_j}$ in S_1 . By Corollary 4.3, (4.7) and the bound $N_\delta(r_j, u) < \kappa_0$, we have

$$\begin{aligned} & \int_{S_1} (-t) |\nabla(u_{r_j}^A - (v_{r_j})_{\mu_j})|^2 G \\ & \leq 2 \int_{S_1} (-t) |\nabla((u_{r_j}^A)_{\mu_j} - (v_{r_j})_{\mu_j})|^2 G + 2 \|(u_{r_j}^A)_{\mu_j} - u_{r_j}^A\|_{\mathcal{F}_0}^2 \\ & \leq Cr_j^{\alpha/2} \int_{S_1} (-t) |\nabla(u_{r_j}^A)_{\mu_j}|^2 G + C \|(u_{r_j}^A)_{\mu_j}\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r_j}} \\ & \quad + C(\alpha) C_0(r_j, \alpha) \|(u_{r_j}^A)_{\mu_j} - u_{r_j}^A\|_{\mathcal{F}_0}^2 \\ & \leq Cr_j^{\alpha/2} \int_{S_1} (-t) |\nabla u_{r_j}^A|^2 G + C \|u_{r_j}^A\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r_j}} + C(\alpha) C_0(r_j, \alpha) \|(u_{r_j}^A)_{\mu_j} - u_{r_j}^A\|_{\mathcal{F}_0}^2 \\ & \leq C\kappa_0 r_j^{\alpha/2} + CN_\delta(r_j, u) e^{-\frac{1}{r_j}} + C(\alpha) C_0(r_j, \alpha) \|(u_{r_j}^A)_{\mu_j} - u_{r_j}^A\|_{\mathcal{F}_0}^2 \\ & \rightarrow 0 \quad \text{as } r_j \rightarrow 0 +. \end{aligned}$$

Similarly, we can obtain

$$\int_{S_1} (u_{r_j}^A - (v_{r_j})_{\mu_j})^2 G \rightarrow 0 \quad \text{as } r_j \rightarrow 0 +.$$

These estimates, combined with the boundedness of $u_{r_j}^A$ in $W_2^{1,0}(S_1, G)$, implies that $(v_{r_j})_{\mu_j}$ is also bounded in $W_2^{1,0}(S_1, G)$ and

$$u_{r_j}^A - (v_{r_j})_{\mu_j} \rightarrow 0 \quad \text{strongly in } W_{2,\text{loc}}^{1,0}(S_1, G),$$

and hence

$$(v_{r_j})_{\mu_j} \rightarrow u_0^A \quad \text{weakly in } W_{2,\text{loc}}^{1,0}(S_1, G).$$

By [DGPT17, Theorem 7.3], we have that u_0^A is a solution of the parabolic Signorini problem in S_1 and that $(v_{r_j})_{\mu_j}$ is bounded in $W_2^{2,1}(S_1, G)$, which gives $(v_{r_j})_{\mu_j}^2 G \rightarrow (u_0^A)^2 G$ strongly in $L^1(S_1)$. This, combined with the observation

$$2 \int_{S_1} (v_{r_j})_{\mu_j}^2 G \geq \int_{S_1} (u_{r_j}^A)^2 G - 2 \int_{S_1} (u_{r_j}^A - (v_{r_j})_{\mu_j})^2 G \rightarrow 1 \quad \text{as } r_j \rightarrow 0+,$$

produces $\int_{S_1} (u_0^A)^2 G = \lim_{r_j \rightarrow 0+} \int_{S_1} (v_{r_j})_{\mu_j}^2 G \geq 1/2$, and hence $u_0^A \not\equiv 0$ in S_1 . This completes the proof. \square

In the subsequent lemma, we show that the additional exponential term present in N_δ is insignificant, as previously mentioned. This enables us to employ the results established in [DGPT17] while proving a lower bound on Almgren's frequency for almost minimizers in Lemma 4.6.

Lemma 4.5. *Suppose u satisfies the almost parabolic Signorini property at $z_0 \in \Gamma(u) \cap Q'_{1/2}$. If $\widehat{N}_{\kappa_0, \delta}(0+, u, z_0) = \kappa < \kappa_0$ for some $0 < \delta < 2$ and $\kappa_0 > 2$, then*

$$\lim_{r \rightarrow 0} N^0(r, u, z_0) = \lim_{r \rightarrow 0} \frac{\int_{S_r(t_0)} 2(t_0 - t) |\nabla u|^2 G_{z_0}}{\int_{S_r(t_0)} u^2 G_{z_0}} = \kappa.$$

Proof. Without loss of generality, we may assume $z_0 = 0$. From $\widehat{N}_{\kappa_0, \delta}(0+, u) = \kappa < \kappa_0$, we have $N_\delta(0+, u) = \kappa$, thus it is enough to show

$$\lim_{r \rightarrow 0} \frac{\|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r} r^{-\delta}}}{\int_{S_r} u^2 G} = 0.$$

To this aim, we assume to the contrary that

$$\limsup_{r \rightarrow 0} \frac{\|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r} r^{-\delta}}}{\int_{S_r} u^2 G} = a_0 \in (0, \kappa].$$

Then we have for some sequence $r = r_j \searrow 0$

$$\frac{\|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r_j} r_j^{-\delta}}}{\int_{S_{r_j}} u^2 G} > \frac{a_0}{2} \quad \text{and} \quad \frac{\int_{S_{r_j}} (-2t) |\nabla u|^2 G}{\int_{S_{r_j}} u^2 G} > \kappa - 2a_0.$$

Fix $\varepsilon \in (0, \alpha)$ and for $\delta' = \delta + \varepsilon/4$, consider $N_{\delta'}(r, u) = \frac{\int_{S_r} (-2t) |\nabla u|^2 G + \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r} r^{-\delta'}}$

Then for $r = r_j$,

$$N_{\delta'}(r_j, u) = \frac{\int_{S_{r_j}} (-2t) |\nabla u|^2 G}{\int_{S_{r_j}} u^2 G} + \frac{\|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r_j} r_j^{-\delta}}}{\int_{S_{r_j}} u^2 G} r_j^{-\varepsilon/4} > \frac{a_0}{2} r_j^{-\varepsilon/4} + \kappa - 2a_0.$$

Thus, we have for any $\kappa_1 > \kappa_0$

$$\widetilde{N}_{\kappa_1, \varepsilon, \delta'}(r_j, u) \geq N_{\delta'}(r_j, u) > \frac{a_0}{2} r_j^{-\varepsilon/4} + \kappa - 2a_0,$$

and hence

$$(4.8) \quad \widehat{N}_{\kappa_1, \delta'}(0+, u) = \lim_{r_j \rightarrow 0} \min\{\widetilde{N}_{\kappa_1, \varepsilon, \delta'}(r_j, u), \kappa_1\} = \kappa_1.$$

On the other hand, since $N_\delta(0+, u) = \kappa$, there is $s_0 > 0$, independent of κ_1 , such that $N_\delta(r, u) < 2\kappa$ for $r < s_0$. Moreover, by Theorem 3.4, there is a constant $c_0 = c_0(\varepsilon) > 0$ such that for any $\kappa_1 > \kappa_0$ and $0 < \delta < 2$, $\widehat{N}_{\kappa_1, \varepsilon, \delta}(r, u)$ is nondecreasing in $0 < r < \frac{2c_0}{\kappa_1^{2/\varepsilon}}$. By taking c_0 smaller if necessary, we may assume $c_0 < s_0$. Note that if κ_1 is sufficiently large, then $1 - \frac{128(\kappa_1+1)}{\varepsilon} \cdot \frac{c_0(\varepsilon)^\varepsilon}{\kappa_1^2} > 1/2$. Moreover, we have for $0 < r < \frac{2c_0}{\kappa_0^{2/\varepsilon}}$

$$N_{\delta'}(r, u) = \frac{\int_{S_r} (-2t) |\nabla u|^2 G}{\int_{S_r} u^2 G} + \frac{\|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r} r^{-\delta}}}{\int_{S_r} u^2 G} r^{-\varepsilon/4} \leq N_\delta(r, u) r^{-\varepsilon/4} \leq 2\kappa r^{-\varepsilon/4}.$$

It then follows that

$$\widetilde{N}_{\kappa_1, \varepsilon, \delta'} \left(\frac{c_0}{\kappa_1^{2/\varepsilon}}, u \right) = \frac{1}{1 - \frac{128(\kappa_1+1)}{\varepsilon} \left(\frac{c_0}{\kappa_1^{2/\varepsilon}} \right)^\varepsilon} N_{\delta'} \left(\frac{c_0}{\kappa_1^{2/\varepsilon}}, u \right)$$

$$\leq 4\kappa \left(\frac{c_0}{\kappa_1^{2/\varepsilon}} \right)^{-\varepsilon/4} \leq C(\varepsilon, \kappa, s_0) \kappa_1^{1/2}.$$

Therefore, we have for large κ_1

$$\widehat{N}_{\kappa_1, \varepsilon, \delta'} \left(\frac{c_0}{\kappa_1^{2/\varepsilon}}, u \right) \leq C(\varepsilon, \kappa, s_0) \kappa_1^{1/2} < \kappa_1.$$

This, along with (4.8), contradicts the monotonicity of $\widehat{N}_{\kappa_1, \varepsilon, \delta'}$ in $\left(0, \frac{2c_0}{\kappa_1^{2/\varepsilon}}\right)$. \square

Lemma 4.6. *Let u be an almost minimizer for the parabolic Signorini problem in Q_1 and $z_0 \in \Gamma(u) \cap Q'_{1/2}$. If $\widehat{N}_{\kappa_0, \delta}(0+, u, z_0) = \kappa < \kappa_0$ for some $0 < \delta < 2$ and $\kappa_0 > 2$, then*

$$\kappa = 3/2 \quad \text{or} \quad \kappa \geq 2.$$

Proof. Without loss of generality, we may assume $z_0 = 0$. Let $u_0^A = \lim_{r_j \rightarrow 0} u_{r_j}^A$ be an Almgren blowup. Recall that it is a solution of the parabolic Signorini problem in S_1 . From Lemma 4.5, we find that for any $0 < \rho < 1$

$$N^0(\rho, u_0^A) = \lim_{r_j \rightarrow 0} N^0(\rho, u_{r_j}^A) = \lim_{r_j \rightarrow 0} N^0(\rho r_j, u) = \kappa,$$

which implies that u_0^A is parabolically homogeneous of degree κ in S_1 (see the proof of [DGPT17, Theorem 7.3]), and by homogeneity, can be extended to S_∞ . In addition, by the Complementarity condition (Lemma 2.2), we have $u_0^A(0) = |\widehat{\nabla} u_0^A(0)| = 0$, where $\widehat{\nabla} u_0^A$ is the even extension of ∇u_0^A from S_1^+ to S_1 . Thus we can repeat the proof of [DGPT17, Proposition 8.1] to get $\kappa > 1$. Then, it follows from [DGPT17, Proposition 8.5] that either $\kappa = 3/2$ or $\kappa \geq 2$. \square

Corollary 4.7. *Let u be an almost minimizer for the parabolic Signorini problem in Q_1 and $z_0 \in \Gamma(u) \cap Q'_{1/2}$. Then for any $\kappa_0 > 2$, $\varepsilon \in (0, \alpha]$ and $0 < \delta < 2$,*

$$W_{3/2, \alpha, \varepsilon, \delta}(r, u, z_0) \geq 0 \quad \text{for } 0 < r < r_0,$$

where r_0 is as in Theorem 3.3.

Proof. The proof follows by using Lemma 4.6 and repeating the argument in [JP21, Corollary 6.3]. \square

5. GROWTH ESTIMATES

In this section we establish the optimal growth of almost minimizers at free boundary points (Lemma 5.4).

Given $\kappa \geq 3/2$, we define the κ -homogeneous rescalings of u at $z_0 = (x_0, t_0) \in \Gamma(u) \cap Q'_{1/2}$ by

$$u_{z_0, r}(x, t) := u_{z_0, r}^{(\kappa)}(x, t) = \frac{u(x_0 + rx, t_0 + r^2 t)}{r^\kappa}, \quad (x, t) \in S_1.$$

Note that $\widehat{N}_{\kappa_0, \delta}(0+, u, z_0)$ and $\widetilde{N}_\delta(0+, u, z_0)$ are independent of α and ε .

Lemma 5.1 (Weak growth estimates). *Suppose u satisfies the almost parabolic Signorini property at $z_0 \in \Gamma(u) \cap Q'_{1/2}$. If $\widehat{N}_{\kappa_0, \delta}(0+, u, z_0) \geq \kappa \geq 1$ for some $\kappa \leq \kappa_0$, $\kappa_0 > 2$ and $0 < \delta < 2$, then for any $0 < \tilde{\varepsilon} \leq \alpha/2 < 1$,*

$$(5.1) \quad \begin{aligned} \int_{S_r(t_0)} u^2 G_{z_0} dxdt &\leq C(\kappa_0, \tilde{\varepsilon}) \|u\|_{\mathcal{F}_{z_0}}^2 r^{2\kappa+2-\tilde{\varepsilon}}, \\ \int_{S_r(t_0)} 2(t_0 - t) |\nabla u|^2 G_{z_0} dxdt &\leq C(\kappa_0, \tilde{\varepsilon}) \|u\|_{\mathcal{F}_{z_0}}^2 r^{2\kappa+2-\tilde{\varepsilon}} \end{aligned}$$

for $0 < r < r_0 = r_0(\kappa_0, \tilde{\varepsilon})$.

Proof. Without loss of generality we assume $z_0 = 0$. Note that for every $\varepsilon = 2\tilde{\varepsilon} \in (0, \alpha]$, the condition $\widehat{N}_{\kappa_0, \delta}(0+, u) \geq \kappa$ implies that $\widehat{N}_{\kappa_0, \varepsilon, \delta}(r, u) \geq \kappa$ for $0 < r < r_0(\kappa_0, \varepsilon)$. Then we also have $\tilde{N}_{\kappa_0, \varepsilon, \delta}(r, u) \geq \kappa$ for such r , and thus

$$(5.2) \quad W_{\kappa, \alpha, \varepsilon, \delta}(r, u) = \frac{e^{ar^\alpha}}{r^{2\kappa+2}} \left(\int_{S_r} u^2 G \right) (1 - br^\varepsilon) \left(\tilde{N}_{\kappa_0, \varepsilon, \delta}(r, u) - \kappa \right) \geq 0.$$

For $u_r = u_{0,r}^{(\kappa)}$, we define

$$(5.3) \quad m(r) := \int_{S_1} u_r^2 G = \frac{1}{r^{2\kappa+2}} \int_{S_r} u^2 G.$$

Using

$$\frac{d}{dr} u_r(x, t) = -\frac{1}{r^{\kappa+1}} \left(\kappa u(rx, r^2 t) - (rx) \cdot \nabla u(rx, r^2 t) - 2(r^2 t) \partial_t u(rx, r^2 t) \right),$$

we can compute

$$\begin{aligned} m'(r) &= 2 \int_{S_1} u_r(x, t) \left(\frac{d}{dr} u_r(x, t) \right) G(x, t) dxdt \\ &= -\frac{2}{r^{2\kappa+1}} \int_{S_1} u(rx, r^2 t) \left(\kappa u(rx, r^2 t) - (rx) \cdot \nabla u(rx, r^2 t) \right. \\ &\quad \left. - 2(r^2 t) \partial_t u(rx, r^2 t) \right) G(x, t) dxdt \\ &= -\frac{2}{r^{2\kappa+3}} \int_{S_r} u(\kappa u - x \cdot \nabla u - 2t \partial_t u) G dxdt. \end{aligned}$$

By applying Theorem 3.3, we further have

$$|m'(r)| = \frac{2}{r^{2\kappa+3}} \left| \int_{S_r} u(\kappa u - x \cdot \nabla u - 2t \partial_t u) G \right| \leq \frac{4r^{-\varepsilon/2}}{\kappa} \frac{d}{dr} W_{\kappa, \alpha, \varepsilon, \delta}(r, u).$$

This, along with (5.2), gives that for $0 < s < r < r_0$

$$\begin{aligned} |m(r) - m(s)| &\leq \int_s^r |m'(\rho)| d\rho \leq \frac{4}{\kappa} \int_s^r \rho^{-\varepsilon/2} \frac{d}{d\rho} W_{\kappa, \alpha, \varepsilon, \delta}(\rho, u) d\rho \\ &\leq 4s^{-\varepsilon/2} \int_s^r \frac{d}{d\rho} W_{\kappa, \alpha, \varepsilon, \delta}(\rho, u) d\rho \leq 4s^{-\varepsilon/2} W_{\kappa, \alpha, \varepsilon, \delta}(r, u). \end{aligned}$$

In particular, we have

$$m(r) \leq m(r_0) + 4r^{-\varepsilon/2} W_{\kappa, \alpha, \varepsilon, \delta}(r_0, u).$$

This implies the first bound. The second one is then derived by utilizing the first one and the monotonicity $W_{\kappa, \alpha, \varepsilon, \delta}(r, u) \leq W_{\kappa, \alpha, \varepsilon, \delta}(r_0, u)$. \square

In the rest of this section, we remove the extra $\tilde{\varepsilon} > 0$ in Lemma 5.1 and obtain the optimal growth in the case of the least frequency $\kappa = 3/2$. To this end, we first derive the polynomial decay estimate of the Weiss-type energy $W_{3/2, \alpha, \varepsilon, \delta}$. Following the approach in the elliptic counterpart [JP21], we achieve this decay by utilizing the epiperimetric inequality. However, it is worth noting that employing the epiperimetric inequality in our context is considerably more complex and technical.

Before we state the parabolic epiperimetric inequality from [Shi20], we introduce two types of “standard” Weiss energy functionals that will be used in this section.

$$V_{3/2}^0(t, v) := \frac{1}{(-t)^{3/2}} \int_{\mathbb{R}^n} \left((-2t) |\nabla v(x, t)|^2 - \frac{3}{2} v(x, t)^2 \right) G(x, t) dx,$$

$$W_{3/2}^0(r, v) := \frac{1}{r^5} \int_{S_r} \left((-2t) |\nabla v|^2 - \frac{3}{2} v^2 \right) G dx dt.$$

Theorem 5.2 (Epiperimetric Inequality [Shi20]). *Let v be a solution of the parabolic Signorini problem in S_1 . Then there is a dimensional constant $\xi \in (0, 1)$ such that*

$$V_{3/2}^0(t/e, v) \leq (1 - \xi) V_{3/2}^0(t, v), \quad -1 < t < 0.$$

Lemma 5.3. *Fix $\kappa_0 > 2$ and $0 < \delta < 2$. Suppose u satisfies the almost parabolic Signorini property at $z_0 \in \Gamma(u) \cap Q'_{1/2}$. Then there exists $\sigma = \sigma(n, \alpha) > 0$ such that for any $\varepsilon \in (0, \alpha]$,*

$$(5.4) \quad 0 \leq W_{3/2, \alpha, \varepsilon, \delta}(r, u, z_0) \leq C r^{\min\{\sigma, 3\varepsilon/4\}}, \quad 0 < r < r_0(\kappa_0, n, \alpha, \varepsilon)$$

with $C = C(\kappa_0, n, \alpha, \varepsilon) \|u\|_{\mathcal{F}_{z_0}}^2$.

Proof. We split our proof into several steps.

Step 1. We may assume $z_0 = 0$. We suppose that for $r \in (0, 1)$ a.e., there exists a parabolic Signorini replacement v of u in S_r . For the simplicity of the presentation, we will derive (5.4) under this assumption in Step 1-Step 3 and consider the general case without the existence assumption in Step 4.

By the epiperimetric inequality (Theorem 5.2) and the monotonicity of $V_{3/2}^0$ ([Shi20, Lemma 2.2]), we get

$$\begin{aligned} r^5 W_{3/2}^0(r, v) &= \int_{-r^2}^0 (-t)^{3/2} V_{3/2}^0(t, v) dt = \sum_{m=0}^{\infty} \int_{-\frac{r^2}{e^m}}^{-\frac{r^2}{e^{m+1}}} (-t)^{3/2} V_{3/2}^0(t, v) dt \\ &\leq \sum_{m=0}^{\infty} (1 - \xi)^m \int_{-\frac{r^2}{e^m}}^{-\frac{r^2}{e^{m+1}}} (-t)^{3/2} V_{3/2}^0(e^m t, v) dt \\ &= \sum_{m=0}^{\infty} \left(\frac{1 - \xi}{e} \right)^m \int_{-r^2}^{-\frac{r^2}{e}} \left(-\frac{s}{e^m} \right)^{3/2} V_{3/2}^0(s, v) ds \\ &\leq \sum_{m=0}^{\infty} \left(\frac{1 - \xi}{e^{5/2}} \right)^m V_{3/2}^0(-r^2, v) \int_{-r^2}^{-\frac{r^2}{e}} (-s)^{3/2} ds \\ &= \frac{r^5 (e^{5/2} - 1)}{(5/2)(e^{5/2} + \xi - 1)} V_{3/2}^0(-r^2, v). \end{aligned}$$

Thus

$$(5.5) \quad W_{3/2}^0(r, v) \leq \frac{1 - \eta}{5/2} V_{3/2}^0(-r^2, v) = \frac{1 - \eta}{5/2} V_{3/2}^0(-r^2, u),$$

where $\eta := \frac{\xi}{e^{5/2} + \xi - 1} \in (0, 1/e)$. On the other hand, by differentiating

$$r^5 W_{3/2}^0(r, u) = \int_{-r^2}^0 (-t)^{3/2} V_{3/2}^0(t, u) dt$$

with respect to r , we obtain after simplification

$$(5.6) \quad V_{3/2}^0(-r^2, u) = (5/2)W_{3/2}^0(r, u) + \frac{r}{2} \frac{d}{dr} W_{3/2}^0(r, u).$$

This, along with (5.5), gives

$$(5.7) \quad W_{3/2}^0(r, v) \leq (1 - \eta)W_{3/2}^0(r, u) + \left(\frac{1 - \eta}{5}\right) r \frac{d}{dr} W_{3/2}^0(r, u).$$

For $\tilde{\varepsilon} = \tilde{\varepsilon}(\alpha) \in (0, \alpha)$ to be determined later, by applying (3.12) with $\kappa = 3/2$ and Lemma 5.1, we have

$$(5.8) \quad \begin{aligned} W_{3/2}^0(r, u) &= \frac{r^\alpha}{r^5} \int_{S_r} (-2t) |\nabla u|^2 + \frac{1}{r^5} \int_{S_r} \left((1 - r^\alpha)(-2t) |\nabla u|^2 - \frac{3}{2} u^2 \right) G \\ &\leq C(\kappa_0, \alpha) \|u\|_{\mathcal{F}_0}^2 r^{\alpha - \tilde{\varepsilon}} + (1 + r^\alpha) W_{3/2}^0(r, v) \\ &\quad + \frac{1}{r^5} \int_{S_r} [3/2 r^\alpha v^2 + (3/2 + 2r^\alpha)(u - v)^2 \\ &\quad - 2(3/2u - x \cdot \nabla u - 2t \partial_t u)(u - v)] G + \frac{2 \|u\|_{\mathcal{F}_0}^2}{r^5} e^{-\frac{1}{r}}. \end{aligned}$$

Combining this and (5.7), we obtain

$$\begin{aligned} W_{3/2}^0(r, u) &\leq (1 + r^\alpha)(1 - \eta)W_{3/2}^0(r, u) + (1 + r^\alpha) \left(\frac{1 - \eta}{5}\right) r \frac{d}{dr} W_{3/2}^0(r, u) \\ &\quad + \frac{1}{r^5} \int_{S_r} [3/2 r^\alpha v^2 + (3/2 + 2r^\alpha)(u - v)^2 \\ &\quad - 2(3/2u - x \cdot \nabla u - 2t \partial_t u)(u - v)] G + C(\kappa_0, \alpha, u) r^{\alpha - \tilde{\varepsilon}}, \end{aligned}$$

which is equivalent to

$$(5.9) \quad \begin{aligned} \frac{d}{dr} W_{3/2}^0(r, u) &\geq \frac{5(\eta - r^\alpha(1 - \eta))}{(1 + r^\alpha)(1 - \eta)} \frac{W_{3/2}^0(r, u)}{r} \\ &\quad + \frac{5}{(1 + r^\alpha)(1 - \eta)r^6} \int_{S_r} [-3/2 r^\alpha v^2 - (3/2 + 2r^\alpha)(u - v)^2 \\ &\quad + 2(3/2u - x \cdot \nabla u - 2t \partial_t u)(u - v)] G \\ &\quad - C(\kappa_0, \alpha, u) r^{\alpha - \tilde{\varepsilon} - 1}. \end{aligned}$$

Step 2. In this step, we simplify (5.9) by estimating the second term in its right-hand side. To this aim, we decompose

$$\begin{aligned} \frac{1}{r^6} \int_{S_r} [-3/2 r^\alpha v^2 - (3/2 + 2r^\alpha)(u - v)^2 + 2(3/2u - x \cdot \nabla u - 2t \partial_t u)(u - v)] G \\ = I + II + III. \end{aligned}$$

Concerning II , we use Lemma 4.2 and Lemma 5.1 to get

$$(5.10) \quad \begin{aligned} II &= -\frac{3/2 + 2r^\alpha}{r^6} \int_{S_r} (u-v)^2 G \geq -\frac{C}{r^6} \left(r^\alpha \int_{S_r} (-t) |\nabla u|^2 + \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r}} \right) \\ &\geq -C(\kappa_0, \alpha, u) r^{\alpha-\tilde{\varepsilon}-1}. \end{aligned}$$

Regarding I , from (5.10) and Lemma 5.1 we infer

$$(5.11) \quad I \geq -\frac{3r^\alpha}{r^6} \left(\int_{S_r} u^2 G + \int_{S_r} (u-v)^2 G \right) \geq -C(\kappa_0, \alpha, u) r^{\alpha-\tilde{\varepsilon}-1}.$$

It remains to consider III . Following the argument in the proof of Lemma 3.2, we have for $\kappa = 3/2$

$$\begin{aligned} &\int_{S_r} (\kappa(u-v) - x \cdot \nabla(u-v) - 2t\partial_t(u-v))(u-v)G \, dxdt \\ &= \frac{1}{\pi^{n/2}} \int_{\mathbb{R}^n \times (-2\ln r, \infty)} \partial_\tau((\tilde{u}-\tilde{v})^2) e^{-|y|^2} e^{-(\kappa+1)\tau} \, dyd\tau \\ &\geq \frac{\kappa+1}{\pi^{n/2}} \int_{\mathbb{R}^n \times (-2\ln r, \infty)} (\tilde{u}-\tilde{v})^2 e^{-|y|^2} e^{-(\kappa+1)\tau} \, dyd\tau \\ &\geq 0. \end{aligned}$$

This, together with Young's inequality and (5.10), yields

$$(5.12) \quad \begin{aligned} III &= \frac{2}{r^6} \int_{S_r} (3/2(u-v) - x \cdot \nabla(u-v) - 2t\partial_t(u-v))(u-v)G \\ &\quad + \frac{2}{r^6} \int_{S_r} (3/2v - x \cdot \nabla v - 2t\partial_tv)(u-v)G \\ &\geq -\frac{1}{r^{6-\tilde{\varepsilon}}} \int_{S_r} (3/2v - x \cdot \nabla v - 2t\partial_tv)^2 G - \frac{1}{r^{6+\tilde{\varepsilon}}} \int_{S_r} (u-v)^2 G. \end{aligned}$$

The second term in the last line is estimated in (5.10). To estimate the first one, we bring the following computation made in the proof of [DGPT17, Theorem 13.1]

$$\frac{d}{dr} W_{3/2}^0(r, v) \geq \frac{2}{r^6} \int_{S_r} (3/2v - x \cdot \nabla v - 2t\partial_tv)^2 G.$$

It then follows that

$$\begin{aligned} &-\frac{1}{r^{6-\tilde{\varepsilon}}} \int_{S_r} (3/2v - x \cdot \nabla v - 2t\partial_tv)^2 G \\ &\geq -\frac{r^{\tilde{\varepsilon}}}{2} \frac{d}{dr} W_{3/2}^0(r, v) \\ &= -r^{\tilde{\varepsilon}-1} V_{3/2}^0(-r^2, v) + 5/2 r^{\tilde{\varepsilon}-1} W_{3/2}^0(r, v) \\ &\geq -r^{\tilde{\varepsilon}-1} V_{3/2}^0(-r^2, u) + \frac{5}{2(1+r^\alpha)} r^{\tilde{\varepsilon}-1} (W_{3/2}^0(r, u) - Cr^{\alpha-\tilde{\varepsilon}} + r(I + II + III)) \\ &\geq -Cr^{\alpha-\tilde{\varepsilon}-1} + O(r^{\tilde{\varepsilon}}) \left(\frac{W_{3/2}^0(r, u)}{r} \right) + O(r^{\tilde{\varepsilon}}) \left(\frac{d}{dr} W_{3/2}^0(r, u) \right) \\ &\quad - r^{\tilde{\varepsilon}} \left(\frac{1}{r^{6-\tilde{\varepsilon}}} \int_{S_r} (3/2v - x \cdot \nabla v - 2t\partial_tv)^2 G \right), \end{aligned}$$

where we applied (5.6) for v in the third line, and used (5.8) in the fourth line and (5.6), (5.10), (5.11), (5.12) in the last step. This implies

$$\begin{aligned} & -\frac{1}{r^{6-\tilde{\varepsilon}}} \int_{S_r} (3/2v - x \cdot \nabla v - 2t\partial_t v)^2 G \\ & \geq -Cr^{\alpha-\tilde{\varepsilon}-1} + O(r^{\tilde{\varepsilon}}) \left(\frac{W_{3/2}^0(r, u)}{r} \right) + O(r^{\tilde{\varepsilon}}) \left(\frac{d}{dr} W_{3/2}^0(r, u) \right), \end{aligned}$$

which, combined with (5.12), gives

$$III \geq -C(\kappa_0, \alpha, u)r^{\alpha-2\tilde{\varepsilon}-1} + O(r^{\tilde{\varepsilon}}) \left(\frac{W_{3/2}^0(r, u)}{r} \right) + O(r^{\tilde{\varepsilon}}) \left(\frac{d}{dr} W_{3/2}^0(r, u) \right).$$

Now, by taking $\tilde{\varepsilon} = \alpha/3$, we conclude

$$\begin{aligned} & I + II + III \\ & \geq -C(\kappa_0, \alpha, u)r^{\alpha/3-1} + O(r^{\alpha/3}) \left(\frac{W_{3/2}^0(r, u)}{r} \right) + O(r^{\alpha/3}) \left(\frac{d}{dr} W_{3/2}^0(r, u) \right). \end{aligned}$$

Therefore, (5.9) can be simplified to

$$(5.13) \quad \frac{d}{dr} W_{3/2}^0(r, u) \geq \left(\frac{5\eta}{1-\eta} + O(r^{\alpha/3}) \right) \frac{W_{3/2}^0(r, u)}{r} - C(\kappa_0, \alpha, u)r^{\alpha/3-1}.$$

Step 3. We consider the Weiss-type energy $W_{3/2, \alpha} = W_{3/2, \alpha, \alpha, 1}$ with $\varepsilon = \alpha$ and $\delta = 1$. By Corollary 4.7 and Lemma 5.1,

$$(5.14) \quad \begin{aligned} W_{3/2}^0(r, u) &= e^{-ar^\alpha} W_{3/2, \alpha}(r, u) - \frac{3/2br^\alpha}{r^5} \int_{S_r} u^2 G - \frac{\|u\|_{\mathcal{F}_0}^2}{r^5} e^{-\frac{1}{r}} r^{-1} \\ &\geq -C_0(\kappa_0, \alpha, u)r^{\alpha/2}, \quad 0 < r < r_0(\kappa_0, \alpha). \end{aligned}$$

We recall $\eta \in (0, 1/e)$ and use (5.13) and (5.14) to get the differential inequality for $W_{3/2}^0(r, u)$:

$$\begin{aligned} \frac{d}{dr} W_{3/2}^0(r, u) &\geq \left(\frac{5\eta}{1-\eta} + O(r^{\alpha/3}) \right) \left(\frac{W_{3/2}^0(r, u) + C_0 r^{\alpha/3}}{r} - \frac{C_0 r^{\alpha/3}}{r} \right) - Cr^{\alpha/3-1} \\ &\geq 5\eta \left(\frac{W_{3/2}^0(r, u) + C_0 r^{\alpha/3}}{r} \right) - Cr^{\alpha/3-1} \\ &\geq 5\eta \frac{W_{3/2}^0(r, u)}{r} - C_1 r^{\alpha/3-1}. \end{aligned}$$

We take $\sigma = \sigma(n, \alpha)$ such that $0 < \sigma < \min\{5\eta, \alpha/3\}$, and use the differential inequality and (5.14) to obtain

$$\begin{aligned} & \frac{d}{dr} \left[W_{3/2}^0(r, u)r^{-\sigma} + \frac{2C_1}{\alpha/3-\sigma} r^{\alpha/3-\sigma} \right] \\ &= r^{-\sigma} \left(\frac{d}{dr} W_{3/2}^0(r, u) - \frac{\sigma}{r} W_{3/2}^0(r, u) \right) + 2C_1 r^{\alpha/3-\sigma-1} \\ &\geq r^{-\sigma-1} (5\eta - \sigma) W_{3/2}^0(r, u) + C_1 r^{\alpha/3-\sigma-1} \\ &\geq -C_2(\kappa_0, \alpha) \|u\|_{\mathcal{F}_0}^2 r^{\alpha/2-\sigma-1} + C_3(\kappa_0, \alpha) \|u\|_{\mathcal{F}_0}^2 r^{\alpha/3-\sigma-1} \end{aligned}$$

$$\geq 0, \quad 0 < r < r_0(\kappa_0, n, \alpha).$$

This readily gives

$$W_{3/2}^0(r, u) \leq C(\kappa_0, n, \alpha, u)r^\sigma.$$

To complete the proof, let $\varepsilon \in (0, \alpha]$ and $\delta \in (0, 2)$ be given. Then, by applying Lemma 5.1 (with $\tilde{\varepsilon} = \varepsilon/4$), we conclude that

$$\begin{aligned} W_{3/2, \alpha, \varepsilon, \delta}(r, u) &= e^{ar^\alpha} W_{3/2}^0(r, u) + \frac{3/2 e^{ar^\alpha} b r^\varepsilon}{r^5} \int_{S_r} u^2 G + \frac{\|u\|_{\mathcal{F}_0}^2 e^{ar^\alpha}}{r^5} e^{-\frac{1}{r}} r^{-\delta} \\ &\leq C(\kappa_0, n, \alpha, \varepsilon) \|u\|_{\mathcal{F}_0}^2 r^{\min\{\sigma, 3\varepsilon/4\}}, \quad 0 < r < r_0(\kappa_0, n, \alpha, \varepsilon). \end{aligned}$$

Step 4. To close the argument, we need to remove the assumption on the existence of the parabolic Signorini replacement made in Step 1. To this end, we consider $u_\mu = u * \varphi_\mu$ as in (4.1). Then, for $r \in (0, 1)$ a.e., the parabolic Signorini replacement v_μ of u_μ in S_r exists. We observe that only the following properties of u are used in Step 1 and Step 2: the almost parabolic Signorini property (equation (1.2)), Lemma 4.2 and the weak growth estimates with $\kappa = 3/2$ (Lemma 5.1). We have already seen in Remark 4.1 and Corollary 4.3 that u_μ satisfies analogues of the first two properties. Moreover, by using the triangle inequality, it is easily seen that u_μ satisfies the following analogue of (5.1) with $\kappa = 3/2$: for any $0 < \tilde{\varepsilon} \leq \alpha/2$

$$\begin{aligned} \int_{S_r} u_\mu^2 G &\leq C(\kappa_0, \alpha) \|u_\mu\|_{\mathcal{F}_0}^2 r^{5-\tilde{\varepsilon}} + C(r, \alpha) \|u - u_\mu\|_{\mathcal{F}_0}^2, \\ \int_{S_r} (-2t) |\nabla u_\mu|^2 G &\leq C(\kappa_0, \alpha) \|u_\mu\|_{\mathcal{F}_0}^2 r^{5-\tilde{\varepsilon}} + C(r, \alpha) \|u - u_\mu\|_{\mathcal{F}_0}^2. \end{aligned}$$

Now, with these properties of u_μ at hand, we can follow the argument in Step 1 and Step 2 with u_μ and v_μ in the place of u and v to deduce an analogue of (5.13):

$$\begin{aligned} \frac{d}{dr} W_{3/2}^0(r, u_\mu) &\geq \left(\frac{5\eta}{1-\eta} + O(r^{\alpha/6}) \right) \frac{W_{3/2}^0(r, u_\mu)}{r} - C(\kappa_0, \alpha) \|u_\mu\|_{\mathcal{F}_0}^2 r^{\alpha/6-1} \\ &\quad - C(r, \alpha, \kappa_0) \|u - u_\mu\|_{\mathcal{F}_0}^2. \end{aligned}$$

Taking $\mu \rightarrow 0$, we obtain the differential inequality (5.13) concerning $W_{3/2}^0(r, u)$ for $r \in (0, 1)$ a.e., but with $\alpha/6$ in the place of $\alpha/3$. Then, (5.4) readily follows by arguing as in Step 3 with obvious modifications. \square

As in [JP21], by using the polynomial decay of $W_{3/2, \alpha, \varepsilon, \delta}$ we can improve Lemma 5.1 when $\kappa = 3/2$ and derive the optimal growth.

Lemma 5.4 (Optimal growth estimate). *Fix $\kappa_0 > 2$. Suppose that $u \in \mathcal{F}_{z_0}$ satisfies the almost parabolic Signorini property at $z_0 \in \Gamma(u) \cap Q_{1/2}^I$. Then,*

$$\begin{aligned} \int_{S_r(t_0)} u^2 G_{z_0} dx dt &\leq C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_{z_0}}^2 r^5, \\ \int_{S_r(t_0)} 2(t_0 - t) |\nabla u|^2 G_{z_0} dx dt &\leq C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_{z_0}}^2 r^5 \end{aligned}$$

for $0 < r < r_0 = r_0(\kappa_0, n, \alpha)$.

Proof. We may assume $z_0 = 0$. Take $\varepsilon = \varepsilon(n, \alpha) > 0$ small so that $3\varepsilon/4 < \sigma$ for $\sigma = \sigma(n, \alpha)$ as in Lemma 5.3. Following the computation in the proof of Lemma 5.1 with $\delta = 1$, we see that for any $0 < s < r < r_0(\kappa_0, n, \alpha)$,

$$|m(r) - m(s)| \leq 4s^{-\varepsilon/2} W_{3/2, \alpha, \varepsilon, 1}(r).$$

By Lemma 5.3, we further have

$$|m(r) - m(s)| \leq C s^{-\varepsilon/2} r^{3\varepsilon/4},$$

with $C = C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_0}^2$. Then, by a dyadic argument, we can obtain that

$$(5.15) \quad |m(r) - m(s)| \leq C r^{\varepsilon/4}.$$

Indeed, let $k = 0, 1, 2, \dots$, be such that $r/2^{k+1} < s \leq r/2^k$. Then,

$$\begin{aligned} |m(r) - m(s)| &\leq \sum_{j=1}^k |m(r/2^{j-1}) - m(r/2^j)| + |m(r/2^k) - m(s)| \\ &\leq C \sum_{j=1}^{k+1} (r/2^j)^{-\varepsilon/2} (r/2^{j-1})^{3\varepsilon/4} = C \left(r^{1/4} 2^{3/4} \right)^\varepsilon \sum_{j=1}^{k+1} 2^{-j/4} \\ &\leq C r^{\varepsilon/4}. \end{aligned}$$

In particular, we have

$$m(r) \leq m(r_0) + C r_0^{\varepsilon/4} \leq C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_0}^2, \quad 0 < r < r_0(\kappa_0, n, \alpha).$$

This implies the first bound. The second bound follows from the first one and the monotonicity $W_{3/2, \alpha, \varepsilon, 1}(r, u) \leq W_{3/2, \alpha, \varepsilon, 1}(r_0, u)$. \square

6. 3/2-HOMOGENEOUS BLOWUPS

In this section, we consider the so-called 3/2-homogeneous blowups of almost minimizers at free boundary points. They are the limits of 3/2-homogeneous rescalings, which are well-defined thanks to the optimal growth estimates. We achieve their uniqueness through controlling the ‘‘rotation’’ of the rescalings.

Concerning the κ -homogeneous rescalings, for the rest of this paper, we focus exclusively on the case $\kappa = 3/2$. Thus we simply write $u_{z_0, r} = u_{z_0, r}^{(3/2)}$.

Fix $z_0 = (x_0, t_0) \in \Gamma(u) \cap Q'_{1/2}$ and $R > 1$, and let $r_0 = r_0(\kappa_0, n, \alpha)$ be as stated in Lemma 5.4. We have for $0 < r < r_0$

$$\begin{aligned} \int_{S_R} (-2t) |\nabla u_{z_0, r}|^2 G &= \frac{1}{r^5} \int_{S_{Rr}(t_0)} 2(t_0 - t) |\nabla u|^2 G_{z_0} \leq C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_{z_0}}^2 R^5, \\ \int_{S_R} u_{z_0, r}^2 G &= \frac{1}{r^5} \int_{S_{Rr}(t_0)} u^2 G_{z_0} \leq C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_{z_0}}^2 R^5. \end{aligned}$$

Thus, for a sequence $r = r_j \rightarrow 0+$, $u_{z_0, r_j} \rightarrow u_{z_0, 0}$ weakly in $W_{2, \text{loc}}^{1,0}(S_R, G)$. Moreover, $u_{z_0, r}$ is an unweighted almost minimizer with a gauge function $\eta_r(\rho) = (r\rho)^\alpha \leq \rho^\alpha$. Given $\varepsilon > 0$ and $K \Subset Q_R \cap \{t \leq -\varepsilon\}$, we infer from Theorem 2.1 that there is a constant $C > 0$ such that for any $0 < r < r_0$,

$$\begin{aligned} \|u_{z_0, r}\|_{C^{\alpha, \alpha/2}(K)} + \|\nabla u_{z_0, r}\|_{C^{\beta, \beta/2}(K \pm \cup K')} &\leq C \|u_{z_0, r}\|_{W_2^{1,0}(Q_R \cap \{t \leq -\varepsilon\})} \\ &\leq C \|u_{z_0, r}\|_{W_2^{1,0}(S_R, G)}, \end{aligned}$$

and hence over a sequence $r = r_j \rightarrow 0+$

$$u_{z_0, r_j} \rightarrow u_{z_0, 0} \quad \text{in } C_{\text{loc}}^{1,0}((Q_R^\pm \cup Q'_R) \cap \{t \leq -\varepsilon\}).$$

Now, taking $\varepsilon \rightarrow 0$ and $R \rightarrow \infty$ and using Cantor's diagonal argument, we can find a subsequence $r = r_j \rightarrow 0+$ such that for some $u_{z_0, 0} \in C_{\text{loc}}^{1,0}(S_\infty^\pm \cup S'_\infty)$

$$u_{z_0, r_j} \rightarrow u_{z_0, 0} \quad \text{in } C_{\text{loc}}^{1,0}(S_\infty^\pm \cup S'_\infty).$$

We call such $u_{z_0, 0}$ a *3/2-homogeneous blowup* of u at z_0 .

Lemma 6.1 (Rotation estimate). *Suppose that u satisfies the almost parabolic Signorini property at $z_0 \in \Gamma(u) \cap Q'_{1/2}$. Then there exists $\sigma = \sigma(n, \alpha) > 0$ such that for any $0 < s < r < r_0 = r_0(\kappa_0, n, \alpha)$ and $-1 < t < 0$,*

$$\int_{\mathbb{R}^n} |u_{z_0, r}(x, t) - u_{z_0, s}(x, t)| G_{z_0}(x, t) dx \leq C(-t)^{3/4 + \sigma} r^{2\sigma}$$

with $C = C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_{z_0}}^2$.

Proof. Without loss of generality, we may assume $z_0 = 0$. We fix $\varepsilon = \alpha$, $\delta = 1$ and $\kappa = 3/2$, and simply write $W_{\kappa, \rho} = W_{\kappa, \alpha, \varepsilon, \delta, \rho}$. By using (3.13) in Theorem 3.3, we infer that for $R > R_0(\kappa_0, \alpha)$

$$\begin{aligned} & W_{\kappa, e^{-R/2}}(e^{3-R/2}, u) - W_{\kappa, e^{-R/2}}(e^{1-R/2}, u) \\ & \geq \frac{2\kappa + 1}{\pi^{n/2}} \int_{R-6}^{R-2} \frac{e^{-(\kappa+1)(r+R)}}{(e^{-(\kappa+1)r} - e^{-(\kappa+1)R})^2} \int_{\mathbb{R}^n} (\tilde{u}(y, R) - \tilde{u}(y, r))^2 e^{-|y|^2} dy dr. \end{aligned}$$

Since $\kappa = 3/2$, we have for $R - 6 < r < R - 2$

$$\frac{e^{-(\kappa+1)(r+R)}}{(e^{-(\kappa+1)r} - e^{-(\kappa+1)R})^2} \geq \frac{e^{-(\kappa+1)(2R-2)}}{(e^{-(\kappa+1)(R-6)} - e^{-(\kappa+1)R})^2} = \frac{e^5}{(e^{15} - 1)^2},$$

thus

$$\begin{aligned} & \int_{R-6}^{R-2} \int_{\mathbb{R}^n} (\tilde{u}(y, R) - \tilde{u}(y, r))^2 e^{-|y|^2} dy dr \\ & \leq C(n) \left(W_{\kappa, e^{-R/2}}(e^{3-R/2}, u) - W_{\kappa, e^{-R/2}}(e^{1-R/2}, u) \right). \end{aligned}$$

To estimate the right-hand side of this previous inequality, we note that by Lemma 5.3, there is $\sigma = \sigma(n, \alpha) > 0$ such that for $W_\kappa = W_{\kappa, 0}$ (i.e., $W_{\kappa, \alpha, \varepsilon, \delta, \rho}$ with $\kappa = 3/2$, $\varepsilon = \alpha$, $\delta = 1$ and $\rho = 0$) and for $R > R_0 = R_0(\kappa_0, n, \alpha)$,

$$0 \leq W_\kappa(e^{-R/2}, u) \leq C e^{-2\sigma R}, \quad W_\kappa(e^{1-R/2}, u) \geq 0, \quad W_\kappa(e^{3-R/2}, u) \leq C e^{-2\sigma R}$$

with $C = C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_0}^2$. Then

$$\begin{aligned} & W_{\kappa, e^{-R/2}}(e^{3-R/2}, u) \\ & = \frac{e^{a e^{(3-R/2)\alpha}}}{e^{-(\kappa+1)(R-6)} - e^{-(\kappa+1)R}} \left(\int_{S_{e^{3-R/2}}} ((-2t)|\nabla u|^2 - \kappa(1 - b e^{(3-R/2)\alpha})u^2) G \right. \\ & \quad \left. - \int_{S_{e^{-R/2}}} ((-2t)|\nabla u|^2 - \kappa(1 - b e^{(3-R/2)\alpha})u^2) G + \|u\|_{\mathcal{F}_0}^2 e^{-e^{R/2-3}} e^{R/2-3} \right) \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{e^{6(\kappa+1)} - 1} \left(e^{6(\kappa+1)} W_\kappa(e^{3-R/2}, u) \right. \\
 &\quad \left. - \frac{e^{ae^{(3-R/2)\alpha}}}{e^{-(\kappa+1)R}} \int_{S_{e^{-R/2}}} ((-2t)|\nabla u|^2 - \kappa(1 - be^{(3-R/2)\alpha})u^2)G \right) \\
 &\leq \frac{1}{e^{6(\kappa+1)} - 1} \left(e^{6(\kappa+1)} W_\kappa(e^{3-R/2}, u) - W_\kappa(e^{-R/2}, u) + O(e^{-\alpha/2R}) \|u\|_{\mathcal{F}_0}^2 \right) \\
 &\leq Ce^{-2\sigma R}.
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 &W_{\kappa, e^{-R/2}}(e^{1-R/2}, u) \\
 &= \frac{1}{e^{2(\kappa+1)} - 1} \left(e^{2(\kappa+1)} W_\kappa(e^{1-R/2}, u) \right. \\
 &\quad \left. - \frac{e^{ae^{(1-R/2)\alpha}}}{e^{-(\kappa+1)R}} \int_{S_{e^{-R/2}}} ((-2t)|\nabla u|^2 - \kappa(1 - be^{(1-R/2)\alpha})u^2)G \right) \\
 &\geq \frac{1}{e^{2(\kappa+1)} - 1} \left(e^{2(\kappa+1)} W_\kappa(e^{1-R/2}, u) - W_\kappa(e^{-R/2}, u) + O(e^{-\alpha/2R}) \|u\|_{\mathcal{F}_0}^2 \right) \\
 &\geq -Ce^{-2\sigma R}.
 \end{aligned}$$

Thus

$$\int_{R-6}^{R-2} \int_{\mathbb{R}^n} (\tilde{u}(y, R) - \tilde{u}(y, r))^2 e^{-|y|^2} dy dr \leq Ce^{-2\sigma R},$$

and hence by Cauchy-Schwarz inequality

$$\int_{R-6}^{R-2} \int_{\mathbb{R}^n} |\tilde{u}(y, R) - \tilde{u}(y, r)| e^{-|y|^2} dy dr \leq C_0 e^{-\sigma R}, \quad R > R_1(\kappa_0, n, \alpha).$$

Then, for $R > R_1$ and $3 < \eta < 6$,

$$\begin{aligned}
 (6.1) \quad &\int_{\mathbb{R}^n} \left| \tilde{u}(y, R + \eta) - \int_R^{R+1} \tilde{u}(y, \tau) d\tau \right| e^{-|y|^2} dy \\
 &\leq \int_{(R+\eta)-6}^{(R+\eta)-2} \int_{\mathbb{R}^n} |\tilde{u}(y, R + \eta) - \tilde{u}(y, \tau)| e^{-|y|^2} dy d\tau \\
 &\leq C_0 e^{-\sigma(R+\eta)}.
 \end{aligned}$$

We claim that for any $k \in \mathbb{N}$, $3 < \eta < 5$ and $R > R_1$,

$$(6.2) \quad \int_{\mathbb{R}^n} \left| \tilde{u}(y, R + k\eta) - \int_R^{R+1} \tilde{u}(y, \tau) d\tau \right| e^{-|y|^2} dy \leq C_0 \sum_{j=1}^k e^{-\sigma(R+\eta j)}.$$

Indeed, we prove it by induction on $k \in \mathbb{N}$. (6.2) is true for $k = 1$ by (6.1). If (6.2) is true for $k - 1$, then the induction hypothesis and (6.1) yield

$$\begin{aligned}
 &\int_{\mathbb{R}^n} \left| \tilde{u}(y, R + k\eta) - \int_R^{R+1} \tilde{u}(y, \tau) d\tau \right| e^{-|y|^2} dy \\
 &\leq \int_{\mathbb{R}^n} \left| \tilde{u}(y, R + \eta + (k-1)\eta) - \int_{R+\eta}^{R+\eta+1} \tilde{u}(y, \rho) d\rho \right| e^{-|y|^2} dy
 \end{aligned}$$

$$\begin{aligned}
& + \int_{\mathbb{R}^n} \left| \int_{R+\eta}^{R+\eta+1} \tilde{u}(y, \rho) d\rho - \int_R^{R+1} \tilde{u}(y, \tau) d\tau \right| e^{-|y|^2} dy \\
& \leq C_0 \sum_{j=1}^{k-1} e^{-\sigma(R+\eta+\eta j)} + \int_{R+\eta}^{R+\eta+1} \int_{\mathbb{R}^n} \left| \tilde{u}(y, \rho) - \int_R^{R+1} \tilde{u}(y, \tau) d\tau \right| e^{-|y|^2} dy d\rho \\
& \leq C_0 \sum_{j=2}^k e^{-\sigma(R+\eta j)} + \int_{R+\eta}^{R+\eta+1} C_0 e^{-\sigma\rho} d\rho \leq C_0 \sum_{j=1}^k e^{-\sigma(R+\eta j)}.
\end{aligned}$$

Now, let $S > R > R_1(\kappa_0, n, \alpha) + 12$ be given. Then we can choose $k \in \mathbb{N}$ and $\eta \in (3, 5)$ such that $S = R - 12 + \eta k$. By (6.2), we have

$$\begin{aligned}
& \int_{\mathbb{R}^n} |\tilde{u}(y, R) - \tilde{u}(y, S)| e^{-|y|^2} dy \\
& \leq \int_{\mathbb{R}^n} \left| \tilde{u}(y, R - 12 + 3 \cdot 4) - \int_{R-12}^{R-11} \tilde{u}(y, \tau) d\tau \right| e^{-|y|^2} dy \\
& \quad + \int_{\mathbb{R}^n} \left| \int_{R-12}^{R-11} \tilde{u}(y, \tau) d\tau - \tilde{u}(y, R - 12 + k\eta) \right| e^{-|y|^2} dy \\
& \leq C_0 \sum_{j=1}^3 e^{-\sigma(R-12+4j)} + C_0 \sum_{j=1}^k e^{-\sigma(R-12+\eta j)} \leq C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_0}^2 e^{-\sigma R}.
\end{aligned}$$

To complete the proof, define $r_0 = r_0(\kappa_0, n, \alpha) = e^{-1/2(R_1(\kappa_0, n, \alpha)+12)}$ and let $-1 < t < 0$ and $0 < s < r < r_0$ be given. By using $u_r(x, t) = \frac{u(rx, r^2 t)}{r^\kappa} = (-t)^{\kappa/2} \tilde{u}\left(\frac{x}{2\sqrt{-t}}, -\ln(-t) - 2\ln r\right)$, we conclude

$$\begin{aligned}
& \int_{\mathbb{R}^n} |u_r(x, t) - u_s(x, t)| G(x, t) dx \\
& = \frac{(-t)^{\kappa/2}}{\pi^{n/2}} \int_{\mathbb{R}^n} |\tilde{u}(y, -\ln(-t) - 2\ln r) - \tilde{u}(y, -\ln(-t) - 2\ln s)| e^{-|y|^2} dy \\
& \leq C(\kappa_0, n, \alpha) \|u\|_{\mathcal{F}_0}^2 (-t)^{3/4+\sigma} r^{2\sigma}. \quad \square
\end{aligned}$$

Lemma 6.2. *Let u, z_0, σ, r_0, C be as in Lemma 6.1. Then, for $0 < r < r_0$ and $-1 < t < 0$,*

$$\int_{\mathbb{R}^n} |u_{z_0, r}(x, t) - u_{z_0, 0}(x)| G(x, t) dx \leq C(-t)^{3/4+\sigma} r^{2\sigma}.$$

In particular, the blowup $u_{z_0, 0}$ is unique.

Proof. If $u_{z_0, 0}$ is the limit of u_{z_0, s_j} , $s_j \rightarrow 0$, then the first part of the lemma follows from Lemma 6.1 by taking $s_j \rightarrow 0$. For the second part, let $\bar{u}_{z_0, 0}$ be another blowup. Then we have

$$\int_{\mathbb{R}^n} |u_{z_0, 0}(x, t) - \bar{u}_{z_0, 0}(x, t)| G(x, t) dx = 0, \quad -1 < t < 0,$$

thus $u_{z_0, 0} = \bar{u}_{z_0, 0}$. □

7. REGULARITY OF THE REGULAR SET

In this last section, we prove one of the most crucial results in this paper, the regularity of the regular set.

Recall that the limit $\hat{N}_{\kappa_0, \delta}(0+, u, z_0) = \lim_{r \rightarrow 0+} \hat{N}_{\kappa_0, \varepsilon, \delta}(r, u, z_0)$ is independent of ε .

Definition 7.1 (Regular points). Let u be an almost minimizer for the parabolic Signorini problem in Q_1 . We say that a free boundary point $z_0 \in Q'_{1/2}$ is *regular* if

$$\hat{N}_{\kappa_0, \delta}(0+, u, z_0) = 3/2 \quad \text{for some } \kappa_0 > 2 \text{ and } 0 < \delta < 2.$$

We denote the set of all regular points of u by $\mathcal{R}(u)$ and call it *regular set*.

In view of Lemma 4.5, we have at every regular point z_0

$$N^0(0+, u, z_0) = 3/2.$$

In addition, regular points have the following characterization.

Remark 7.2. z_0 is a regular point if and only if

$$\lim_{\delta \rightarrow 0} N_\delta(0+, u, z_0) = \inf_{0 < \delta < 2} N_\delta(0+, u, z_0) = 3/2.$$

Proof. If z_0 is a regular point, then $3/2 = \hat{N}_{\kappa_0, \delta_0}(0+, u, z_0) = N_{\delta_0}(0+, u, z_0)$ for some $\kappa_0 > 2$ and $0 < \delta_0 < 2$. This, along with Lemma 4.6 and the fact that $\delta \mapsto N_\delta(0+, u, z_0)$ is nondecreasing, implies $3/2 \leq \hat{N}_{\kappa_0, \delta}(0+, u, z_0) \leq \hat{N}_{\kappa_0, \delta_0}(0+, u, z_0) = 3/2$ for every $\delta \in (0, \delta_0)$, which readily gives $3/2 = N_\delta(0+, u, z_0)$ for any $0 < \delta < \delta_0$. Therefore, we get $\lim_{\delta \rightarrow 0} N_\delta(0+, u, z_0) = \inf_{0 < \delta < 2} N_\delta(0+, u, z_0) = 3/2$.

To prove the opposite direction, we fix $\kappa_0 > 2$. Take $\delta_1 > 0$ such that $N_\delta(0+, u, z_0) < 7/4$ for $\delta \in (0, \delta_1)$. Then, by Lemmas 4.5 and 4.6, $3/2 \leq \hat{N}_{\kappa_0, \delta}(0+, u, z_0) = N^0(0+, u, z_0) \leq N_\delta(0+, u, z_0)$ for $0 < \delta < \delta_1$. Taking $\delta \rightarrow 0$ yields $N^0(0+, u, z_0) = 3/2$. This in turn gives that $\hat{N}_{\kappa_0, \delta}(0+, u, z_0) = 3/2$ for $0 < \delta < \delta_1$, and we conclude that z_0 is a regular point. \square

With the monotonicity of the frequency $\hat{N}_{\kappa_0, \varepsilon, \delta}$ (Theorem 3.4) and the frequency gap (Lemma 4.6) at hand, we can prove the relative openness of the regular set by following the argument in [JP21, Corollary 9.5].

Corollary 7.3. *The regular set $\mathcal{R}(u)$ is a relatively open subset of $\Gamma(u)$.*

Lemma 7.4 (Nondegeneracy at regular points). *Suppose that u satisfies the almost parabolic Signorini property at $z_0 = (x_0, t_0) \in \mathcal{R}(u)$. Then*

$$\liminf_{t \rightarrow 0} \int_{S_1} (u_{z_0, t})^2 G = \liminf_{t \rightarrow 0} \frac{1}{t^5} \int_{S_r(t_0)} u^2 G_{z_0} > 0.$$

Proof. By using (5.15) and the Weiss-type monotonicity formula, we can employ the contradiction argument as in [JP21, Lemma 9.2] to prove Lemma 7.4. \square

Proposition 7.5. *If u satisfies the almost parabolic Signorini property at $z_0 \in \mathcal{R}(u)$, then*

$$u_{z_0, 0}(x, t) = c_{z_0} \operatorname{Re}(x' \cdot e_{z_0} + i|x_n|)^{3/2} \quad \text{in } S_\infty$$

for some $c_{z_0} > 0$ and $e_{z_0} \in \partial B'_1$.

Proof. Without loss of generality, we may assume $z_0 = 0$. Let $r_j \rightarrow 0+$ be a sequence such that $u_{r_j} \rightarrow u_0$ in $C_{\text{loc}}^{1,0}(S_\infty^\pm \cup S'_\infty)$. Fix $R > 1$, and consider j large so that $Rr_j < 1$. For such r_j , we take $\mu_j > 0$ small so that $u_{\mu_j} = u * \varphi_{\mu_j}$ as in (4.1) satisfies $C_0(r_j, \alpha) \|u - u_{\mu_j}\|_{\mathcal{F}_0}^2 \rightarrow 0$ as $r_j \rightarrow 0$, where $C_0(r_j, \alpha)$ is as in Corollary 4.3. We let v_{μ_j} be the parabolic Signorini replacement of u_{μ_j} in S_R , and denote its 3/2-homogeneous rescaling by $(v_{\mu_j})_{r_j}(x, t) = \frac{v_{\mu_j}(r_j x, r_j^2 t)}{r_j^{3/2}}$. Then, by Corollary 4.3 and Lemma 5.4, we have

$$\begin{aligned} \int_{S_R} (-t) |\nabla(u_{r_j} - (v_{\mu_j})_{r_j})|^2 G &= \frac{1}{r_j^5} \int_{S_{Rr_j}} (-t) |\nabla(u - v_{\mu_j})|^2 G \\ &\leq \frac{2}{r_j^5} \left(\int_{S_{Rr_j}} (-t) |\nabla(u_{\mu_j} - v_{\mu_j})|^2 G + \|u - u_{\mu_j}\|_{\mathcal{F}_0}^2 \right) \\ &\leq \frac{C}{r_j^5} \left((Rr_j)^{\alpha/2} \int_{S_{Rr_j}} (-t) |\nabla u_{\mu_j}|^2 G + \|u_{\mu_j}\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r_j}} + C_0(r_j, \alpha) \|u - u_{\mu_j}\|_{\mathcal{F}_0}^2 \right) \\ &\leq \frac{C}{r_j^5} \left((Rr_j)^{\alpha/2} \int_{S_{Rr_j}} (-t) |\nabla u|^2 G + \|u\|_{\mathcal{F}_0}^2 e^{-\frac{1}{r_j}} + C_0(r_j, \alpha) \|u - u_{\mu_j}\|_{\mathcal{F}_0}^2 \right) \\ &\rightarrow 0 \quad \text{as } r_j \rightarrow 0. \end{aligned}$$

Similarly, we can obtain

$$\int_{S_R} (u_{r_j} - (v_{\mu_j})_{r_j})^2 G \rightarrow 0 \quad \text{as } r_j \rightarrow 0.$$

Thus $(v_{\mu_j})_{r_j} \rightarrow u_0$ weakly in $W_{2,\text{loc}}^{1,0}(S_R, G)$, and hence u_0 is a solution of the parabolic Signorini problem in S_R . Since $R > 1$ is arbitrary, we see that u_0 is the solution in S_∞ .

Next, we compare u_r and Almgren rescalings u_r^A

$$u_r = u_r^A \lambda(r), \quad \lambda(r) = \frac{\left(\frac{1}{r^2} \int_{S_r} u^2 G \right)^{1/2}}{r^{3/2}}.$$

It follows from Lemma 5.4 and Lemma 7.4 that

$$0 < \liminf_{r \rightarrow 0+} \lambda(r) \leq \limsup_{r \rightarrow 0+} \lambda(r) < \infty.$$

Thus, for a sequence $r_j \rightarrow 0+$, $u_0 = \lambda_0 u_0^A$ for some constant $\mu_0 \in (0, \infty)$. We have shown in the proof of Lemma 4.6 that u_0^A is 3/2-parabolically homogeneous in S_1 . Therefore, u_0 is also 3/2-parabolically homogeneous in S_1 , which can be extended to S_∞ by applying the unique continuation for caloric function in S_∞^\pm . In view of [DGPT17, Proposition 8.5], we conclude that

$$u_0(x, t) = c \operatorname{Re}(x' \cdot e + i|x_n|)^{3/2} \quad \text{in } S_\infty, \quad c > 0, e \in \partial B'_1. \quad \square$$

Lemma 7.6 (Continuous dependence of blowups). *Let $u \in \mathcal{F}$ be an almost minimizer for the parabolic Signorini problem in Q_1 . If $z_1, z_2 \in \mathcal{R}(u)$ and $\|z_1 - z_2\| < r_1$, then*

$$\int_{\partial B_1} |u_{z_1,0} - u_{z_2,0}| dSx \leq C \|z_1 - z_2\|^\gamma,$$

with $r_1 = r_1(\kappa_0, n, \alpha)$, $C = C(\kappa_0, n, \alpha, u)$ and $\gamma = \gamma(n, \alpha) > 0$.

Proof. Let $r_0 = r_0(\kappa_0, n, \alpha)$ and $\sigma = \sigma(n, \alpha)$ be as in Lemma 6.2. We have for every $0 < r < r_0$

$$\begin{aligned}
 & \int_{B_1} |u_{z_1,0}(x) - u_{z_2,0}(x)|G(x, -1) dx \\
 & \leq \int_{B_1} |u_{z_1,0}(x) - u_{z_1,r}(x, -1)|G(x, -1) dx \\
 & \quad + \int_{B_1} |u_{z_2,0}(x) - u_{z_2,r}(x, -1)|G(x, -1) dx \\
 & \quad + \int_{B_1} |u_{z_1,r}(x, -1) - u_{z_2,r}(x, -1)|G(x, -1) dx \\
 & \leq Cr^{2\sigma} + \frac{C(n)}{r^{3/2}} \int_{B_1} |u(x_1 + rx, t_1 - r^2) - u(x_2 + rx, t_2 - r^2)| dx.
 \end{aligned}$$

Since $(x_1 + rx, t_1 - r^2)$ and $(x_2 + rx, t_2 - r^2)$ are contained in $Q_{3/4}$ for every $x \in B_1$, we have by Theorem 2.1

$$|u(x_1 + rx, t_1 - r^2) - u(x_2 + rx, t_2 - r^2)| \leq C\|z_1 - z_2\|^{1/2}.$$

By taking $r = \|z_1 - z_2\|^{\frac{1}{4\sigma+3}}$ (which is possible if $r_1^{\frac{1}{4\sigma+3}} < r_0$), we obtain

$$\begin{aligned}
 \int_{B_1} |u_{z_1,0}(x) - u_{z_2,0}(x)|G(x, -1) dx & \leq C \left(r^{2\sigma} + \frac{\|z_1 - z_2\|^{1/2}}{r^{3/2}} \right) \\
 & = C(\kappa_0, n, \alpha, u)\|z_1 - z_2\|^\gamma, \quad \gamma = \frac{2\sigma}{4\sigma+3}.
 \end{aligned}$$

Now, the lemma follows by the boundedness of $G(x, -1)$ and the homogeneity of $u_{z_1,0}$ and $u_{z_2,0}$. \square

The following lemma follows from Proposition 7.5 and Lemma 7.6 by repeating the argument in [GPS16, Lemma 7.5].

Lemma 7.7. *Let $u \in \mathcal{F}$ be an almost minimizer for the parabolic Signorini problem in Q_1 , and $z_0 \in \mathcal{R}(u) \cap Q'_{1/4}$. Then there exist $\rho > 0$, depending on z_0 , and $\gamma = \gamma(n, \alpha) > 0$ such that $Q'_\rho(z_0) \cap \Gamma(u) \subset \mathcal{R}(u)$ and if $u_{z_j,0}(x) = c_{z_j} \operatorname{Re}(x' \cdot e_{z_j} + i|x_n|)^{3/2}$ is the unique 3/2-parabolically homogeneous blowup at $z_j \in Q'_\rho(z_0) \cap \Gamma(u)$, $j = 1, 2$, then*

$$\begin{aligned}
 |c_{z_1} - c_{z_2}| & \leq C_0 \|z_1 - z_2\|^\gamma, \\
 |e_{z_1} - e_{z_2}| & \leq C_0 \|z_1 - z_2\|^\gamma
 \end{aligned}$$

with a constant C_0 depending on z_0 .

We are now ready to prove the central result in this paper, the regularity of the regular set.

Theorem 7.8 (Regularity of the regular set). *Let $u \in \mathcal{F}$ be an almost minimizer for the parabolic Signorini problem in Q_1 . If $z_0 = (x_0, t_0) \in \mathcal{R}(u) \cap Q'_{1/4}$, there exists $\rho > 0$, depending on z_0 , such that possibly after a rotation in \mathbb{R}^{n-1} , one has $Q'_\rho(z_0) \cap \Gamma(u) \subset \mathcal{R}(u)$, and*

$$Q'_\rho(z_0) \cap \Gamma(u) = \{(x', t) \in Q'_\rho(z_0) : x_{n-1} = g(x'', t)\},$$

for a function g with $\nabla''g \in C^{\gamma, \gamma/2}$ for some $\gamma = \gamma(n, \alpha) \in (0, 1)$.

Proof. Since the proof of this theorem follows the lines of [JP21, Theorem 9.7], we shall provide only the outline of the proof.

Since $\mathcal{R}(u)$ is relatively open in $\Gamma(u)$, we have $Q'_{2\rho}(z_0) \cap \Gamma(u) \subset \mathcal{R}(u)$ for small $\rho > 0$. We claim that for any $\varepsilon > 0$, there exists $r_\varepsilon > 0$ such that for any $\bar{z} \in Q'_\rho(z_0) \cap \Gamma(u)$ and $0 < r < r_\varepsilon$, there holds

$$(7.1) \quad \|u_{\bar{z},r} - u_{\bar{z},0}\|_{C^{1,0}(\overline{Q_1^\pm})} < \varepsilon.$$

Indeed, towards a contradiction, suppose there are sequences $r_j \rightarrow 0$ and $\bar{z}_j \in Q'_\rho(z_0) \cap \Gamma(u)$ such that for some $\varepsilon_0 > 0$

$$\|u_{\bar{z}_j,r_j} - u_{\bar{z}_j,0}\|_{C^{1,0}(\overline{Q_1^\pm})} \geq \varepsilon_0.$$

Up to a subsequence, we have $\bar{z}_j \rightarrow \bar{z}_0 \in \overline{Q'_\rho(z_0)} \cap \Gamma(u)$. We can argue as in the beginning of Section 6 to deduce that over another subsequence

$$(7.2) \quad u_{\bar{z}_j,r_j} \rightarrow w \quad \text{in } C^{1,0}(\overline{Q_1^\pm})$$

for some $w \in C^{1,0}(\overline{Q_1^\pm})$. Moreover, we have by Lemma 6.2 that for any $s \in (-1, 0)$

$$u_{\bar{z}_j,r_j} - u_{\bar{z}_j,0} \rightarrow 0 \quad \text{in } L^1(B_1 \times (-1, s)),$$

which implies by using Cantor's diagonal argument

$$u_{\bar{z}_j,r_j} - u_{\bar{z}_j,0} \rightarrow 0 \quad \text{a.e. in } Q_1.$$

On the other hand, from Lemma 7.7, we find

$$u_{\bar{z}_j,0} \rightarrow u_{\bar{z}_0,0} \quad \text{in } C^{1,0}(\overline{Q_1^\pm}).$$

The previous two convergences, combined with (7.2), imply $w = u_{\bar{z}_0,0}$ and contradict our assumption.

Next, for a given $\varepsilon > 0$ and a unit vector $e \in \mathbb{R}^{n-1}$, define the cone

$$\mathcal{C}_\varepsilon(e) = \{x' \in \mathbb{R}^{n-1} : x' \cdot e > \varepsilon|x'|\}.$$

By utilizing Lemma 7.7, the estimate (7.1) and the complementarity condition (Lemma 2.2), we can follow Steps 2-3 in the proof of [JP21, Theorem 9.7] to obtain the following: for any $\varepsilon > 0$, there is $r_\varepsilon > 0$ such that for any $\bar{z} = (\bar{x}, \bar{t}) \in Q'_\rho(z_0) \cap \Gamma(u)$, we have

$$\begin{aligned} \bar{x} + (\mathcal{C}_\varepsilon(e_{\bar{z}}) \cap B'_{r_\varepsilon}) &\subset \{u(\cdot, 0, \bar{t}) > 0\}, \\ \bar{x} - (\mathcal{C}_\varepsilon(e_{\bar{z}}) \cap B'_{r_\varepsilon}) &\subset \{u(\cdot, 0, \bar{t}) = 0\}. \end{aligned}$$

Finally, by using these inclusions and Lemma 7.7, we can repeat the arguments in Steps 4-5 in [JP21, Theorem 9.7] to conclude the theorem. \square

APPENDIX A. EXISTENCE OF WEAK SOLUTIONS

In this section, we prove the existence and uniqueness of the weak solution to the parabolic Signorini problem in S_1 , provided that the initial datum belongs to W_∞^2 .

Theorem A.1. *If $\varphi_0 \in W_\infty^2(\mathbb{R}^n)$, then there exists a unique weak solution of*

$$(A.1) \quad \begin{cases} \partial_t v - \Delta v = 0 & \text{in } S_1^+ \cup S_1^-, \\ v \geq 0, \quad \partial_{\nu^+} v + \partial_{\nu^-} v \geq 0, \quad v(\partial_{\nu^+} v + \partial_{\nu^-} v) = 0 & \text{on } S_1', \\ v(\cdot, -1) = \varphi_0 & \text{on } \mathbb{R}^n, \end{cases}$$

where ν^\pm is the outer unit normal to S_1^\pm on S_1' .

Proof. For the change of coordinates

$$\tilde{v}(y, \tau) := v(2e^{-\frac{\tau}{2}}y, -e^{-\tau}), \quad (y, \tau) \in \mathbb{R}^n \times [0, \infty),$$

(A.1) is equivalent to

$$\begin{cases} \partial_\tau \tilde{v} + \frac{y}{2} \cdot \nabla \tilde{v} - \frac{1}{4} \Delta \tilde{v} = 0 & \text{in } (\mathbb{R}_+^n \cup \mathbb{R}_-^n) \times (0, \infty), \\ \tilde{v} \geq 0, \quad \partial_{\nu^+} \tilde{v} + \partial_{\nu^-} \tilde{v} \geq 0, \quad \tilde{v}(\partial_{\nu^+} \tilde{v} + \partial_{\nu^-} \tilde{v}) = 0 & \text{on } \mathbb{R}^{n-1} \times (0, \infty), \\ \tilde{v}(\cdot, 0) = \tilde{\varphi}_0 & \text{on } \mathbb{R}^n, \end{cases}$$

where $\tilde{\varphi}_0(y) = \varphi_0(2y)$. Note that \tilde{v} is a weak solution of the above equation if and only if it satisfies for a.e. $\tau \in (0, \infty)$ the variational inequality

$$\int_{\mathbb{R}^n} \partial_\tau \tilde{v}(w - \tilde{v})e^{-|y|^2} + \frac{y}{2} \cdot \nabla \tilde{v}(w - \tilde{v})e^{-|y|^2} + \frac{1}{4} \nabla \tilde{v} \cdot \nabla \left((w - \tilde{v})e^{-|y|^2} \right) \geq 0,$$

which is equivalent to

$$\int_{\mathbb{R}^n} \partial_\tau \tilde{v}(w - \tilde{v})e^{-|y|^2} + \frac{1}{4} \nabla \tilde{v} \cdot \nabla (w - \tilde{v})e^{-|y|^2} \geq 0,$$

for any $w \in L^2(0, \infty; W^{1,2}(\mathbb{R}^n, e^{-|y|^2}))$ with $w = \tilde{\varphi}_0$ on $\mathbb{R}^n \times \{0\}$, $w \geq 0$ on $\mathbb{R}^{n-1} \times (0, \infty)$ and $w - \tilde{v} \in L^2(0, \infty; W_0^{1,2}(\mathbb{R}^n, e^{-|y|^2}))$. In addition, for $a(v, v) := \frac{1}{4} \int_{\mathbb{R}^n} \nabla v \cdot \nabla v e^{-|y|^2} dy$, the coercivity

$$a(v, v) + C \int_{\mathbb{R}^n} v^2 e^{-|y|^2} \geq \alpha \int_{\mathbb{R}^n} (|\nabla v|^2 + v^2) e^{-|y|^2}$$

is satisfied. Therefore, the existence and the uniqueness of the weak solution \tilde{v} follow from [DL76, Chapter 1, Theorem 5.1]. \square

APPENDIX B. EXAMPLES OF ALMOST MINIMIZERS

In this section, we provide examples of solutions to certain equations that satisfy almost parabolic Signorini properties, both the unweighted and the weighted versions. These examples rely on the following technical lemma. For $\varepsilon \in (0, 1)$, we write $Q_{r,\rho}^\varepsilon(z_0) := B_{r^\varepsilon}(x_0) \times (t_0 - r^2, t_0 - \rho^2]$.

Lemma B.1. *For $\varepsilon = 1/3$ and a point $z_0 = (x_0, t_0) \in Q'_1$, suppose that a function $u \in W_2^{1,1}(Q_1) \cap L^2(-1, t_0; W^{1,2}(B_1, G_{z_0}))$ satisfies the following property: for any $Q_{r,\rho}^\varepsilon(z_0) \Subset Q_{1/2}$, and $v \in L^2(t_0 - r^2, t_0 - \rho^2; W^{1,2}(B_{r^\varepsilon}(x_0), G_{z_0}))$ with $v \geq 0$ on $Q_{r,\rho}^\varepsilon(z_0) \cap Q'_1$ and $v - u \in L^2(t_0 - r^2, t_0 - \rho^2; W_0^{1,2}(B_{r^\varepsilon}(x_0), G_{z_0}))$*

(B.1)

$$\begin{aligned} & \int_{Q_{r,\rho}^\varepsilon(z_0)} \left((1 - Cr^{\varepsilon\alpha})(t_0 - t)|\nabla u|^2 + ((x_0 - x) \cdot \nabla u + 2(t_0 - t)\partial_t u)(u - v) \right) G_{z_0} \\ & \leq \int_{Q_{r,\rho}^\varepsilon(z_0)} \left((1 + Cr^{\varepsilon\alpha})(t_0 - t)|\nabla v|^2 + Cr^{\varepsilon\alpha} \frac{|x_0 - x|^2}{t_0 - t} (u - v)^2 \right) G_{z_0}, \end{aligned}$$

where $C > 0$ are constants, independent of z_0 , ρ and r . Let $\psi \in C_0^\infty(\mathbb{R}^n)$ be a cutoff function satisfying

$$0 \leq \psi \leq 1, \quad \psi = 1 \quad \text{on } B_{1/2}, \quad \text{supp } \psi \subset B_1.$$

Then there exists a constant $r_0 > 0$ such that $\tilde{u} := u\psi$ satisfies the weighted almost parabolic Signorini property (1.2) at z_0 for $0 \leq \rho < r < r_0$, with a gauge function $\eta(r) = Cr^{\alpha/3}$.

Remark B.2. Since our main objective in this paper is the free boundary $\Gamma(u)$, in Lemma B.1, we can make the assumption that $\|u\|_{W_2^{1,1}(Q_{1/2})} > 0$. Otherwise, we have $u \equiv 0$ in $Q_{1/2}$ and there is no free boundary on $Q'_{1/2}$. Moreover, the condition (B.1) only concerns u within $Q_{r,\rho}^\varepsilon(z_0)$ and $Q_{r,\rho}^\varepsilon(z_0) \subset Q_{1/2}$, which allows us to freely modify the value of u in $Q_1 \setminus Q_{1/2}$. Therefore, we may assume that for some dimensional constant $C > 0$

$$\|u\|_{W_2^{1,1}(Q_1)} \leq C \|u\|_{W_2^{1,1}(Q_{1/2})}.$$

Proof. Step 1. Without loss of generality, we may assume $z_0 = 0$. (B.1) can be rewritten as

$$(B.2) \quad (1 - Cr^{\varepsilon\alpha})I + II \leq (1 + Cr^{\varepsilon\alpha})III + Cr^{\varepsilon\alpha}IV,$$

where

$$\begin{aligned} I &= \int_{Q_{r,\rho}^\varepsilon} (-t)|\nabla u|^2 G, & II &= \int_{Q_{r,\rho}^\varepsilon} (-x \cdot \nabla u - 2t\partial_t u)(u - v)G, \\ III &= \int_{Q_{r,\rho}^\varepsilon} (-t)|\nabla v|^2 G, & IV &= \int_{Q_{r,\rho}^\varepsilon} \frac{|x|^2}{-t}(u - v)^2 G. \end{aligned}$$

For $0 \leq \rho < r < 1$, let $w \in L^2(-r^2, -\rho^2; W^{1,2}(\mathbb{R}^n, G))$ with $w \geq 0$ on $S'_r \setminus S'_\rho$ and $\tilde{u} - w \in L^2(-r^2, -\rho^2; W_0^{1,2}(\mathbb{R}^n, G))$. By approximation, we may assume that w has a bounded support. We consider dilations of ψ

$$\psi_r(x) = \psi_{r,\varepsilon}(x) := \psi\left(\frac{x}{r^\varepsilon}\right),$$

and define

$$(B.3) \quad v(x, t) := u(x, t) + \psi_r(x)(w(x, t) - \tilde{u}(x, t)), \quad (x, t) \in B_{r^\varepsilon} \times (-r^2, -\rho^2).$$

Then $v - u \in L^2(-r^2, -\rho^2; W_0^{1,2}(B_{r^\varepsilon}, G))$ and $v = u + w - \tilde{u} = w \geq 0$ on $Q_{r,\rho}^\varepsilon \cap Q'_1$. Thus v is a valid competitor for u , and hence (B.2) holds for such v . In the below we estimate and rewrite I , II , III and IV in terms of \tilde{u} and w .

Step 2. We first deal with I . We compute

$$\begin{aligned} (B.4) \quad & \int_{S_r \setminus S_\rho} (-t)|\nabla \tilde{u}|^2 G = \int_{B_1 \times (-r^2, -\rho^2)} (-t)|\psi \nabla u + u \nabla \psi|^2 G \\ &= \int_{B_1 \times (-r^2, -\rho^2)} (-t)|\nabla u|^2 G \\ & \quad + \int_{(B_1 \setminus B_{1/2}) \times (-r^2, -\rho^2)} (-t) \left((\psi^2 - 1)|\nabla u|^2 + 2u\psi \nabla u \cdot \nabla \psi + u^2 |\nabla \psi|^2 \right) G \\ &= \int_{Q_{r,\rho}^\varepsilon} (-t)|\nabla u|^2 G + \int_{(B_1 \setminus B_{r^\varepsilon}) \times (-r^2, -\rho^2)} (-t)|\nabla u|^2 G \\ & \quad + \int_{(B_1 \setminus B_{1/2}) \times (-r^2, -\rho^2)} (-t) \left((\psi^2 - 1)|\nabla u|^2 + 2u\psi \nabla u \cdot \nabla \psi + u^2 |\nabla \psi|^2 \right) G. \end{aligned}$$

To estimate the last two terms, we claim that for $t \in (-r^2, -\rho^2)$ with $r < r_0 = r_0(n)$ small, we have

$$(B.5) \quad G(x, t) \leq e^{\frac{1}{17t}} \text{ for } |x| \geq 1/2 \quad \text{and} \quad G(x, t) \leq e^{\frac{1}{32t}r^{2\varepsilon}} \text{ for } |x| \geq \frac{1}{2}r^\varepsilon.$$

Indeed, if $|x| \geq 1/2$, then we easily have

$$G(x, t) \leq \frac{e^{\frac{|x|^2}{4t}}}{(-t)^{n/2}} \leq \frac{e^{\frac{1}{16t}}}{(-t)^{n/2}} \leq e^{\frac{1}{17t}},$$

which gives the first estimate. For the second one, we define

$$\zeta_r(s) := \frac{e^{-\frac{s^{2\varepsilon}}{32s}}}{s^{n/2}}, \quad 0 < s \leq r^2.$$

Then $\zeta_r(r^2) = \frac{e^{-\frac{1}{32r^{2-2\varepsilon}}}}{r^n} < 1$ and $\frac{d}{ds}\zeta_r(s) = \frac{ne^{-\frac{s^{2\varepsilon}}{32s}}}{2s^{\frac{n}{2}+2}} \left(\frac{r^{2\varepsilon}}{16n} - s \right) \geq 0$, $0 < s \leq r^2$, which gives $\zeta_r(s) < 1$ for $0 < s \leq r^2$. Thus, if $|x| \geq \frac{1}{2}r^\varepsilon$ and $t \in (-r^2, 0)$, then

$$G(x, t) \leq \frac{e^{\frac{1}{16t}r^{2\varepsilon}}}{(-t)^{n/2}} = \zeta_r(-t)e^{\frac{1}{32t}r^{2\varepsilon}} \leq e^{\frac{1}{32t}r^{2\varepsilon}}.$$

By using the claim (B.5), we obtain

$$\begin{aligned} & \left| \int_{(B_1 \setminus B_{1/2}) \times (-r^2, -\rho^2)} (-t) \left((\psi^2 - 1) |\nabla u|^2 + 2u\psi \nabla u \cdot \nabla \psi + u^2 |\nabla \psi|^2 \right) G \right| \\ & \leq C(n) \int_{(B_1 \setminus B_{1/2}) \times (-r^2, -\rho^2)} (|\nabla u|^2 + u^2) G \\ & \leq C(n) \|u\|_{W_2^{1,1}(Q_1)}^2 e^{-\frac{1}{17r^2}}, \end{aligned}$$

and

$$\int_{(B_1 \setminus B_{r^\varepsilon}) \times (-r^2, -\rho^2)} (-t) |\nabla u|^2 G \leq C(n) \|u\|_{W_2^{1,1}(Q_1)}^2 e^{-\frac{1}{32r^{2-2\varepsilon}}}.$$

Combining these estimates with (B.4) yields

$$I \geq \int_{S_r \setminus S_\rho} (-t) |\nabla \tilde{u}|^2 G - C(n) \|u\|_{W_2^{1,1}(Q_1)}^2 e^{-\frac{1}{32r^{2-2\varepsilon}}}.$$

Step 3. To estimate II , we observe that $u = \tilde{u}$ in $Q_{r,\rho}^\varepsilon$, $\tilde{u} = 0$ in $(\mathbb{R}^n \setminus B_1) \times (-r^2, -\rho^2)$, and

$$(B.6) \quad r^{-\varepsilon} G(x, t) \leq r^{-\varepsilon} e^{\frac{1}{32t}r^{2\varepsilon}} \leq r^{-\varepsilon} e^{-\frac{1}{16r^{2-2\varepsilon}}} \leq e^{-\frac{1}{32r^{2-2\varepsilon}}}$$

for $(x, t) \in (B_1 \setminus B_{\frac{1}{2}r^\varepsilon}) \times (-r^2, -\rho^2)$ with $r < r_0$ small. By using (B.3), (B.6) and Young's inequality, we deduce

$$\begin{aligned} II &= \int_{Q_{r,\rho}^\varepsilon} \psi_r(-x \cdot \nabla \tilde{u} - 2t\partial_t \tilde{u})(\tilde{u} - w)G \\ &= \int_{Q_{r,\rho}^\varepsilon} (-x \cdot \nabla \tilde{u} - 2t\partial_t \tilde{u})(\tilde{u} - w)G \\ &\quad + \int_{(B_{r^\varepsilon} \setminus B_{\frac{1}{2}r^\varepsilon}) \times (-r^2, -\rho^2)} (\psi_r - 1)(-x \cdot \nabla \tilde{u} - 2t\partial_t \tilde{u})(\tilde{u} - w)G \end{aligned}$$

$$\begin{aligned}
&= \int_{S_r \setminus S_\rho} (-x \cdot \nabla \tilde{u} - 2t \partial_t \tilde{u})(\tilde{u} - w)G \\
&\quad - \int_{(B_1 \setminus B_{r^\varepsilon}) \times (-r^2, -\rho^2)} (-x \cdot \nabla \tilde{u} - 2t \partial_t \tilde{u})(\tilde{u} - w)G \\
&\quad + \int_{(B_{r^\varepsilon} \setminus B_{\frac{1}{2}r^\varepsilon}) \times (-r^2, -\rho^2)} (\psi_r - 1)(-x \cdot \nabla \tilde{u} - 2t \partial_t \tilde{u})(\tilde{u} - w)G \\
&\geq \int_{S_r \setminus S_\rho} (-x \cdot \nabla \tilde{u} - 2t \partial_t \tilde{u})(\tilde{u} - w)G - r^\varepsilon \int_{(B_1 \setminus B_{\frac{1}{2}r^\varepsilon}) \times (-r^2, -\rho^2)} (\tilde{u} - w)^2 G \\
&\quad - r^{-\varepsilon} \int_{(B_1 \setminus B_{\frac{1}{2}r^\varepsilon}) \times (-r^2, -\rho^2)} (-x \cdot \nabla \tilde{u} - 2t \partial_t \tilde{u})^2 G \\
&\geq \int_{S_r \setminus S_\rho} (-x \cdot \nabla \tilde{u} - 2t \partial_t \tilde{u})(\tilde{u} - w)G - r^\varepsilon \int_{S_r \setminus S_\rho} (\tilde{u} - w)^2 G \\
&\quad - \|u\|_{W_2^{1,1}(Q_1)}^2 e^{-\frac{1}{33r^2-2\varepsilon}}.
\end{aligned}$$

Step 4. Before we estimate *III*, we prove

$$\begin{aligned}
\text{(B.7)} \quad &\int_{(B_{r^\varepsilon} \setminus B_{\frac{1}{2}r^\varepsilon}) \times (-r^2, -\rho^2)} |\nabla \psi_r|^2 (w - \tilde{u})^2 G \\
&\leq C(n)r^{4-6\varepsilon} \int_{S_r \setminus S_\rho} (w - \tilde{u})^2 G + C(n)r^{2-4\varepsilon} \int_{S_r \setminus S_\rho} (-t)(|\nabla w|^2 + |\nabla \tilde{u}|^2)G.
\end{aligned}$$

To this end, we apply the Log-Sobolev Inequality. [EP08, Lemma 1.2] can be rewritten as (by letting $g(y) := f\left(\frac{y}{\sqrt{-2t}}\right)$)

$$\log\left(\frac{1}{\mathcal{A}}\right) \int_{\mathbb{R}^n} g^2 G(\cdot, t) \leq -4t \int_{\mathbb{R}^n} |\nabla g|^2 G(\cdot, t), \quad \text{where } \mathcal{A} := \int_{\{|g|>0\}} G(\cdot, t),$$

for every $t < 0$ and $g \in W_2^{1,0}(\mathbb{R}^n, G(\cdot, t))$. We plug in $g = \partial_{x_i} \psi_r (w(\cdot, t) - \tilde{u}(\cdot, t))$, $1 \leq i \leq n$, for each $t \in (-r^2, -\rho^2)$. Then, by using

$$\mathcal{A} \leq \int_{B_{r^\varepsilon} \setminus B_{\frac{1}{2}r^\varepsilon}} G(x, t) dx \leq e^{\frac{1}{32t}r^{2\varepsilon}} \int_{B_{r^\varepsilon} \setminus B_{\frac{1}{2}r^\varepsilon}} dx \leq e^{\frac{1}{32t}r^{2\varepsilon}},$$

where the second inequality holds due to (B.5), we have

$$\begin{aligned}
&\int_{\mathbb{R}^n \times \{t\}} |\partial_{x_i} \psi_r|^2 (w - \tilde{u})^2 G \\
&\leq \frac{128t^2}{r^{2\varepsilon}} \int_{\mathbb{R}^n \times \{t\}} |\nabla(\partial_{x_i} \psi_r)(w - \tilde{u}) + \partial_{x_i} \psi_r \nabla(w - \tilde{u})|^2 G \\
&\leq \frac{C(n)t^2}{r^{2\varepsilon}} \left(\frac{1}{r^{4\varepsilon}} \int_{\mathbb{R}^n \times \{t\}} (w - \tilde{u})^2 G + \frac{1}{r^{2\varepsilon}} \int_{\mathbb{R}^n \times \{t\}} (|\nabla w|^2 + |\nabla \tilde{u}|^2)G \right) \\
&= \frac{C(n)t^2}{r^{6\varepsilon}} \int_{\mathbb{R}^n \times \{t\}} (w - \tilde{u})^2 G + \frac{C(n)(-t)}{r^{4\varepsilon}} \int_{\mathbb{R}^n \times \{t\}} (-t)(|\nabla w|^2 + |\nabla \tilde{u}|^2)G.
\end{aligned}$$

By integrating in $t \in (-r^2, -\rho^2)$ and summing for $1 \leq i \leq n$, we derive (B.7).

Now, by using (B.3), (B.6), (B.7) and Young's inequality, we have

$$\begin{aligned}
 III &= \int_{Q_{r,\rho}^\varepsilon} (-t)|\psi_r \nabla w + \nabla \psi_r(w - \tilde{u}) + \nabla u - \psi_r \nabla \tilde{u}|^2 G \\
 &\leq (1 + r^\varepsilon) \int_{Q_{r,\rho}^\varepsilon} (-t)\psi_r^2 |\nabla w|^2 G \\
 &\quad + \left(1 + \frac{1}{r^\varepsilon}\right) \int_{(B_{r^\varepsilon} \setminus B_{\frac{1}{2}r^\varepsilon}) \times (-r^2, -\rho^2)} |\nabla \psi_r(w - \tilde{u}) + (1 - \psi_r)\nabla u|^2 G \\
 &\leq (1 + r^\varepsilon) \int_{S_r \setminus S_\rho} (-t)|\nabla w|^2 G \\
 &\quad + \frac{4}{r^\varepsilon} \int_{(B_{r^\varepsilon} \setminus B_{\frac{1}{2}r^\varepsilon}) \times (-r^2, -\rho^2)} (|\nabla \psi_r(w - \tilde{u})|^2 + |\nabla u|^2) G \\
 &\leq (1 + r^\varepsilon + Cr^{2-5\varepsilon}) \int_{S_r \setminus S_\rho} (-t)|\nabla w|^2 G + Cr^{2-5\varepsilon} \int_{S_r \setminus S_\rho} (-t)|\nabla \tilde{u}|^2 G \\
 &\quad + Cr^{4-7\varepsilon} \int_{S_r \setminus S_\rho} (w - \tilde{u})^2 G + C\|u\|_{W_2^{1,1}(Q_1)}^2 e^{-\frac{1}{33r^{2-2\varepsilon}}}.
 \end{aligned}$$

Step 5. It remains to consider *IV*. By using the equality $\nabla G = \frac{x}{2t}G$ and applying the integrations by parts and Young's inequality, we get

$$\begin{aligned}
 &\int_{S_r \setminus S_\rho} \frac{|x|^2}{-t} (\tilde{u} - w)^2 G \\
 &= \int_{S_r \setminus S_\rho} -2x(\tilde{u} - w)^2 \nabla G = \int_{S_r \setminus S_\rho} [2n(\tilde{u} - w)^2 + 4x \cdot \nabla(\tilde{u} - w)(\tilde{u} - w)] G \\
 &\leq 2n \int_{S_r \setminus S_\rho} (\tilde{u} - w)^2 G + \frac{1}{2} \int_{S_r \setminus S_\rho} \frac{|x|^2}{-t} (\tilde{u} - w)^2 G + C \int_{S_r \setminus S_\rho} (-t)|\nabla(\tilde{u} - w)|^2 G.
 \end{aligned}$$

This gives

$$IV \leq \int_{S_r \setminus S_\rho} \frac{|x|^2}{-t} (\tilde{u} - w)^2 G \leq C \int_{S_r \setminus S_\rho} (\tilde{u} - w)^2 G + C \int_{S_r \setminus S_\rho} (-t)|\nabla(\tilde{u} - w)|^2 G.$$

Step 6. By combining (B.2) with the estimates for *I-IV* and recalling $\varepsilon = 1/3$, we obtain

$$\begin{aligned}
 &\int_{S_r \setminus S_\rho} \left[(1 - Cr^{\alpha/3})(-t)|\nabla \tilde{u}|^2 + (-x \cdot \nabla \tilde{u} - 2t\partial_t \tilde{u})(\tilde{u} - w) \right] G \\
 &\leq \int_{S_r \setminus S_\rho} \left[(1 + Cr^{\alpha/3})(-t)|\nabla w|^2 + Cr^{\alpha/3}(\tilde{u} - w)^2 \right] G + C\|u\|_{W_2^{1,1}(Q_1)} e^{-\frac{1}{34r^{4/3}}}.
 \end{aligned}$$

Finally, since $C\|u\|_{W_2^{1,1}(Q_1)} \leq C\|u\|_{W_2^{1,1}(Q_{1/2})} \leq C\|\tilde{u}\|_{W_2^{1,1}(Q_1)}$ by Remark B.2, we have for small $r_0 > 0$

$$C\|u\|_{W_2^{1,1}(Q_1)} e^{-\frac{1}{34r^{4/3}}} \leq \|\tilde{u}\|_{W_2^{1,1}(Q_1)} e^{-\frac{1}{r}}, \quad 0 < r < r_0.$$

This completes the proof. \square

Now we are ready to introduce some examples of almost minimizers, with the help of Lemma B.1.

Example B.3. Given $0 < \alpha < 1$, let A be a variable coefficient matrix satisfying $|A(x, t) - I| \leq C(|x|^2 + |t|)^{\alpha/2}$. Let $u \in W_2^{1,1}(Q_1)$ be a solution of the parabolic A -Signorini problem in Q_1

$$\begin{aligned} -\operatorname{div}(A\nabla u) + \partial_t u &= 0 \quad \text{in } Q_1^\pm, \\ u &\geq 0, \quad \langle A\nabla u, \nu^+ \rangle + \langle A\nabla u, \nu^- \rangle \geq 0, \\ u(\langle A\nabla u, \nu^+ \rangle + \langle A\nabla u, \nu^- \rangle) &= 0 \quad \text{on } Q_1', \end{aligned}$$

where $\nu^\pm = \mp e_n$. We interpret this in the weak sense that u satisfies for a.e. $t \in (-1, 0)$ the variational inequality

$$(B.8) \quad \int_{B_1} \langle A\nabla u, \nabla(u-w) \rangle + \partial_t u(u-w) \leq 0,$$

for any $w \in W^{1,2}(B_1)$ with $w = u$ on ∂B_1 and $w \geq 0$ on B_1' . Then

- (i) u satisfies the unweighted almost parabolic Signorini property at 0 with a gauge function $\eta(r) = Cr^\alpha$.
- (ii) $\tilde{u} = u\psi$ satisfies the weighted almost parabolic Signorini property at 0 with a gauge function $\eta(r) = Cr^{\alpha/3}$.

Proof. We first treat (i). For any $0 < r < 1$, let $w \in W_2^{1,0}(Q_r)$ be such that $w = u$ on $\partial_p Q_r$ and $w \geq 0$ on Q_r' . By extending $w = u$ in $Q_1 \setminus Q_r$, we get from (B.8) that

$$\int_{Q_r} \langle A\nabla u, \nabla(u-w) \rangle + \partial_t u(u-w) \leq 0.$$

Thus

$$\begin{aligned} &\int_{Q_r} |\nabla u|^2 + \partial_t u(u-w) \\ &= \int_{Q_r} \langle A\nabla u, \nabla(u-w) \rangle + \partial_t u(u-w) \\ &\quad + \int_{Q_r} \langle \nabla u, \nabla w \rangle + \langle (A-I)\nabla u, \nabla w \rangle + \langle (I-A)\nabla u, \nabla u \rangle \\ &\leq \frac{1}{2} \int_{Q_r} (|\nabla u|^2 + |\nabla w|^2) + \frac{1}{2} \int_{Q_r} (r^{-\alpha}|(A-I)\nabla u|^2 + r^\alpha|\nabla w|^2) \\ &\quad + \frac{1}{2} \int_{Q_r} (r^{-\alpha}|(I-A)\nabla u|^2 + r^\alpha|\nabla u|^2) \\ &\leq \frac{1+Cr^\alpha}{2} \int_{Q_r} (|\nabla u|^2 + |\nabla w|^2). \end{aligned}$$

This gives the unweighted almost parabolic Signorini property of u at 0.

To prove the weighted property (ii), we observe that u also satisfies for a.e. $t \in (-1, 0)$ the following variational inequality

$$(B.9) \quad \int_{B_1} [(-2t)\langle A\nabla u, \nabla(u-v) \rangle - \langle x, A\nabla u \rangle(u-v) + (-2t)\partial_t u(u-v)]G(\cdot, t) \leq 0,$$

for any competitor $v \in W_2^{1,0}(B_1, G)$ with $v = u$ on ∂B_1 and $v \geq 0$ on B_1' . In fact, this follows by inserting $w = u + (v-u)e^{\frac{|x|^2}{4t}}$ in (B.8) and multiplying $\frac{-2t}{(-4\pi t)^{n/2}}$ in both sides. To prove (B.1) for $z_0 = 0$, which readily implies (ii) by Lemma B.1,

we fix $\varepsilon = 1/3$. Then, for any $0 \leq \rho < r < 1$ and $v \in W_2^{1,0}(Q_{r,\rho}^\varepsilon, G)$ such that $v - u \in L^2(-r^2, -\rho^2; W_0^{1,2}(B_{r^\varepsilon}))$ and $v \geq 0$ on $Q_{r,\rho}^\varepsilon \cap Q'_1$, we extend $v = u$ in $(B_1 \setminus B_{r^\varepsilon}) \times (-r^2, -\rho^2)$ and use (B.9) to obtain

$$\int_{Q_{r,\rho}^\varepsilon} (-2t) \langle A \nabla u, \nabla(u-v) \rangle G - \langle x, A \nabla u \rangle (u-v)G + (-2t) \partial_t u (u-v)G \leq 0.$$

Using $2 \nabla u \cdot \nabla(u-v) \geq |\nabla u|^2 - |\nabla v|^2$, $|\nabla u \cdot \nabla(u-v)| \leq 3/2 |\nabla u|^2 + |\nabla v|^2$, and $|A - I| \leq Cr^{\varepsilon\alpha}$ in $Q_{r,\rho}^\varepsilon$, we get

$$\begin{aligned} & \int_{Q_{r,\rho}^\varepsilon} (-2t) \langle A \nabla u, \nabla(u-v) \rangle G \\ &= \int_{Q_{r,\rho}^\varepsilon} (-2t) \nabla u \cdot \nabla(u-v)G + \int_{Q_{r,\rho}^\varepsilon} (-2t) \langle (A - I) \nabla u, \nabla(u-v) \rangle G \\ &\geq (1 - Cr^{\varepsilon\alpha}) \int_{Q_{r,\rho}^\varepsilon} (-t) |\nabla u|^2 G - (1 + Cr^{\varepsilon\alpha}) \int_{Q_{r,\rho}^\varepsilon} (-t) |\nabla v|^2 G. \end{aligned}$$

Combining the above two estimates yields

$$\begin{aligned} & (1 - Cr^{\varepsilon\alpha}) \int_{Q_{r,\rho}^\varepsilon} (-t) |\nabla u|^2 G + \int_{Q_{r,\rho}^\varepsilon} (-x \cdot \nabla u - 2t \partial_t u) (u-v)G \\ &\leq (1 + Cr^{\varepsilon\alpha}) \int_{Q_{r,\rho}^\varepsilon} (-t) |\nabla v|^2 G + \int_{Q_{r,\rho}^\varepsilon} \langle x, (A - I) \nabla u \rangle (u-v)G. \end{aligned}$$

Finally, by estimating the last term with Young's inequality

$$\int_{Q_{r,\rho}^\varepsilon} \langle x, (A - I) \nabla u \rangle (u-v)G \leq Cr^{\varepsilon\alpha} \int_{Q_{r,\rho}^\varepsilon} (-t) |\nabla u|^2 G + \frac{|x|^2}{-t} (u-v)^2 G,$$

we conclude (B.1) for $z_0 = 0$. \square

Example B.4. Let u be a solution of the parabolic Signorini problem for the Laplacian with drift with the velocity field $b \in L^\infty(-1, 0; L^p(B_1))$, $p > n$:

$$\begin{aligned} -\Delta u + b(x, t) \cdot \nabla u + \partial_t u &= 0 \quad \text{in } Q_1^\pm \\ -\partial_{x_n} u \geq 0, \quad u \geq 0 \quad u \partial_{x_n} u &= 0 \quad \text{on } Q'_1, \end{aligned}$$

even in x_n -variable. We understand this in the weak sense that u satisfies the variational inequality: for any $-1 < t < 0$,

$$\int_{B_1 \times \{t\}} \nabla u \cdot \nabla(v-u) + b(x, t) \cdot \nabla u (v-u) + \partial_t u (v-u) \geq 0,$$

for any competitor $v \in W^{1,2}(B_1)$ such that $v \geq 0$ on B'_1 and $v = u$ on ∂B_1 . Then

- (i) u is an unweighted almost minimizer for the parabolic Signorini problem in Q_1 with a gauge function $\eta(r) = Cr^{1-n/p}$.
- (ii) $\tilde{u} = u\psi$ is a weighted almost minimizer for the parabolic Signorini problem in Q'_1 with a gauge function $\eta(r) = Cr^{\frac{1}{3}(1-n/p)}$.

Proof. Since (i) is proved in [JP21, Example A.1] for more general case with variable coefficients, it is sufficient to prove (ii). For this purpose, as in Example B.3, we prove (B.1) for every $z_0 \in Q'_1$. Indeed, without loss of generality we may assume

that $z_0 = 0$. By the similar argument as in Example B.3, u also satisfies for a.e. $t \in (-1, 0)$ the variational inequality

$$\int_{B_1 \times \{-t\}} [(-2t)\nabla u \cdot \nabla(u-v) + (-x \cdot \nabla u - 2t\partial_t u)(u-v) + (-2t)b \cdot \nabla u(u-v)] G \leq 0$$

for any $v \in W^{1,2}(B_1, G(\cdot, t))$ with $v = u$ on ∂B_1 and $v \geq 0$ on B_1' . For $\varepsilon = 1/3$ and $0 \leq \rho < r < 1$, let $v \in W_2^{1,0}(Q_{r,\rho}^\varepsilon, G)$ be such that $v-u \in L^2(-r^2, -\rho^2; W_0^{1,2}(B_{r^\varepsilon}, G))$ and $v \geq 0$ on $Q_{r,\rho}^\varepsilon \cap Q_1'$. Extending v to $B_1 \times (-r^2, -\rho^2)$ by $v = u$ on $(B_1 \setminus B_{r^\varepsilon}) \times (-r^2, -\rho^2)$ and using the above variational inequality, we get

$$\begin{aligned} & \int_{Q_{r,\rho}^\varepsilon} ((-t)|\nabla u|^2 + (-x \cdot \nabla u - 2t\partial_t u)(u-v)) G \\ & \leq \int_{Q_{r,\rho}^\varepsilon} ((-t)|\nabla u|^2 + (-2t)\nabla u \cdot \nabla(v-u) + (-2t)b \cdot \nabla u(v-u)) G \\ & = \int_{Q_{r,\rho}^\varepsilon} (-(-t)|\nabla u|^2 + (-2t)\nabla u \cdot \nabla v) G + \int_{-r^2}^{-\rho^2} (-2t) \int_{B_{r^\varepsilon}} b \cdot \nabla u(v-u) G \, dx dt \\ & \leq \int_{Q_{r,\rho}^\varepsilon} (-t)|\nabla v|^2 G + \int_{-r^2}^{-\rho^2} (-2t) M \|\nabla u G^{1/2}\|_{L^2(B_{r^\varepsilon})} \|(v-u)G^{1/2}\|_{L^{p^*}(B_{r^\varepsilon})} dt, \end{aligned}$$

where $M := \sup\{\|b(\cdot, t)\|_{L^p(B_1)} : -1 < t < 0\}$ and $p^* = \frac{2p}{p-2}$. For $\gamma = 1 - n/p$, we have by Sobolev's inequality,

$$\begin{aligned} \|(v-u)G^{1/2}\|_{L^{p^*}(B_{r^\varepsilon})} & \leq C_{n,p} r^{\varepsilon\gamma} \|\nabla((v-u)G^{1/2})\|_{L^2(B_{r^\varepsilon})} \\ & \leq Cr^{\varepsilon\gamma} \left(\|\nabla(v-u)G^{1/2}\|_{L^2(B_{r^\varepsilon})} + \|(v-u)\frac{x}{t}G^{1/2}\|_{L^2(B_{r^\varepsilon})} \right). \end{aligned}$$

Thus

$$\begin{aligned} & \int_{-r^2}^{-\rho^2} (-2t) M \|\nabla u G^{1/2}\|_{L^2(B_{r^\varepsilon})} \|(v-u)G^{1/2}\|_{L^{p^*}(B_{r^\varepsilon})} dt \\ & \leq Cr^{\varepsilon\gamma} \int_{-r^2}^{-\rho^2} (-2t) \|\nabla u G^{1/2}\|_{L^2(B_{r^\varepsilon})} \left(\|\nabla(v-u)G^{1/2}\|_{L^2(B_{r^\varepsilon})} \right. \\ & \quad \left. + \|(v-u)\frac{x}{t}G^{1/2}\|_{L^2(B_{r^\varepsilon})} \right) dt \\ & \leq Cr^{\varepsilon\gamma} \int_{Q_{r,\rho}^\varepsilon} (-2t) (|\nabla u|^2 + |\nabla v|^2) G + Cr^{\varepsilon\gamma} \int_{Q_{r,\rho}^\varepsilon} \frac{|x|^2}{(-t)} (v-u)^2 G, \end{aligned}$$

where constants $C > 0$ depend only on n, p and M . Therefore,

$$\begin{aligned} & \int_{Q_{r,\rho}^\varepsilon} (1 - Cr^{\varepsilon\gamma}) (-t) |\nabla u|^2 G + (-x \cdot \nabla u - 2t\partial_t u)(u-v) G \\ & \leq \int_{Q_{r,\rho}^\varepsilon} (1 + Cr^{\varepsilon\gamma}) (-t) |\nabla v|^2 G + Cr^{\varepsilon\gamma} \frac{|x|^2}{(-t)} (v-u)^2 G. \end{aligned}$$

This completes the proof. \square

Disclosure statement. The authors report there are no competing interests to declare.

REFERENCES

- [AS24] Vedansh Arya and Wenhui Shi, *Optimal regularity for the variable coefficients parabolic Signorini problem* (2024), 67 pp, arXiv:2401.13305 preprint. [↑2](#)
- [ACS08] I. Athanasopoulos, L. A. Caffarelli, and S. Salsa, *The structure of the free boundary for lower dimensional obstacle problems*, Amer. J. Math. **130** (2008), no. 2, 485–498, doi:[10.1353/ajm.2008.0016](#). MR2405165 [↑2](#)
- [CSS08] Luis A. Caffarelli, Sandro Salsa, and Luis Silvestre, *Regularity estimates for the solution and the free boundary of the obstacle problem for the fractional Laplacian*, Invent. Math. **171** (2008), no. 2, 425–461, doi:[10.1007/s00222-007-0086-6](#). MR2367025 [↑2](#)
- [DGPT17] Donatella Danielli, Nicola Garofalo, Arshak Petrosyan, and Tung To, *Optimal regularity and the free boundary in the parabolic Signorini problem*, Mem. Amer. Math. Soc. **249** (2017), no. 1181, v + 103, doi:[10.1090/memo/1181](#). MR3709717 [↑2](#), 5, 6, 15, 20, 22, 26, 34
- [DL76] Georges Duvaut and Jacques-Louis Lions, *Inequalities in mechanics and physics*, Springer-Verlag, Berlin-New York, 1976. Translated from the French by C. W. John, Grundlehren der Mathematischen Wissenschaften, 219. MR0521262 [↑37](#)
- [EP08] Anders Edquist and Arshak Petrosyan, *A parabolic almost monotonicity formula*, Math. Ann. **341** (2008), no. 2, 429–454. MR2385663 [↑40](#)
- [GP09] Nicola Garofalo and Arshak Petrosyan, *Some new monotonicity formulas and the singular set in the lower dimensional obstacle problem*, Invent. Math. **177** (2009), no. 2, 415–461, doi:[10.1007/s00222-009-0188-4](#). MR2511747 [↑2](#)
- [GPS16] Nicola Garofalo, Arshak Petrosyan, and Mariana Smit Vega Garcia, *An epiperimetric inequality approach to the regularity of the free boundary in the Signorini problem with variable coefficients*, J. Math. Pures Appl. (9) **105** (2016), no. 6, 745–787, doi:[10.1016/j.matpur.2015.11.013](#) (English, with English and French summaries). MR3491531 [↑35](#)
- [JP21] Seongmin Jeon and Arshak Petrosyan, *Almost minimizers for the thin obstacle problem*, Calc. Var. Partial Differential Equations **60** (2021), Paper No. 124, 59, doi:[10.1007/s00526-021-01986-8](#). MR4277328 [↑4](#), 5, 7, 15, 22, 24, 28, 33, 36, 43
- [JP23] ———, *Regularity of almost minimizers for the parabolic thin obstacle problem*, Nonlinear Anal. **237** (2023), Paper No. 113386, 29, doi:[10.1016/j.na.2023.113386](#). MR4645649 [↑3](#), 4, 6, 7
- [JPS24] Seongmin Jeon, Arshak Petrosyan, and Mariana Smit Vega Garcia, *Almost minimizers for the thin obstacle problem with variable coefficients*, Interfaces Free Bound., posted on 2024, doi:[10.4171/IFB/507](#). [↑16](#)
- [PSU12] Arshak Petrosyan, Henrik Shahgholian, and Nina Uraltseva, *Regularity of free boundaries in obstacle-type problems*, Graduate Studies in Mathematics, vol. 136, American Mathematical Society, Providence, RI, 2012. MR2962060 [↑2](#)
- [Poo96] Chi-Cheung Poon, *Unique continuation for parabolic equations*, Comm. Partial Differential Equations **21** (1996), 521–539, doi:[10.1080/03605309608821195](#). MR1387458 [↑5](#), 15
- [Shi20] Wenhui Shi, *An epiperimetric inequality approach to the parabolic Signorini problem*, Disc. Cont. Dyn. Syst. **40** (2020), 1813–1846, doi:[10.3934/dcds.2020095](#). MR4063946 [↑5](#), 24
- [Wei99] Georg S. Weiss, *A homogeneity improvement approach to the obstacle problem*, Invent. Math. **138** (1999), no. 1, 23–50, doi:[10.1007/s002220050340](#). MR1714335 [↑7](#)

DEPARTMENT OF MATHEMATICS EDUCATION, HANYANG UNIVERSITY, 222 WANGSIMNI-RO, SEONGDONG-GU, SEOUL 04763, REPUBLIC OF KOREA
Email address: seongminjeon@hanyang.ac.kr

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907, USA
Email address: arshak@purdue.edu