

## 8.2 Integration By Parts

NOT on exam!

Hw includes 8.1 (Basic integration)

Integration by substitution: Chain rule in reverse

Integration by parts: Product rule in reverse

$u, v$  both functions of  $x$

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

in the differential form

$$d(uv) = u dv + v du$$

$$u dv = d(uv) - v du$$

integrate

$$\int u dv = \int d(uv) - \int v du$$

$$\int u dv = uv - \int v du$$

Integration by parts

---

$$\int u dv = uv - \int v du$$

Given integral, identify

$u$  and  $dv$ , then use

$$uv - \int v du$$

## Example

$$\int x \cdot \ln x \, dx$$

not something we can integrate directly

Substitution doesn't work either

now try by parts


identify  $u$ ,  $dv$

$$\int \underline{x} \cdot \underline{\ln x} \, dx \quad \text{two parts to consider: } x \text{ and } \ln x$$

$u$ : has a simpler derivative

$dv$ : easy to integrate

a good rule of thumb for order of choosing  $u$



Pick in that order

back to  $\int x \cdot \ln x \, dx$   
algebraic  $\swarrow$  logarithmic

L.I.A.T.E: pick  $u = \ln x$  since it's L  
which is before A ( $x$ )

$$u = \ln x$$

$dv = x \, dx \rightarrow$  any left over after choosing  $u$

$$\frac{du}{dx} = \frac{1}{x}$$

$v = \int dv$  integrate

$$du = \frac{1}{x} \, dx$$

$$v = \int x \, dx = \frac{1}{2} x^2$$

don't need  $+ C$

formula:  $\int u \, dv = uv - \int v \, du$

$$\begin{aligned} &= (\ln x) \left( \frac{1}{2} x^2 \right) - \int \left( \frac{1}{2} x^2 \right) \left( \frac{1}{x} \, dx \right) \\ &= \frac{1}{2} x^2 \ln x - \int \frac{1}{2} x \, dx = \frac{1}{2} x^2 \ln x - \frac{1}{4} x^2 + C \end{aligned}$$

to check: take derivative  
it must match  $x \ln x$

Example

$$\int x \cos x \, dx$$

parts:  $\underline{x}$  .  $\underline{\cos x}$  order for u: LIATE

$\swarrow$   
algebraic (A)  
 $\searrow$   
trig (T)

A before T so we choose  
 $u = x$

deriv.

$$\left\{ \begin{array}{l} u = x \\ du = dx \end{array} \right. \quad \left\{ \begin{array}{l} dv = \cos x \, dx \\ v = \sin x \end{array} \right.$$

} integrate  
+ c is not needed here

$$uv - \int v \, du$$

$$= x \sin x - \int \sin x \, dx = x \sin x - (-\cos x) + C$$

$$= \boxed{x \sin x + \cos x + C}$$

definite integral  $\int_0^{\pi/2} x \cos x \, dx$

pick u, dv as usual

then  $uv \Big|_0^{\pi/2} - \int_0^{\pi/2} v \, du = x \sin x \Big|_0^{\pi/2} - \int_0^{\pi/2} \sin x \, dx$

$$= \int_1^{\frac{\pi}{2}} \sin \frac{\pi}{2} + \cos x \Big|_0^{\frac{\pi}{2}} = \frac{\pi}{2} + \cos\left(\frac{\pi}{2}\right) - \cos(0) = \boxed{\frac{\pi}{2} - 1}$$


---

Example

$$\int_1^2 x^2 \cos x \, dx$$

L(ATE)

so  $u = x^2$  (A before T)

$$u = x^2 \quad dv = \cos x \, dx$$

$$du = 2x \, dx \quad v = \sin x$$

$$uv - \int v \, du$$

$$= x^2 \sin x - \int (\sin x)(2x) \, dx$$

$$= x^2 \sin x - 2 \int x \cdot \sin x \, dx$$

by parts again,  $x \rightarrow A$   $\sin x \rightarrow T$

$$U = x \quad dV = \sin x \, dx$$

$$dU = dx \quad V = -\cos x$$

$$UV - \int V \, dU$$

$$= x^2 \sin x - 2 (uv - \int v du)$$

$$= x^2 \sin x - 2 (-x \cos x - \int -\cos x dx)$$

$$= x^2 \sin x - 2 (-x \cos x + \int \cos x dx)$$

$$= x^2 \sin x - 2 (-x \cos x + \sin x) + C$$

$$= x^2 \sin x + 2x \cos x - 2 \sin x + C$$

$$\int x^n \cos x dx \rightarrow n \text{ rounds}$$

Example

$$\int e^x \cos x \, dx$$

E ←      ← T

LIATE

T before E so,  $u = \cos x$

$$u = \cos x \quad dv = e^x dx$$

$$du = -\sin x dx \quad v = e^x$$

$$\int e^x \cos x \, dx = uv - \int v du$$

$$= e^x \cos x - \int e^x \cdot -\sin x \, dx$$

$$= e^x \cos x + \int e^x \sin x \, dx$$

by parts

$$U = \sin x \quad dV = e^x dx$$

$$dU = \cos x dx \quad V = e^x$$

$$\int e^x \cos x \, dx = e^x \cos x + (uv - \int v du)$$

$$= e^x \cos x + e^x \sin x - \int e^x \cos x \, dx$$

orig. integral

by parts in infinite loop

STOP!

$$\int e^x \cos x dx = e^x \cos x + e^x \sin x - \int e^x \cos x dx$$

add  $\int e^x \cos x dx$  to both sides

$$2 \int e^x \cos x dx = e^x (\cos x + \sin x)$$

so,  $\int e^x \cos x dx = \frac{1}{2} e^x (\cos x + \sin x) + C$

this is an example of a case where going against LIATE

still works:

$$u = e^x \quad dv = \cos x dx$$

$$du = e^x dx \quad v = \sin x$$



**EXAM 1 REVIEW  
SESSION**

**SEPT 18th  
(Monday)  
5:30pm-7:30pm  
MATH 175**

