

10.3 Infinite Series

NOT on exam 2

$$\sum_{k=1}^{\infty} a_k = a_1 + a_2 + a_3 + \dots$$

a_k : k th term, usually formula is given

many kinds of series to study

today: geometric series
telescoping series

Geometric Series: common ratio between the terms

for example, $\frac{1}{4} + \frac{1}{12} + \frac{1}{36} + \frac{1}{108} + \dots$ ratio: $r = \frac{1}{3}$

we can write a geometric series in this form:

$$a + ar + ar^2 + ar^3 + ar^4 + \dots = \sum_{k=0}^{\infty} ar^k$$

$$\frac{1}{4} + \frac{1}{12} + \frac{1}{36} + \frac{1}{108} + \dots = \frac{1}{4} \left(1 + \frac{1}{3} + \frac{1}{3^2} + \frac{1}{3^3} + \dots \right)$$

$$= \frac{1}{4} + \frac{1}{4} \left(\frac{1}{3} \right) + \frac{1}{4} \left(\frac{1}{3} \right)^2 + \frac{1}{4} \left(\frac{1}{3} \right)^3 + \dots =$$

$$\boxed{\sum_{k=0}^{\infty} \frac{1}{4} \left(\frac{1}{3} \right)^k}$$

the starting k value doesn't really matter

$$\sum_{k=0}^{\infty} ar^k = a + ar + ar^2 + ar^3 + \dots$$

notice $\sum_{k=1}^{\infty} ar^{k-1}$ is the same series

$$\begin{aligned} &= ar^{1-1} + ar^{2-1} + ar^{3-1} + \dots \\ &= ar^0 + ar^1 + ar^2 + \dots \end{aligned}$$

$k=1$

$k=2$

$k=3$

$$= ar^0 + ar^1 + ar^2 + \dots = a + ar + ar^2 + ar^3 + \dots$$

Same

the partial sums of a geometric series

$$\frac{1}{4} + \frac{1}{12} + \frac{1}{36} + \frac{1}{108} + \frac{1}{324} + \frac{1}{972} + \dots$$

1st partial sum

$$S_1 = \frac{1}{4}$$

$$S_2 = \frac{1}{4} + \frac{1}{12}$$

$$S_3 = \frac{1}{4} + \frac{1}{12} + \frac{1}{36}$$

...

$$\sum_{k=0}^{\infty} ar^k = a + ar + ar^2 + ar^3 + \dots$$

$$S_1 = a = ar^0 \quad \textcircled{1}$$

$$S_2 = a + ar \quad \textcircled{1}$$

$$S_3 = a + ar + ar^2 \quad \textcircled{2}$$

$$S_4 = a + ar + ar^2 + ar^3 \quad \textcircled{3}$$

⋮

$$S_n = a + ar + ar^2 + \dots + ar^{n-1} \quad \text{---} \quad \textcircled{1}$$

let's find a formula for S_n w/o adding n terms
multiply by r

$$rS_n = ar + ar^2 + ar^3 + \dots + ar^n \quad \text{---} \quad \textcircled{2}$$

$$\textcircled{1} - \textcircled{2}$$

$$S_n - rS_n = a - ar^n$$

$$S_n(1-r) = a - ar^n \quad \rightarrow$$

$$S_n = \frac{a - ar^n}{1-r}$$

pattern: n th partial sum ends
w/ $n-1$ power of r

$$\frac{1}{4} + \frac{1}{36} + \frac{1}{108} + \dots = \sum_{k=0}^{\infty} \frac{1}{4} \left(\frac{1}{3}\right)^k \rightarrow a = \frac{1}{4}, r = \frac{1}{3}$$

7th partial sum (w/o adding 7 terms)

$$S_7 = \frac{a - ar^7}{1-r} = \frac{\frac{1}{4} - \frac{1}{4} \left(\frac{1}{3}\right)^7}{1 - \frac{1}{3}} = \dots = \boxed{\frac{1093}{2916}}$$

bigger question: does $S_n \rightarrow$ finite value as $n \rightarrow \infty$

in other words, does $\sum_{k=0}^{\infty} ar^k$ converge? $\lim_{n \rightarrow \infty} S_n = L$?

$$\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \frac{a(1-r^n)}{1-r} = \frac{a}{1-r} \lim_{n \rightarrow \infty} (1-r^n)$$

limit exists only
if $r^n \rightarrow 0$ as $n \rightarrow \infty$
exists

and that happens if

$$|r| < 1 \text{ or } -1 < r < 1$$

so, for $\sum_{k=0}^{\infty} ar^k$, it converges if $|r| < 1$

and its sum is $\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \frac{a(1-r^{n+1})}{1-r}$ goes to 0

$$= \boxed{\frac{a}{1-r}}$$

first term
common ratio

try on $\frac{1}{4} + \frac{1}{12} + \frac{1}{36} + \frac{1}{108} + \dots = \sum_{k=0}^{\infty} \frac{1}{4} \left(\frac{1}{3}\right)^k$

$$= \frac{a}{1-r} = \frac{\frac{1}{4}}{1-\frac{1}{3}} = \frac{\frac{1}{4}}{\frac{2}{3}} = \boxed{\frac{3}{8}}$$

$$1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \frac{1}{16} - \dots = \sum_{k=0}^{\infty} \left(-\frac{1}{2}\right)^k = \frac{a}{1-r} = \frac{1}{1-\frac{1}{2}} = \frac{2}{1}$$

Telescoping Series

Example

$$\sum_{k=1}^{\infty} \frac{1}{(k+1)(k+2)}$$

$$= \frac{1}{12 \cdot 13} + \frac{1}{13 \cdot 14} + \frac{1}{14 \cdot 15} + \frac{1}{15 \cdot 16} + \dots = ?$$

$k=1$ $k=2$

$$= \frac{A}{k+1} + \frac{B}{k+2}$$

not clear
where this is
going

partial fraction: $\frac{1}{(k+1)(k+2)} = \frac{1}{k+1} - \frac{1}{k+2}$

$$\sum_{k=1}^{\infty} \left(\frac{1}{k+1} - \frac{1}{k+2} \right)$$

$$= \left(\frac{1}{12} - \cancel{\frac{1}{13}} \right) + \left(\cancel{\frac{1}{13}} - \cancel{\frac{1}{14}} \right) + \left(\cancel{\frac{1}{14}} - \frac{1}{15} \right) + \dots$$

$$S_3 = \frac{1}{12} - \frac{1}{15} \quad \text{only first and last survive}$$

$$S_4 = \frac{1}{12} - \frac{1}{16}$$

$$S_5 = \frac{1}{12} - \frac{1}{17}$$

now generalize for S_n

$$S_n = \frac{1}{12} - \frac{1}{12+n}$$

does this series converge? yes, because $\lim_{n \rightarrow \infty} S_n$ exists

what does it converge to? $\frac{1}{12}$

if we keep adding terms, the sum approaches $\frac{1}{12}$

the starting k does NOT affect convergence

if a series converges, changing starting k results in a

for example, $\sum_{k=0}^{\infty} \left(\frac{1}{2}\right)^k$ converges because $|r| < 1$ convergent series


$$= \left(1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8}\right) + \left(\frac{1}{16} + \frac{1}{32} + \frac{1}{64} + \dots\right) = 2 = \frac{1}{1-\frac{1}{2}} = 2$$

finite series

$$\sum_{k=4}^{\infty} \left(\frac{1}{2}\right)^k$$

ALWAYS converges

must also converge because it's a part of a convergent series



EXAM REVIEW
MON 10/16
5:30 - 7:30 pm
University
Church 114

