

11.1 Approximating Functions with Polynomials

NOT ON exam 3

$$\text{Power Series: } \sum_{k=0}^{\infty} C_k (x-a)^k$$

$$= C_0 + C_1 (x-a) + C_2 (x-a)^2 + C_3 (x-a)^3 + C_4 (x-a)^4 + \dots$$

a : center of power series

C_k : coefficients of the k^{th} order term

the power series we will investigate is the Taylor Series

idea: write a power series that behaves like a function $f(x)$ of our choice

Taylor series matches the function value and all derivatives
at $x = a$

so near $x = a$, Taylor series acts like the real $f(x)$

Taylor series of $f(x)$

$$f(x) = C_0 + C_1(x-a) + C_2(x-a)^2 + C_3(x-a)^3 + C_4(x-a)^4 + \dots$$

match function value at $x=a$

$$f(a) = C_0 + C_1 \cancel{(a-a)} + C_2 \cancel{(a-a)^2} + C_3 \cancel{(a-a)^3} + \dots \rightarrow \boxed{f(a) = C_0} = 0! C_0$$

now match derivatives at $x=a$

$$f'(x) = C_1 + 2C_2(x-a) + 3C_3(x-a)^2 + 4C_4(x-a)^3 + \dots$$

$$f'(a) = C_1 \rightarrow \boxed{f'(a) = C_1} = 1! C_1$$

$$f''(x) = 2C_2 + 3 \cdot 2C_3(x-a) + 4 \cdot 3C_4(x-a)^2 + \dots$$

$$f''(a) = 2C_2 \rightarrow \boxed{f''(a) = 2C_2} = 2! C_2$$

$$f'''(x) = 3 \cdot 2C_3 + 4 \cdot 3 \cdot 2C_4(x-a) + \dots$$

$$f'''(a) = 3 \cdot 2C_3 \rightarrow \boxed{f'''(a) = 3 \cdot 2C_3} = 3! C_3$$

$$f^{(4)}(x) = 4 \cdot 3 \cdot 2C_4 + \dots$$

$$f^{(4)}(a) = 4 \cdot 3 \cdot 2C_4 \rightarrow \boxed{f^{(4)}(a) = 4 \cdot 3 \cdot 2C_4} = 4! C_4$$

generalize: $f^{(k)}(a) = k! C_k \rightarrow$

$$C_k = \frac{f^{(k)}(a)}{k!}$$

Taylor series of $f(x)$ at $x=a$ is

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k$$

$$= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!} (x-a)^2 + \frac{f'''(a)}{3!} (x-a)^3 + \frac{f^{(4)}(a)}{4!} (x-a)^4 + \dots$$

if we stop at $k=1 \rightarrow$ Linear approximation

$$f(x) \approx f(a) + f'(a)(x-a)$$

think of Taylor series as an extension of linear approx.
more shape features added each k

Example Find the Taylor series of $f(x) = e^x$ at $x = a$ $a = 0$

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k$$

$$= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!} (x-a)^2 + \frac{f'''(a)}{3!} (x-a)^3 + \dots$$

$$f(x) = e^x \quad a=0$$

$$f(0) = e^0 = 1$$

$$f'(x) = e^x$$

$$f'(0) = e^0 = 1$$

$$f''(x) = e^x$$

$$f''(0) = 1$$

\vdots

$$f^{(k)}(0) = 1$$

so we get

$$\begin{aligned} f(x) &= 1 + 1 \cdot (x-0) + \frac{1}{2!} (x-0)^2 + \frac{1}{3!} (x-0)^3 + \dots \\ &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots \\ &= 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots \approx e^x \text{ near } x=0 \end{aligned}$$

the more terms we include, the better the approx.

as $k \rightarrow \infty \Rightarrow$ series converges to e^x

if we cut off after k , we see the k th-order Taylor Polynomial (P_k)

$$P_0 = 1$$

$$P_1 = 1 + x \quad (\text{linear approx})$$

$$P_2 = 1 + x + \frac{x^2}{2}$$

$$P_3 = 1 + x + \frac{x^2}{2} + \frac{x^3}{6}$$


$$\left. \begin{array}{l} P_0 \\ P_1 \\ P_2 \\ P_3 \end{array} \right\} \approx e^x \text{ near } x=0$$

one use of this: approx. $e^{0.5}$

w/o calculator, $e^{0.5} = ?$

but $f(x) \approx e^x \approx 1+x$

so $e^{0.5} \approx 1+0.5 \approx 1.5$ (true value is $e^{0.5} = 1.6487$)



Example Find the 4th-order Taylor polynomial of $f(x) = \cos(2x)$

$$\text{near } x = a = \frac{\pi}{8}$$

(so we want a 4th order polynomial that behaves like $\cos(2x)$ near $x = \pi/8$)

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k \quad a = \pi/8$$

up to $k=4$

$$f(x) = f\left(\frac{\pi}{8}\right) + f'\left(\frac{\pi}{8}\right)\left(x - \frac{\pi}{8}\right) + \frac{1}{2!} f''\left(\frac{\pi}{8}\right)\left(x - \frac{\pi}{8}\right)^2 + \frac{1}{3!} f'''\left(\frac{\pi}{8}\right)\left(x - \frac{\pi}{8}\right)^3 + \frac{1}{4!} f^{(4)}\left(\frac{\pi}{8}\right)\left(x - \frac{\pi}{8}\right)^4$$

$$f(x) = \cos(2x) \rightarrow f\left(\frac{\pi}{8}\right) = \cos\left(\frac{\pi}{4}\right) = \frac{\sqrt{2}}{2}$$

$$f'(x) = -2 \sin(2x) \rightarrow f'\left(\frac{\pi}{8}\right) = -2 \sin\left(\frac{\pi}{4}\right) = -\sqrt{2}$$

$$f''(x) = -4 \cos(2x) \rightarrow f''\left(\frac{\pi}{8}\right) = -4 \cos\left(\frac{\pi}{4}\right) = -2\sqrt{2}$$

$$f'''(x) = 8 \sin(2x) \rightarrow f'''(\frac{\pi}{2}) = 8 \cdot \sqrt{2} = 4\sqrt{2}$$

$$f^{(4)}(x) = 16 \cos(2x) \rightarrow f^{(4)}(\frac{\pi}{2}) = 8\sqrt{2}$$

So, near $x = \frac{\pi}{2}$, $\cos(2x)$ behaves like

$$\frac{\sqrt{2}}{2} - \sqrt{2} \left(x - \frac{\pi}{2}\right) - \frac{2\sqrt{2}}{2!} \left(x - \frac{\pi}{2}\right)^2 + \frac{4\sqrt{2}}{3!} \left(x - \frac{\pi}{2}\right)^3 + \frac{8\sqrt{2}}{4!} \left(x - \frac{\pi}{2}\right)^4$$

$f(\frac{\pi}{2}) \quad f'(\frac{\pi}{2})$


$$= \underbrace{\frac{\sqrt{2}}{2}}_{P_0} - \underbrace{\sqrt{2} \left(x - \frac{\pi}{2}\right)}_{P_1} - \underbrace{\sqrt{2} \left(x - \frac{\pi}{2}\right)^2}_{P_2} + \underbrace{\frac{2\sqrt{2}}{3} \left(x - \frac{\pi}{2}\right)^3}_{P_3} + \underbrace{\frac{\sqrt{2}}{3} \left(x - \frac{\pi}{2}\right)^4}_{P_4}$$

P_1

P_2

P_3

P_4



EXAM REVIEW
MON 11/6
6:30 - 8:30 pm
WTHR 200

