


6.5 + 6.6 Length and Surface Area




Same idea as last time: find the length of one small piece
then accumulate using integration



is approximately the length of
the green piece

the length L of the curve (green)

$$L \approx \sqrt{(\Delta x)^2 + (\Delta y)^2}$$



$$\sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(\Delta x)^2 \left[1 + \left(\frac{\Delta y}{\Delta x} \right)^2 \right]} = \sqrt{1 + \left(\frac{\Delta y}{\Delta x} \right)^2} (\Delta x)$$

now shrink the interval: $\Delta x \rightarrow dx$

$$\frac{\Delta y}{\Delta x} \rightarrow \frac{dy}{dx}$$

so, the exact length of the small piece is $\sqrt{1 + \left(\frac{dy}{dx} \right)^2} dx$

now accumulate all from $x=a$ to $x=b$

so, the exact length of $y=f(x)$ from $x=a$ to $x=b$ is

$$\boxed{\int_a^b \sqrt{1 + \left(\frac{dy}{dx} \right)^2} dx} \quad \text{or} \quad \int_a^b \sqrt{1 + (y')^2} dx$$


$y=f(x)$ must be continuous and differentiable on $a \leq x \leq b$

$f'(x)$ must exist on $a \leq x \leq b$

example

$$y = \frac{2}{3} (x^2 - 1)^{3/2}$$

from $x=1$ to $x=\sqrt{10}$



$$L = \int_a^b \sqrt{1 + (y')^2} dx$$

$$y' = \frac{2}{3} \cdot \frac{3}{2} (x^2 - 1)^{1/2} \cdot 2x = 2x (x^2 - 1)^{1/2}$$

$$\int_1^{\sqrt{10}} \sqrt{1 + [2x(x^2 - 1)^{1/2}]^2} dx = \int_1^{\sqrt{10}} \sqrt{1 + 4x^2(x^2 - 1)} dx$$

$$= \int_1^{\sqrt{10}} \underbrace{\sqrt{4x^4 - 4x^2 + 1}}_{(2x^2 - 1)^2} dx = \dots = \boxed{\frac{17}{3}\sqrt{10} + \frac{1}{3}}$$

back to $\sqrt{(\Delta x)^2 + (\Delta y)^2}$

now factor out $(\Delta y)^2$

$$\sqrt{(\Delta y)^2 \left(\frac{(\Delta x)^2}{(\Delta y)^2} + 1 \right)}$$


$$= \sqrt{1 + \left(\frac{\Delta x}{\Delta y} \right)^2} (\Delta y)$$

shrink interval: $\Delta y \rightarrow dy$, $\frac{\Delta x}{\Delta y} \rightarrow \frac{dx}{dy}$

the equivalent formula for length of $x = f(y)$ from $y=c$ to $y=d$

$$\int_c^d \sqrt{1 + \left(\frac{dx}{dy} \right)^2} dy$$

is




Sometimes the switch in variable is for convenience

Sometimes we have to switch, for example, when $f'(x)$ does not exist

Somewhere on $a \leq x \leq b$ in $\int_a^b \sqrt{1 + [f'(x)]^2} dx$

example $y = x^{2/3}$ from $x=0$ to $x=1$



$$y' = \frac{2}{3} x^{-1/3} = \frac{2}{3 x^{1/3}}$$

$$\int_0^1 \sqrt{1 + \left(\frac{2}{3x^{1/3}}\right)^2} dx$$

DNE at $x=0$

try switching to $x = g(y)$

$$y = x^{2/3}$$

$$x = y^{3/2}$$

now use $\int_c^d \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$

$c=0, d=1$ for this example

(y of starting point is 0)

(y of ending point is 1)



$x = y^{3/2} \quad \frac{dx}{dy} = \frac{3}{2} y^{1/2}$ which always exists on $0 \leq y \leq 1$

$$\int_0^1 \sqrt{1 + \left(\frac{3}{2} y^{1/2}\right)^2} dy = \int_0^1 \sqrt{1 + \frac{9}{4} y} dy$$

$u = 1 + \frac{9}{4} y$ and so on



$$= \dots = \frac{1}{27} (13\sqrt{13} - 8) \approx 1.44$$

6.6 Surface Area of Solid of Revolution




revolve around x-axis

Volume: disk or shell methods
 Surface area = ?



rectangle → strip on surface


idea: find area of one strip
 then integrate to accumulate
 All strips



cut, then unwrap d_1 (diameter from r_1)



Shrink interval, the curved strip will be
 approximately a rectangle



$$= \int \sqrt{1 + [f'(x)]^2} dx$$

each strip is area $2\pi \cdot f(x) \cdot \sqrt{1 + [f'(x)]^2} dx$

accumulate from $x=a$ to $x=b$

$$\int_a^b 2\pi f(x) \sqrt{1 + [f'(x)]^2} dx$$


around x-axis

equivalent form
 (around y-axis)

$$\int_c^d 2\pi g(y) \sqrt{1 + [g'(y)]^2} dy$$

example Region bounded by $y = \sqrt{x}$, $x=1$, $x=4$, $y=0$

revolved around x -axis



$$\int_a^b 2\pi f(x) \sqrt{1+(f')^2} dx$$

$$f = \sqrt{x} = x^{1/2} \quad f' = \frac{1}{2}x^{-1/2} = \frac{1}{2\sqrt{x}}$$

$$\int_1^4 2\pi \sqrt{x} \sqrt{1 + \left(\frac{1}{2\sqrt{x}}\right)^2} dx$$

$$= 2\pi \int_1^4 \sqrt{x} \sqrt{1 + \frac{1}{4x}} dx = 2\pi \int_1^4 \sqrt{x \left(1 + \frac{1}{4x}\right)} dx$$

$$= 2\pi \int_1^4 \sqrt{x + \frac{1}{4}} dx \quad u = x + \frac{1}{4} \text{ and so on}$$

$$= \dots = \frac{\pi}{6} (17\sqrt{17} - 5\sqrt{5})$$