

9.3 Fourier Cosine and Sine Series (part 2)

$f(x)$ periodic period $2l$

even: Fourier cosine series $b_n = 0$ $a_n = \frac{2}{l} \int_0^l f(x) \cos\left(\frac{n\pi x}{l}\right) dx$

odd: Fourier sine series $a_n = 0$ $b_n = \frac{2}{l} \int_0^l f(x) \sin\left(\frac{n\pi x}{l}\right) dx$

Boundary-value problems: conditions specified at boundaries

instead of initial points (initial-value problems)

Example: $x'' + 4x = 0$

$x(0) = x(\pi) = 0$ w/o derivatives
"Dirichlet condition"
boundaries

NOT both at 0

(e.g. $x(0) = 0, x'(0) = 0$)

or $x'(0) = x'(\pi) = 0$ both derivatives

"Neumann condition"

"old way" solve $x'' + 4x = 0 \rightarrow r = \pm 2i$

$x(t) = C_1 \cos 2t + C_2 \sin 2t$, then use BC's to find C's

nonhomogeneous is more complicated

$$x'' + 2x = 1$$

"old way" solve homogeneous $x'' + 2x = 0$

then solve for the particular solution

due to nonhomogeneous term (undetermined coeffs

or variation of params)

Alternative way: use Fourier series (sines and cosines are

very good at meeting

BC's because they

are periodic)

expand the nonhomogeneous term as Fourier series

expand x as Fourier series w/ unknown coeffs

then solve for the coeffs

Example

$$X'' + 2X = 1 \quad X(0) = X(\pi) = 0$$

first, expand the nonhomogeneous term $f(t) = 1$


as a Fourier series

$$f(t) = 1 \quad 0 < t < \pi$$


add either odd or even extensions whichever

meet the BC's $X(0) = X(\pi) = 0$

$$f(t) = 1 \quad 0 < t < \pi \quad \text{period } 2\pi$$




with even extensions



note w/ even extensions
the expansion will

NOT meet the BC's

with odd extensions



meet the BC's?

yes. Because the series converges to the average value at discontinuities

so, we need to expand $f(t) = 1$ for $0 < t < \pi$ period 2π

as a Fourier Sine series

$$a_n = 0 \quad b_n = \frac{2}{L} \int_0^L f(t) \sin\left(\frac{n\pi t}{L}\right) dt$$

$$= \dots = \frac{2}{n\pi} [1 - (-1)^n]$$

$$f(t) = \sum_{n=0}^{\infty} \frac{2}{n\pi} [1 - (-1)^n] \sin(nt)$$

now left side of $X'' + 2X = 1$

Expand X as a general (unknown) Fourier series w/ some period as the right side

$$x(t) = \frac{1}{2} A_0 + \sum_{n=1}^{\infty} A_n \cos\left(\frac{n\pi t}{L}\right) + B_n \sin\left(\frac{n\pi t}{L}\right)$$

$$x = \frac{1}{2} A_0 + \sum_{n=1}^{\infty} A_n \cos(nt) + B_n \sin(nt)$$

Sub into left side

$$x' = \sum_{n=1}^{\infty} -n A_n \sin(nt) + n B_n \cos(nt)$$

of $x'' + 2x = 1$

$$x'' = \sum_{n=1}^{\infty} -n^2 A_n \cos(nt) - n^2 B_n \sin(nt)$$

$$\sum_{n=1}^{\infty} \frac{2}{n\pi} [1 - (-1)^n] \sin(nt)$$

goal: find A_n and B_n

$$\sum_{n=1}^{\infty} \underbrace{-n^2 A_n \cos(nt) - n^2 B_n \sin(nt)}_{x''} + A_0 + \sum_{n=1}^{\infty} \underbrace{2A_n \cos(nt) + 2B_n \sin(nt)}_{2x}$$

$$= \sum_{n=1}^{\infty} \frac{2}{n\pi} [1 - (-1)^n] \sin(nt)$$

collect like terms

$$A_0 + \sum_{n \neq 1}^{\infty} (2-n^2)A_n \cos(nt) + \sum_{n \neq 1}^{\infty} (2-n^2)B_n \sin(nt) = \sum_{n \neq 1}^{\infty} \frac{2}{n\pi} [1 - (-1)^n] \sin(nt)$$

NO cos terms

NO sines

$$\text{so, } A_0 = 0 \quad A_n = 0$$

$$\text{and } B_n = \frac{2}{n\pi(2-n^2)} [1 - (-1)^n]$$

$$\text{so, the solution to } x'' + 2x = 1 \quad x(0) = x(\pi) = 0$$

$$\text{is } x(t) = \sum_{n \neq 1}^{\infty} \frac{2}{n\pi(2-n^2)} [1 - (-1)^n] \sin(nt)$$

first few terms:

$$x(t) = \frac{4}{\pi} \sin(t) - \frac{4}{21\pi} \sin(3t) - \frac{4}{105\pi} \sin(5t) - \dots$$