

Lecture 21

• Last time :

- root/coroot and weight/coweight lattices
- Weyl chambers, their walls
- positive Weyl chamber C_+ for any polarization $R = R_+ \cup R_-$

• Finish pages 4-5 of Lecture 20 notes.

↳ this provides the positive answer to Q1 from [Lect 20, p.3] which asked if different polarizations give rise to equivalent sets of simple roots

• We shall now also provide an affirmative answer to Q2 from [Lect 20, p.3] which asked if the entire root system R can be recovered from Π , the set of simple roots. We set $s_i := s_{\alpha_i} \forall \alpha_i \in \Pi$, called simple reflections.

To this end, we start from the following simple result:

Lemma 1: Let R be a root system with a fixed polarization $R = R_+ \cup R_-$, and let $\Pi = \{\alpha_1, \dots, \alpha_r\} \subset R_+$ be the set of simple roots. Then for every Weyl chamber C , there exist $i_1, \dots, i_r \in \{1, \dots, r\}$ s.t. $C = s_{i_1} s_{i_2} \dots s_{i_r}(C_+)$, where C_+ = positive Weyl chamber

↳ Pick two points $t \in C$ and $t_+ \in C_+$ generically enough, connect them by a line segment, and consider its intersections with root hyperplanes (the "generic" condition is to guarantee that all these intersections are "simple") Let N be the number of such intersections, and we will argue by induction on N

Base ($N=1$): In this case, C is adjacent to C_+ , but all walls of C_+ are $L_{\alpha_1}, \dots, L_{\alpha_r}$, hence, $\exists i_1: C = s_{i_1}(C_+)$ by [Lect 20, Lemma 3a]

Induction Step ($N-1 \rightsquigarrow N$): Let C' be the chamber first entered as we move from t along the above segment, so that $C = s_{\alpha_j}(C')$ where L_{α_j} is a wall of C' . By induction hypothesis, C' can be written as

$C' = u(C_+)$ for some $u = s_{i_2} s_{i_3} \dots s_{i_k}(C_+)$ with $i_2, \dots, i_k \in \{1, \dots, r\}$.

In particular, the wall $L_{\alpha_j} = u(L_{\alpha_j})$ for some $1 \leq j \leq r \Rightarrow s_{\alpha_j} = u s_{\alpha_j} u^{-1}$

Lecture #21

(continuation)

Thus: $C = s_\alpha(C') = s_\alpha(u(C_+)) = u s_j u^{-1}(C_+) = u s_j(C_+)$

This implies the result, with $l = k+1$ and $s_l = j$

With this result, we can immediately prove:

Theorem 1: In the notations of Lemma 1:
a) The simple reflections $\{s_i\}_{i=1}^r$ generate $W = \text{Weyl gp}$
b) $W(\Pi) = R$

a) $\forall \alpha \in R$, the hyperplane L_α is a wall of some Weyl chamber C .

By Lemma 1, $C = s_{i_1} \dots s_{i_\ell}(C_+)$ for some $i_1, \dots, i_\ell \Rightarrow L_\alpha = s_{i_1} \dots s_{i_\ell}(L_{\alpha_j})$

for some j . Hence: $\alpha = \pm w(\alpha_j)$ with $w = s_{i_1} \dots s_{i_\ell}$

note: $s_i(p) = -p$
AND
 $s_\alpha = w s_j w^{-1}$
b) \Rightarrow a)

Upshot: The whole root system R can be reconstructed from Π as $W(\Pi)$, where $W = \langle s_1, \dots, s_r \rangle$ - subgroup of $O(E)$ generated by simple reflections.

Example (A_{n-1} -type root system): $W = S_n$, $s_i = (i, i+1)$. Thus part a) of Thm 1 says that any permutation can be written as a product of transpositions.

Length function

We shall say that a root hyperplane L_α separates two Weyl chambers C, C' if they lie on different sides of L_α , i.e. $\alpha(C) \neq \alpha(C')$ have different signs.

[Warning: We don't assume L_α to be a wall of either C or C']

Def 1: The length $l(w) \in \mathbb{Z}_{\geq 0}$ of $w \in W$ is the number of root hyperplanes separating C_+ and $w(C_+)$

As $\alpha(C_+) > 0 \forall \alpha \in R_+$ and $\alpha(w(C_+)) = (w^{-1}\alpha)(C_+)$, we can equivalently write

$l(w) = \#\{\alpha \in R_+ \mid w^{-1}(\alpha) \in R_-\}$

Furthermore, if α is as above, then $\beta = -w^{-1}(\alpha)$ satisfies $\beta(C_+) > 0$ & $w(\beta)(C_+) < 0$ and vice versa, so that $l(w) = l(w^{-1}) \Rightarrow l(w) = \#\{\alpha \in R_+ \mid w(\alpha) \in R_-\}$

Lecture #21

Example: $w = s_i$ - simple reflection. In this case, C_+ and $s_i(C_+)$ are clearly separated only by $L_{\alpha_i} \Rightarrow l(s_i) = 1$. Therefore:

$\{\alpha \in R_+ \mid s_i(\alpha) \in R_+\} = \{\alpha_i\} \Rightarrow s_i$ permutes the set $R_+ \setminus \{\alpha_i\}$

Lemma 2: Let $\rho = \frac{1}{2} \sum_{\alpha \in R_+} \alpha$. Then $\alpha_i(\rho) = 1 \forall i$. Hence, $\rho = \sum_{i=1}^r w_i$

↑ this weight is very open in the repr. theory (as we will soon see)

Example $\rho = \frac{1}{2} \alpha_i + \frac{1}{2} \sum_{\alpha \in R_+ \setminus \{\alpha_i\}} \alpha \Rightarrow s_i(\rho) = -\frac{1}{2} \alpha_i + \frac{1}{2} \sum_{\alpha \in R_+ \setminus \{\alpha_i\}} \alpha = \rho - \alpha_i \Rightarrow \alpha_i(\rho) = 1 \Rightarrow \rho = \sum_{i=1}^r w_i$

We shall now derive an equivalent definition of the length $l(w)$:

Theorem 2: Let $w = s_{i_1} s_{i_2} \dots s_{i_\ell}$ be a representation of $w \in W$ as a product of simple reflections that has minimal possible length. Then $\ell = l(w)$.

As in the proof of Thm 1, consider a chain of Weyl chambers $C_k = s_{i_1} \dots s_{i_k}(C_+)$ so that $C_0 = C_+$, $C_\ell = w(C_+)$, and $0 \leq k \leq \ell$. As we already saw, C_k & C_{k-1} are adjacent Weyl chambers (separated by L_{β_k} with $\beta_k = s_{i_1} \dots s_{i_{k-1}}(\alpha_{i_k})$). This implies the inequality $l(w) \leq \ell$, as every separating hyperplane should be precisely one of L_{β_k} above. ↑ CHECK

On the other hand, picking generic points $t_+ \in C_+$ and $t_+^w \in w(C_+)$, the line segment connecting them will intersect every separating root hyperplane and no other root hyperplanes. Then, the prop of Thm 1 shows that w can be written as a product of $l(w)$ simple reflections $\Rightarrow \ell \leq l(w) \Rightarrow \ell = l(w)$.

Def 2: An expression $w = s_{i_1} s_{i_2} \dots s_{i_\ell}$ is called reduced if $\ell = l(w)$

Corollary 1: The action $W \curvearrowright \{\text{Weyl chambers}\}$ is simply transitive.

We know this action is transitive. Hence, it suffices to show $w(C_+) = C_+ \Rightarrow w = 1$. But by definition, $l(w) = 0 \xrightarrow{\text{Thm 2}} w = 1$

Lecture #21

Let \bar{C}_+ be the closure of the positive Weyl chamber. Then, by above every W -orbit of $W \backslash \mathfrak{E}$ has an element from \bar{C}_+ . In fact a stronger result holds:

Proposition 1: $E/W \cong \bar{C}_+$, i.e. every W -orbit of $W \backslash \mathfrak{E}$ contains exactly 1 elt in \bar{C}_+

Assume the contrary, i.e. $\exists \lambda, \mu \in \bar{C}_+ \exists w \in W$ s.t. $\mu = w(\lambda)$. We assume that w is the shortest possible, and $w \neq 1$. Consider a reduced decomposition $w = s_{i_1} \dots s_{i_\ell}$. Note that $s_{i_1} w = s_{i_2} \dots s_{i_\ell} \Rightarrow \ell(s_{i_1} w) \leq \ell - 1 < \ell(w)$.

Thus, $\exists \gamma \in \mathfrak{R}_+$ s.t. $w(\gamma) \in \mathfrak{R}_-$ but $s_{i_1} w(\gamma) \in \mathfrak{R}_+ \Rightarrow w(\gamma) = -\alpha_{i_1} \Rightarrow$

$\Rightarrow \gamma = -s_{i_\ell} s_{i_{\ell-1}} \dots s_{i_2} \underbrace{s_{i_1}(-\alpha_{i_1})}_{-\alpha_{i_1}} = s_{i_\ell} \dots s_{i_2}(\alpha_{i_1}) \Rightarrow S_\gamma = s_{i_\ell} \dots s_{i_2} s_{i_1} s_{i_2} \dots s_{i_\ell}$

Then: $0 \leq (\lambda, \gamma) = \underbrace{(w(\lambda), w(\gamma))}_{\mu} = (\mu, \underbrace{w(\gamma)}_{\in \mathfrak{R}_-}) \leq 0 \Rightarrow (\lambda, \gamma) = 0 \Rightarrow S_\gamma(\lambda) = \lambda$.

But then, we get:

$\mu = w(\lambda) = s_{i_1} \dots s_{i_\ell}(\lambda) = \underbrace{s_{i_2} \dots s_{i_\ell} s_{i_\ell} \dots s_{i_2} s_{i_1} s_{i_2} \dots s_{i_\ell}}_{=id}(\lambda) = s_{i_2} \dots s_{i_\ell} S_\gamma(\lambda) = s_{i_2} \dots s_{i_\ell}(\lambda)$

Hence, contradiction with w being of minimal length!

Given a polarization, $C_- := -C_+$ is called the negative Weyl chamber.

Corollary 2: $\exists! w_0 \in W$ s.t. $C_- = w_0(C_+)$.
Moreover, $\ell(w_0) = \#\mathfrak{R}_+$ and $\ell(w) < \ell(w_0) \forall w \in W \setminus \{w_0\}$.
Finally, $w_0^2 = 1$.

- uniqueness & existence of w_0 follow from Cor 1.
- $\ell(w_0) = \#\{\alpha \in \mathfrak{R}_+ \mid w_0(\alpha) \in \mathfrak{R}_-\} = \#\mathfrak{R}_+$
- $\ell(w) \leq \ell(w_0)$ and equality means $w(\mathfrak{R}_+) = \mathfrak{R}_- \Rightarrow w(C_+) = C_- \Rightarrow w = w_0$
- $w_0^2(C_+) = C_+ \Rightarrow w_0^2 = 1$ by Cor 1

Example (A_{n-1} -root system): w_0 is the permutation $\begin{pmatrix} 1 & 2 & 3 & \dots & n-1 & n \\ n & n-1 & n-2 & & 2 & 1 \end{pmatrix}$.