

Combinatorial analysis

Samy Tindel

Purdue University

Probability - MA 416

Mostly taken from *A first course in probability*
by S. Ross

Outline

- 1 Introduction
- 2 The basic principle of counting
- 3 Permutations
- 4 Combinations
- 5 Multinomial coefficients

Outline

- 1 Introduction
- 2 The basic principle of counting
- 3 Permutations
- 4 Combinations
- 5 Multinomial coefficients

A simple example of counting

A communication system:

- Setup: n antennas lined up
- Functional system:
↔ when no 2 consecutive defective antennas
- We know that m antennas are defective

Problem: compute

P (functional system)

A simple example of counting (2)

Particular instance of the previous situation:

- Take $n = 4$ and $m = 2$
- Possible configurations:

0011	0101	0110
1001	1010	1100

- We get 3 working configurations among 6, and thus

$$\mathbf{P}(\text{functional system}) = \frac{1}{2}$$

Conclusion: need an effective way to count, that is

Combinatorial analysis

Outline

- 1 Introduction
- 2 The basic principle of counting**
- 3 Permutations
- 4 Combinations
- 5 Multinomial coefficients

Basic principle of counting

Theorem 1.

Suppose 2 experiments to be performed and

- For Experiment 1, we have m possible outcomes
- For each outcome of Experiment 1
 \hookrightarrow We have n outcomes for Experiment 2

Then

Total number of possible outcomes is $m \times n$

Proof

Sketch of the proof: Set

$(i, j) \equiv$ Outcome i for Experiment 1 & Outcome j for Experiment 2

Then enumerate possibilities

Application of basic principle of counting

Example: Small community with

- 10 women
- Each woman has 3 children

We have to pick one pair as mother & child of the year

Question:

How many possibilities?

Generalized principle of counting

Theorem 2.

Suppose r experiments to be performed and

- For Experiment 1, we have n_1 possible outcomes
- For each outcome of Experiment i
 \hookrightarrow We have n_{i+1} outcomes for Experiment $i + 1$

Then total number of possible outcomes is

$$\prod_{i=1}^r n_i = n_1 \times n_2 \times \cdots \times n_r$$

Application of basic principle of counting

Example 1: Find # possible 7 place license plates if

- First 3 places are letters
- Final 4 places are numbers

Answer: 175,760,000

Example 2: Find # possible 7 place license plates if

- First 3 places are letters
- Final 4 places are numbers
- No repetition among letters or numbers

Answer: 78,624,000

Outline

- 1 Introduction
- 2 The basic principle of counting
- 3 Permutations**
- 4 Combinations
- 5 Multinomial coefficients

Permutations

Definition:

A permutation of n objects is an ordered sequence of those n objects.

Property:

Two permutations only differ according to the order of the objects

Counting:

Let P_n be the number of permutations for n objects. Then

$$P_n = n! = n \times (n - 1) \cdots \times 2 = \prod_{j=1}^n j$$

Example of permutation

Example: 3 balls, Red, Black, Green

Permutations: RBG, RGB, BRG, BGR, GBN, GBR

↔ 6 possibilities

Formula: $P_3 = 3! = 6$

Proof for the counting number P_n

Sketch of the proof:

Direct application of Theorem 2

Example of permutation (1)

Problem:

Count possible arrangements of letters in PEPPER

Example of permutation (2)

Solution 1:

Consider all letters as distinct objects

$$P_1 E_1 P_2 P_3 E_2 R$$

Then

$$P_6 = 6! = 720 \text{ possibilities}$$

Example of permutation (3)

Solution 2:

Do not distinguish P's and E's.

Then

$$\frac{P_6}{P_3 P_2} = \frac{6!}{3! 2!} = 60 \text{ possibilities}$$

Outline

- 1 Introduction
- 2 The basic principle of counting
- 3 Permutations
- 4 Combinations**
- 5 Multinomial coefficients

Combinations

Definition:

A combination of p objects among n objects is non ordered subset of p objects.

Property:

Two combinations only differ according to nature of their objects

Counting:

The number of combinations of p objects among n objects is

$$\binom{n}{p} = \frac{n!}{p! (n - p)!}$$

Proof of counting

Combination when order is relevant:

Number of possibilities is

$$n \times (n - 1) \cdots \times (n - p + 1) = \frac{n!}{(n - p)!}$$

Combination when order is irrelevant:

We divide by # permutations of p objects

Number of possibilities is

$$\frac{n \times (n - 1) \cdots \times (n - p + 1)}{p!} = \frac{n!}{p!(n - p)!} = \binom{n}{p}$$

Example of combination (1)

Situation:

- We have a group of 5 women and 7 men
- We wish to form a committee with 2 women and 3 men

Problem:

- Find the number of possibilities

Example of combination (2)

Number of possibilities:

$$\binom{5}{2} \binom{7}{3} = 350$$

Example of combination (3)

Situation 2:

- We have a group of 5 women and 7 men
- We wish to form a committee with 2 women and 3 men
- 2 men refuse to serve together

Problem:

- Find the number of possibilities

Example of combination (4)

New number of possibilities:

$$\binom{5}{2} \left\{ \binom{7}{3} - \binom{2}{2} \binom{5}{1} \right\} = 300$$

Binomial theorem

Theorem 3.

Let

- $x_1, x_2 \in \mathbb{R}$
- $n \geq 1$

Then

$$(x_1 + x_2)^n = \sum_{k=0}^n \binom{n}{k} x_1^k x_2^{n-k}$$

Combinatorial proof

First expansion:

$$(x_1 + x_2)^n = \sum_{(i_1, \dots, i_n) \in \{1, 2\}^n} x_{i_1} x_{i_2} \cdots x_{i_n}$$

Definition of a family of sets:

$$A_k = \{(i_1, \dots, i_n) \in \{1, 2\}^n; \text{there are } k \text{ } j\text{'s such that } i_j = 1\}.$$

New expansion: we have (convention: $|A_k| \equiv \text{Card}(A_k)$)

$$\begin{aligned}(x_1 + x_2)^n &= \sum_{k=0}^n |A_k| x_1^k x_2^{n-k} \\ &= \sum_{k=0}^n \binom{n}{k} x_1^k x_2^{n-k}\end{aligned}$$

Application of the binomial theorem

Proposition 4.

Let

- A a set with $|A| = n$
- $\mathcal{P}_n \equiv$ collection of all subsets of A

Then

$$|\mathcal{P}_n| = 2^n$$

Proof

Decomposition of $|\mathcal{P}_n|$: Write

$$\begin{aligned} |\mathcal{P}_n| &= \sum_{k=0}^n |\text{Subsets of } A \text{ with } k \text{ elements}| \\ &= \sum_{k=0}^n \binom{n}{k} \end{aligned}$$

Application of the binomial theorem:

$$\begin{aligned} |\mathcal{P}_n| &= (1 + 1)^n \\ &= 2^n \end{aligned}$$

Outline

- 1 Introduction
- 2 The basic principle of counting
- 3 Permutations
- 4 Combinations
- 5 Multinomial coefficients**

Multinomial coefficients

Divisions of n objects into r groups with size n_1, \dots, n_r : We have

- n objects and r groups
- We want n_j objects in group j and $\sum_{j=1}^r n_j = n$

Notation: Set

$$\binom{n}{n_1, \dots, n_r} = \frac{n!}{\prod_{j=1}^r (n_j!)}$$

Counting: We have

Divisions of n objects into r groups with size n_1, \dots, n_r

$$\binom{\overline{n}}{n_1, \dots, n_r}$$

Proof of counting

Number of choices for the i th group:

$$\binom{n - \sum_{j=1}^{i-1} n_j}{n_i}$$

Number of divisions: We have

Divisions of n objects into r groups with size n_1, \dots, n_r

$$\begin{aligned} &= \prod_{i=1}^r \binom{n - \sum_{j=1}^{i-1} n_j}{n_i} \\ &= \binom{n}{n_1, \dots, n_r} \end{aligned}$$

Example of multinomial coefficient (1)

Situation: Police department with 10 officers and

- 5 have to patrol the streets
- 2 are permanently working at the station
- 3 are on reserve at the station

Problem:

How many divisions do we get?

Example of multinomial coefficient (2)

Answer:

$$\frac{10!}{5! 2! 3!} = 2520$$

Tournament example

Situation: Tournament with $n = 2^m$ players

↔ How many outcomes?

Particular case:

Take $m = 3$, thus $n = 8$

Number of rounds: 3

Tournament example (2)

Counting number of outcomes for the first round:

$$\overbrace{\binom{8}{2, 2, 2, 2}}^{\text{\# pairings with order}} \overbrace{\frac{1}{4!}}^{\text{No ordering}} \overbrace{2^4}^{\text{Possible outcomes}} = \frac{8!}{4!}$$

Counting number of outcomes for second and third round:

$$\frac{4!}{2!} \quad \text{and} \quad \frac{2!}{1!}$$

Conclusion:

$$\frac{8!}{4!} \frac{4!}{2!} \frac{2!}{1!} = 8! = 40,320 \text{ possible outcomes}$$

Multinomial theorem

Theorem 5.

Let

- $x_1, x_2, \dots, x_r \in \mathbb{R}$
- $n \geq 1$

Then

$$(x_1 + x_2 + \dots + x_r)^n = \sum_{(n_1, \dots, n_r) \in A_{n,r}} \binom{n}{n_1, \dots, n_r} x_1^{n_1} x_2^{n_2} \dots x_r^{n_r}$$

where

$$A_{n,r} = \{(n_1, \dots, n_r) \in \mathbb{N}^r; n_1 + n_2 + \dots + n_r = n\}$$