

**MA59800ANT ANALYTIC THEORY OF FUNCTION FIELDS.
PROBLEMS 4**

TO BE HANDED IN BY 6PM MONDAY 28TH OCTOBER 2024

Key: **A-questions** are short questions testing basic skill sets; **B-questions** integrate essential methods of the course; **C-questions** are more challenging for enthusiasts, with hints available on request.

Throughout, we take p to be a prime number, put $q = p^h$, and denote $\mathbb{F}_q[t]$ by \mathbb{A} .

A1. Consider the polynomials $u \in \mathbb{A}$ of the shape

$$u(t) = t^n + u_{n-1}t^{n-1} + \dots + u_k t^k + \dots + u_1 t + u_0,$$

with $u_i \in \mathbb{F}_q$.

(a) Show that as $n \rightarrow \infty$, there are infinitely many irreducible polynomials of this shape with u_0, u_1, \dots, u_{k-1} all fixed to be specified values, provided that $u_0 \neq 0$. Compute the Dirichlet density of this set of polynomials.

(b) Show that, as $n \rightarrow \infty$, there are infinitely many irreducible polynomials of this shape with u_{n-1}, \dots, u_{n-k} all fixed to be specified values. Compute the Dirichlet density of this set of polynomials.

A2. (a) Show that, whenever $c \in \mathbb{F}_q$ and $d \in \mathbb{F}_q \setminus \{0\}$, there are infinitely many monic irreducible polynomials $\pi(t) \in \mathbb{A}$ having the property that $\pi(c) = d$.

(b) Show that, whenever c_1 and c_2 are distinct elements of \mathbb{F}_q , and $d_1, d_2 \in \mathbb{F}_q \setminus \{0\}$, then there are infinitely many monic irreducible polynomials $\pi(t) \in \mathbb{A}$ having the property that $\pi(c_1) = d_1$ and $\pi(c_2) = d_2$.

B3. Assume the Riemann Hypothesis for Dirichlet L -functions over $\mathbb{F}_q[t]$. Let $\Lambda(u)$ denote the von Mangoldt function in $\mathbb{A} = \mathbb{F}_q[t]$. Suppose that $a, m \in \mathbb{F}_q[t]$ satisfy $(a, m) = 1$ and $\deg(m) \geq 1$. Obtain asymptotic formulae for the following quantities as $n \rightarrow \infty$:

$$\sum_{\substack{u \in \mathbb{F}_q[t]^+ \\ \deg(u) = n \\ u \equiv a \pmod{m}}} \Lambda(u) \quad \text{and} \quad \sum_{\substack{u \in \mathbb{F}_q[t]^+ \\ \deg(u) = n \\ u \equiv a \pmod{m}}} \frac{\Lambda(u)}{|u|}.$$

B4. Recall that the forward difference operator Δ_1 is defined for any \mathbb{K}_∞ -valued function ψ of the variable x via the relation

$$\Delta_1(\psi(x); h) = \psi(x + h) - \psi(x).$$

The j -fold forward difference operator is then defined for $j \geq 2$ inductively via the relation

$$\Delta_j(\psi(x); h_1, \dots, h_j) = \Delta_1(\Delta_{j-1}(\psi(x); h_1, \dots, h_{j-1}); h_j).$$

Show that for $1 \leq j \leq k$, one has

$$\Delta_j(x^k; h_1, \dots, h_j) = \sum_{i_0, \dots, i_j} \frac{k!}{i_0! \dots i_j!} x^{i_0} h_1^{i_1} \dots h_j^{i_j},$$

where the summation is over $i_0 \geq 0, i_1 \geq 1, \dots, i_j \geq 1$ and $i_0 + \dots + i_j = k$. Hence deduce that

$$\Delta_j(x^k; h_1, \dots, h_j) = h_1 \dots h_j p_j(x; h_1, \dots, h_j),$$

where p_j is a polynomial in x of degree $k - j$ with leading coefficient $k!/(k - j)!$.

B5. Let H and X be positive integers. Suppose that $k \geq 2$ and $\alpha \in \mathbb{K}_\infty$, and define

$$T(\alpha) = \sum_{0 \leq \deg(h) < H} \left| \sum_{\deg(x) < X} e(h\alpha x^k) \right|.$$

(a) Show that

$$T(\alpha)^{2^{k-1}} \leq (q^H)^{2^{k-1}-1} \sum_{0 \leq \deg(h) < H} \left| \sum_{\deg(x) < X} e(h\alpha x^k) \right|^{2^{k-1}}.$$

(b) Suppose that $a, g \in \mathbb{F}_q[t]$ satisfy $(a, g) = 1$, g monic, and $|\alpha - a/g| < 1/|g|^2$. Show that, whenever $\text{char}(\mathbb{F}_q) > k$, one has

$$T(\alpha) \ll (q^{H+X})^{1+\varepsilon} (|g|^{-1} + q^{-X} + |g|(q^{H+kX})^{-1})^{2^{1-k}}.$$

C6. Suppose that $\text{char}(\mathbb{F}_q) \neq 2$, and put $\alpha = \sqrt{1+t^2} \in \mathbb{K}_\infty$.

(a) Show that the continued fraction expansion of α is $\alpha = [t; 2t, 2t, \dots]$.

(b) Suppose that the convergents to the continued fraction expansion of α are a_n/g_n ($n \in \mathbb{N}$). By examining $a_n^2 - (1+t^2)g_n^2$, find an infinite sequence of polynomials $(x_n, y_n)_{n=1}^\infty$ having the property that $x_n^2 - (t^2 + 1)y_n^2 = 1$.

(c) Find an explicit constant $c > 0$ having the property that, whenever a and g are elements of $\mathbb{F}_q[t]$ with g monic and $(a, g) = 1$, then $|\sqrt{1+t^2} - a/g| > c/|g|^2$.

C7. Suppose that $\theta \in \mathbb{K}_\infty$ is irrational.

(a) Show that the sequence of squares of fractional parts of θx , namely

$$(\{\theta x\}^2)_{x \in \mathbb{F}_q[t]},$$

is *not* equidistributed in \mathbb{T} .

(b) Suppose that $\text{char}(\mathbb{F}_q)$ is odd. Show that there exists an irrational element $\theta \in \mathbb{K}_\infty$ having the property that

$$(x \lfloor x\sqrt{1+t^2} \rfloor \theta)_{x \in \mathbb{F}_q[t]},$$

is *not* equidistributed in \mathbb{T} .

©Trevor D. Wooley, Purdue University 2024. This material is copyright of Trevor D. Wooley at Purdue University unless explicitly stated otherwise. It is provided exclusively for educational purposes at Purdue University, and is to be downloaded or copied for your private study only.