

ESTIMATES FOR SMOOTH WEYL SUMS ON MAJOR ARCS

JÖRG BRÜDERN AND TREVOR D. WOOLEY

ABSTRACT. We present estimates for smooth Weyl sums of use on sets of major arcs in applications of the Hardy-Littlewood method. In particular, we derive mean value estimates on major arcs for smooth Weyl sums of degree k delivering essentially optimal bounds for moments of order u whenever $u > 2\lfloor k/2 \rfloor + 4$.

1. INTRODUCTION

The introduction by Vaughan [10, 11] of smooth numbers into the armoury of the circle method practitioner created a flexible new tool for the analysis of diagonal problems involving k -th powers, and indeed the sharpest bounds currently available in Waring's problem require the use of smooth Weyl sums (see [2, 14, 15, 16, 17, 18], for example). While powerful mean value estimates for these modified Weyl sums are available for handling the minor arc contributions in applications of the circle method, the absence of correspondingly powerful major arc estimates is a source of technical complication with the potential to block certain applications. Our goal in this note is to derive estimates for smooth Weyl sums of major arc type sufficiently powerful that the vast majority of technical complications may be avoided. Indeed, our recent work on sums of squares and higher powers has already made use of these new major arc estimates (see the discussion surrounding [3, equation (3.15)]).

In order to proceed further we must introduce some notation. We employ the letter p to denote a prime number, and as usual, we abbreviate $e^{2\pi iz}$ to $e(z)$. When P and R are real numbers with $2 \leq R \leq P$, we write

$$\mathcal{A}(P, R) = \{n \in [1, P] \cap \mathbb{Z} : p \mid n \Rightarrow p \leq R\}.$$

Also, when $\nu > 1$, we put $\mathcal{A}_\nu(P, R) = \mathcal{A}(P, R) \cap (P/\nu, P]$. In addition, fixing an integer $k \geq 2$, we define the smooth Weyl sums

$$g_\nu(\alpha; P, R) = \sum_{x \in \mathcal{A}_\nu(P, R)} e(\alpha x^k) \quad \text{and} \quad g(\alpha; P, R) = \sum_{x \in \mathcal{A}(P, R)} e(\alpha x^k).$$

Finally, we define the multiplicative function $\kappa(q) = \kappa_k(q)$ by putting

$$\kappa(p^{uk+v}) = \begin{cases} kp^{-u-1/2}, & \text{when } u \geq 0 \text{ and } v = 1, \\ p^{-u-1}, & \text{when } u \geq 0 \text{ and } 2 \leq v \leq k. \end{cases}$$

2020 *Mathematics Subject Classification.* 11L07, 11L15, 11P55, 11P05.

Key words and phrases. Smooth Weyl sums, exponential sums, Waring's problem, Hardy-Littlewood method.

First author supported by Deutsche Forschungsgemeinschaft Project Number 462335009. Second author supported by NSF grants DMS-1854398 and DMS-2001549. The second author is grateful to Institut Mittag-Leffler for excellent working conditions which assisted the completion of this memoir.

We note that, as in [9, Lemma 3], when $k \geq 3$ one has $q^{-1/2} \leq \kappa(q) \ll q^{-1/k}$. When $k = 2$ we have the slightly weaker bounds $q^{-1/2} \leq \kappa(q) \leq 2^{\omega(q)} q^{-1/2}$. Here $\omega(q)$ is the number of distinct primes dividing q , and the implicit constant in Vinogradov's notation depends on k .

In §2 we establish the estimates for the exponential sums $g_\nu(\alpha) = g_\nu(\alpha; P, R)$ and $g(\alpha) = g(\alpha; P, R)$ contained in the following theorem. Here and throughout, we find it useful to introduce the notation

$$L = \log P, \quad L_2 = \log \log(3R) \quad \text{and} \quad \mathcal{L} = \log(2 + P^k |\alpha - a/q|). \quad (1.1)$$

Theorem 1.1. *Let k be a natural number with $k \geq 2$, and let ν and ε be real numbers with $\nu > 1$ and $\varepsilon > 0$. Suppose that R and P are real numbers with $2 \leq R \leq P$. Then, whenever $\alpha \in \mathbb{R}$, $a \in \mathbb{Z}$ and $q \in \mathbb{N}$ satisfy $(a, q) = 1$, one has*

$$g_\nu(\alpha) \ll \frac{q^\varepsilon \kappa(q)^{1/2} P L (L_2 \mathcal{L})^{1/2}}{(1 + P^k |\alpha - a/q|)^{1/2}} + q^\varepsilon P^{3/4} R^{1/2} (LL_2)^{1/4} (q + P^k |q\alpha - a|)^{1/8}.$$

Moreover, under the same conditions, one has

$$g(\alpha) \ll \frac{q^\varepsilon \kappa(q)^{1/2} P L L_2^{1/2}}{(1 + P^k |\alpha - a/q|)^{1/k}} + q^\varepsilon P^{3/4} R^{1/2} (LL_2)^{1/4} (q + P^k |q\alpha - a|)^{1/8} \quad (k \geq 3)$$

and

$$g(\alpha) \ll \frac{q^\varepsilon \kappa(q)^{1/2} P L L_2^{1/2} \mathcal{L}^{3/2}}{(1 + P^2 |\alpha - a/q|)^{1/2}} + q^\varepsilon P^{3/4} R^{1/2} (LL_2)^{1/4} (q + P^2 |q\alpha - a|)^{1/8} \quad (k = 2).$$

We note that when $q + P^k |q\alpha - a| > P^2 R^{-4} (LL_2)^{-2}$, the estimates supplied by this theorem are worse than trivial. There is thus no loss of generality in assuming henceforth that $R \leq P^{1/2} (LL_2)^{-1/2}$, and that $1 + P^k |\alpha - a/q| \leq P^2$. We then have

$$\mathcal{L} = \log(2 + P^k |\alpha - a/q|) \ll L. \quad (1.2)$$

This estimate allows one to remove the annoying presence of \mathcal{L} from the bounds provided by Theorem 1.1. Alternatively, one may observe that for each $\varepsilon > 0$ one has $\mathcal{L} \ll (1 + P^k |\alpha - a/q|)^\varepsilon$. Hence, subject to the hypotheses of Theorem 1.1, one has

$$g_\nu(\alpha) \ll \frac{q^\varepsilon \kappa(q)^{1/2} P L L_2^{1/2}}{(1 + P^k |\alpha - a/q|)^{1/2-\varepsilon}} + q^\varepsilon P^{3/4} R^{1/2} (LL_2)^{1/4} (q + P^k |q\alpha - a|)^{1/8}, \quad (1.3)$$

a bound that often suffices in practice.

In order to put into context previous work on this subject, we introduce a fairly general Hardy-Littlewood dissection of the unit interval. Let Q be a real number with $1 \leq Q \leq P^{k/2}$. When $q \in \mathbb{N}$ and $a \in \mathbb{Z}$, we take

$$\mathfrak{M}(q, a; Q) = \{\alpha \in [0, 1) : |q\alpha - a| \leq QP^{-k}\}.$$

We write $\mathfrak{M}(Q)$ for the union of the intervals $\mathfrak{M}(q, a; Q)$ over coprime integers a and q with $0 \leq a \leq q \leq Q$, and we put $\mathfrak{m}(Q) = [0, 1) \setminus \mathfrak{M}(Q)$. In applications of the circle method, the most frequently encountered model Hardy-Littlewood dissection is that into major arcs $\mathfrak{M}(P)$ and minor arcs $\mathfrak{m}(P)$ thus defined. On noting that whenever $\alpha \in \mathfrak{M}(Q)$,

one has $q + P^k|q\alpha - a| \leq 2Q$, we discern from Theorem 1.1 and the bound (1.2) that whenever $\alpha \in \mathfrak{M}(q, a; P) \subseteq \mathfrak{M}(P)$, one has

$$g_\nu(\alpha; P, R) \ll \frac{q^\varepsilon \kappa(q)^{1/2} P (\log P)^{3/2+\varepsilon}}{(1 + P^k|\alpha - a/q|)^{1/2}} + P^{7/8+\varepsilon} R^{1/2}. \quad (1.4)$$

In the special case $k = 3$, a related conclusion is available from [1, Lemma 2.2], though with the exponent $9/10$ in place of $7/8$. We note, however, that the estimate obtained in [1] is applicable for a wider set of parameters α . Rather than attempt to generalise the argument underlying the proof of this lemma to general exponents k , which would in any case yield substantially weaker estimates than we derive in (1.4), we instead turn to earlier work of the second author joint with Vaughan [13] for inspiration. As enhanced in [6, Lemma 5.4], this approach shows that whenever $2 \leq R \leq M \leq P$ and $q + P^k|q\alpha - a| \leq TM$, then

$$g(\alpha; P, R) \ll q^\varepsilon (\log P)^3 \left(\frac{P}{(q + P^k|q\alpha - a|)^{1/(2k)}} + (PMR)^{1/2} + PR^{1/2} \left(\frac{T}{M} \right)^{1/4} \right).$$

Experts will recognise that a factor $q^{-1/(2k)}$ in this estimate is replaced in Theorem 1.1 by a factor $\kappa(q)^{1/2}$, an enhancement that offers substantially stronger control of mean value estimates restricted to sets of major arcs. There is also a useful improvement in the dependence on $1 + P^k|\alpha - a/q|$.

Very recently, using methods quite different from those that we apply here, Shparlinski [7, Theorem 1.3] estimated $g(\alpha; P, R)$ in the special case where the argument is a rational number with prime denominator p . In his work, it is assumed that $p > (4P)^{3/4}$. Then, for $a \in \mathbb{Z}$ with $(a, p) = 1$ and $2 \leq R \leq p^{1/2}$, it is shown that for each $\varepsilon > 0$ one has

$$g(a/p; P, R) \ll p^{5/48+\varepsilon} P^{29/32} R^{3/104}.$$

Note that this estimate is worse than the trivial bound when $p > P^{9/10}$ so that we may concentrate on the situation with $(4P)^{3/4} < p \leq P^{9/10}$. In this range, Theorem 1.1 supplies the estimate

$$g(a/p; P, R) \ll Pp^{\varepsilon-1/4} + P^{3/4+\varepsilon} p^{1/8} R^{1/2} \ll P^{3/4+\varepsilon} p^{1/8} R^{1/2}.$$

In the most interesting case where R is a very small power of P , our bound is decidedly smaller, and our bound is non-trivial for p almost as large as P^2 .

We turn our attention now to the factor $\log P$ occurring in the first term on the right hand side of our estimate for $g_\nu(\alpha)$ in Theorem 1.1. This has the potential to prevent the immediate use of this theorem in pursuit of near optimal bounds for moments of smooth Weyl sums over major arcs of the shape $\mathfrak{M}(Q)$, when Q is no larger than a modest power of $\log P$. In §3 we obtain estimates for smooth Weyl sums that eliminate this deficiency. The bounds feature Euler's totient, denoted $\phi(q)$, as well as the closely associated multiplicative function

$$\psi(q) = q/\phi(q) = \prod_{p|q} (1 - 1/p)^{-1}.$$

We note for future reference that $\psi(q) = O(\log \log q)$ for $q \geq 3$ (see [5, Theorem 2.9], for example).

Theorem 1.2. *Let k be a natural number with $k \geq 2$, and let ν be a real number with $\nu > 1$. Suppose that R and P are real numbers with $2 \leq R \leq P^{1/2}$. Then, for each $A \geq 1$, there is a positive number c with the following property. Whenever $a \in \mathbb{Z}$, $q \in \mathbb{N}$ and $\alpha \in \mathbb{R}$ satisfy $(a, q) = 1$ and $q \leq (\log P)^A$, one has*

$$g_\nu(\alpha; P, R) \ll \frac{\kappa(q)\psi(q)P}{1 + P^k|\alpha - a/q|} + P \exp(-c(\log P)^{1/2})(1 + P^k|\alpha - a/q|)$$

and

$$g(\alpha; P, R) \ll \frac{\kappa(q)\psi(q)P}{(1 + P^k|\alpha - a/q|)^{1/k}} + P \exp(-c(\log P)^{1/2})(1 + P^k|\alpha - a/q|).$$

By way of comparison, the earlier work of the second author joint with Vaughan [13, Lemma 8.5] provides a conclusion of similar type, though with the factor $\kappa(q)$ replaced by $q^{\varepsilon-1/k}$. While this previous approach certainly addresses the issue raised in the preamble to Theorem 1.2, our new conclusion permits control of major arc moments in which the number of generating functions can be very nearly halved.

Following the proof of an auxiliary pruning lemma in §4, we obtain estimates for mean values of the smooth Weyl sum $g_\nu(\alpha; P, R)$ on sets of major arcs in §5. These estimates provide near optimal upper bounds for corresponding moments of the exponential sum $g(\alpha; P, R)$.

Theorem 1.3. *Let k and t be natural numbers with $k \geq 3$ and $t \geq \lfloor k/2 \rfloor$. Furthermore, let $\omega < k/2$ and ω' be positive numbers with*

$$\omega < \frac{2t + 4}{t + 10} \quad \text{and} \quad \omega' < \frac{2k}{k + 4},$$

and put $\Omega = \min\{\omega, \omega'\}$. Then, for any $\varepsilon > 0$ there is a real number $\eta > 0$ with the property that whenever $2 \leq R \leq P^\eta$, one has

$$\int_{\mathfrak{M}(P^\omega)} |g(\alpha; P, R)|^{2t+4} d\alpha \ll P^{2t+4-k+\varepsilon}. \quad (1.5)$$

Moreover, when u is a real number with $u > 2t + 4$, then provided that

$$1 \leq Q \leq P^\Omega \quad \text{and} \quad \tau < (u - 2t - 4)/(2k),$$

one has

$$\int_{\mathfrak{M}(P^\Omega) \setminus \mathfrak{M}(Q)} |g(\alpha; P, R)|^u d\alpha \ll P^{u-k} Q^{-\tau}.$$

This theorem delivers the near optimal estimate (1.5). Moreover, one may allow the exponent ω to approach 2 as $t \rightarrow \infty$. By contrast, earlier treatments would restrict ω to satisfy the constraint $0 \leq \omega \leq 1$. Furthermore, when $k \geq 4$ one has $t \geq 2$, and in such circumstances ω can be nearly as large as $2/3$. One finds that ω can be taken to be any number with $0 \leq \omega < 1$ as soon as $t \geq 6$, and this is assured when $k \geq 12$. When $k = 3$ and $t = 1$, the estimate (1.5) was demonstrated in [1, Corollary 3.2] with $\omega < 2/5$. Our new result covers the range $\omega < 6/11$.

We next record a corollary to Theorem 1.3 that is employed in our recent work [3] on the representation of integers as sums of a square and a number of k -th powers. Here, and throughout, we define

$$\mathfrak{N}(Q) = \mathfrak{M}(Q) \setminus \mathfrak{M}(Q/2). \quad (1.6)$$

Corollary 1.4. *Let k be a natural number with $k \geq 3$, put $t = \lfloor k/2 \rfloor$, and suppose that u is a real number with $u > 2t + 4$. Furthermore, let τ be a real number with $0 < \tau < (u - 2t - 4)/(2k)$. Then, there is a real number $\eta > 0$ with the property that whenever $2 \leq R \leq P^\eta$ and $1 \leq Q \leq P^{1/2}$, one has*

$$\int_{\mathfrak{N}(Q)} |g(\alpha; P, R)|^u d\alpha \ll P^{u-k} Q^{-\tau}.$$

We finish this memoir in §6 with an account of the consequences of our new estimates for Waring's problem. We are able to establish lower bounds for the contribution of the major arcs in Waring's problem of the expected order, provided that the number of variables available is at least $2\lfloor k/2 \rfloor + 5$, when $k \geq 3$. In the classical situation in which variables are not restricted, and in particular are not restricted to be smooth, analogous conclusions would require at least $k+2$ variables. Previous work using smooth Weyl sums, meanwhile, would require at least $2k+3$ variables. The superiority of our new conclusions is clear. We refer the reader to Theorems 6.1 and 6.2 for details of these new results.

In this memoir, our basic parameter is P , a sufficiently large positive number. We adopt the usual convention that, whenever a statement involves the letter ε , then it is asserted that the statement holds for any positive value of ε . Implicit constants in the notations of Vinogradov and Landau may depend on ε , as well as ambient parameters implicitly fixed, such as k and u . We write $p^h \parallel n$ to denote that $p^h \mid n$, but $p^{h+1} \nmid n$.

The authors are grateful to the referees for their careful reading and comments useful in improving the exposition of this paper.

2. AN ENHANCED MAJOR ARC ESTIMATE

Our goal in this section is the proof of Theorem 1.1. In order to establish the bounds for $g_\nu(\alpha; P, R)$ recorded therein, we adapt the analysis of [13, §7], taking advantage of more precise auxiliary estimates deemed unnecessary in this earlier treatment. Our account, nonetheless, follows closely the argument of the proof of [13, Lemma 7.2]. Throughout, in order to ease our exposition, we make use of the notation recorded in (1.1). We begin with a lemma concerning the multiplicative function $\kappa(q)$ required later in this section.

Lemma 2.1. *Let $q \in \mathbb{N}$ and suppose that $d \mid q$. Let e_0 be a divisor of d , and write $d/e_0 = d_1 d_2^2 \cdots d_k^k$, where d_1, \dots, d_{k-1} are pairwise coprime and squarefree. Then*

$$\kappa(q/d) \leq k^{4\omega(q)} \kappa(q) e_0 d_1 \cdots d_k.$$

Proof. We claim that whenever a and b are non-negative integers with $a + b \geq 1$, then

$$\kappa(p^a) \kappa(p^b) \leq k^2 \kappa(p^{a+b}). \quad (2.1)$$

In order to establish this assertion, we note that in view of the definition of $\kappa(q)$, it suffices to consider the situation in which $1 \leq a, b \leq k$. We divide into three cases. When $2 \leq a, b \leq k$, we have

$$\kappa(p^a) \kappa(p^b) = p^{-2} \quad \text{and} \quad \kappa(p^{a+b}) \geq p^{-2}.$$

When $a = 1$ and $2 \leq b \leq k$, and also when $b = 1$ and $2 \leq a \leq k$, we have

$$\kappa(p^a) \kappa(p^b) = k p^{-3/2} \quad \text{and} \quad \kappa(p^{a+b}) \geq p^{-3/2}.$$

Finally, when $a = b = 1$, one has

$$\kappa(p^a)\kappa(p^b) = k^2 p^{-1} \quad \text{and} \quad \kappa(p^{a+b}) = p^{-1}.$$

This confirms the upper bound (2.1) in all cases.

The multiplicative property of $\kappa(q)$ now confirms that whenever $q \in \mathbb{N}$ and $d \mid q$, one has

$$\kappa(d)\kappa(q/d) \leq k^{2\omega(q)}\kappa(q),$$

and similarly

$$\kappa(e_0)\kappa(d_1 d_2^2 \cdots d_k^k) \leq k^{2\omega(d)}\kappa(d).$$

But $\kappa(e_0) \geq e_0^{-1}$ and $\kappa(d_1 d_2^2 \cdots d_k^k) \geq (d_1 d_2 \cdots d_k)^{-1}$. Hence, we conclude that

$$(e_0 d_1 d_2 \cdots d_k)^{-1} \kappa(q/d) \leq k^{2\omega(d)+2\omega(q)}\kappa(q) \leq k^{4\omega(q)}\kappa(q).$$

The conclusion of the lemma follows immediately. \square

Next, we provide two simple estimates for auxiliary sums.

Lemma 2.2. *Suppose that X and J are positive numbers. Then*

$$\sum_{1 \leq j \leq J} (1 + jX)^{-1} \leq 2J(1 + XJ)^{-1}(1 + \log(1 + JX)).$$

Proof. When X and J are positive numbers with $X \leq J^{-1}$, then

$$\sum_{1 \leq j \leq J} (1 + jX)^{-1} \leq J \leq 2J(1 + XJ)^{-1}.$$

When instead $J^{-1} < X$, then

$$\begin{aligned} \sum_{1 \leq j \leq J} (1 + jX)^{-1} &\leq X^{-1}(1 + \log(1 + JX)) \\ &\leq 2J(1 + XJ)^{-1}(1 + \log(1 + JX)). \end{aligned}$$

The desired conclusion therefore follows in both cases. \square

Lemma 2.3. *Let k and J be integers with $k \geq 2$ and $1 \leq 2^J \leq P$, and let λ be a real number with $1/k < \lambda \leq 1$. When j is a non-negative integer, put $X_j = 2^{-j}P$. Then, for all real numbers β , one has*

$$\sum_{0 \leq j \leq J} X_j (1 + X_j^k |\beta|)^{-\lambda} \ll \frac{P}{(1 + P^k |\beta|)^{1/k}} \quad (2.2)$$

while

$$\sum_{0 \leq j \leq J} X_j (1 + X_j^k |\beta|)^{-1/k} \ll \frac{P \log(2 + P^k |\beta|)}{(1 + P^k |\beta|)^{1/k}}. \quad (2.3)$$

Proof. Suppose that $1/k \leq \lambda \leq 1$, and let K_λ denote the sum on the left hand side of (2.2) (or (2.3) when $\lambda = 1/k$). If $|\beta| \leq P^{-k}$, then

$$K_\lambda \leq \sum_{j \geq 0} 2^{-j} P \leq 2P,$$

which is acceptable. Next suppose that $|\beta| \geq 1$. Then, writing J_0 for the largest integer satisfying $2^{J_0} \leq P$, we find that

$$K_\lambda \leq \sum_{2^j \leq P} X_j^{1-\lambda k} |\beta|^{-\lambda} = P^{1-\lambda k} |\beta|^{-\lambda} \sum_{0 \leq j \leq J_0} 2^{(\lambda k - 1)j}.$$

When $\lambda > 1/k$, we conclude that $K_\lambda \ll |\beta|^{-\lambda}$, which is superior to our claim in (2.2). Further, noting that $J_0 \leq 2 \log P$, we find that $K_{1/k} \ll |\beta|^{-1/k} \log P$. This confirms (2.3) when $|\beta| \geq 1$.

We may now suppose that $P^{-k} \leq |\beta| \leq 1$. Then, there is an integer I with $I \leq J_0$ and

$$2^{-I} P \geq |\beta|^{-1/k} \geq 2^{-I-1} P.$$

The contribution of the summands in K_λ where $j > I$ (if any) does not exceed

$$\sum_{j > I} 2^{-j} P \leq 2^{-I} P \leq 2|\beta|^{-1/k}.$$

When $j \leq I$, one has $X_j^k |\beta| \geq 1$, and hence we see that

$$K_\lambda \leq 2|\beta|^{-1/k} + |\beta|^{-\lambda} \sum_{0 \leq j \leq I} X_j^{1-\lambda k} \leq 2|\beta|^{-1/k} + |\beta|^{-\lambda} P^{1-\lambda k} \sum_{0 \leq j \leq I} 2^{(\lambda k - 1)j}.$$

We now proceed as in the case $|\beta| \geq 1$ to estimate the rightmost sum over j , and this completes the proof of the lemma. \square

We are now equipped to prove our first estimate for $g_\nu(\alpha; P, R)$. In this context, we remind the reader of the notation introduced in (1.1).

Lemma 2.4. *Let k be a natural number with $k \geq 2$, and let ν be a real number with $\nu > 1$. Suppose that R, P and M are real numbers with*

$$2 \leq R \leq M \leq P/\nu.$$

Then, whenever $\alpha \in \mathbb{R}$ and $a \in \mathbb{Z}$, $q \in \mathbb{N}$ are coprime, one has

$$\begin{aligned} g_\nu(\alpha; P, R) &\ll \frac{q^\varepsilon \kappa(q)^{1/2} PL(L_2 \mathcal{L})^{1/2}}{(1 + P^k |\alpha - a/q|)^{1/2}} + q^\varepsilon (LL_2)^{1/2} (PMR)^{1/2} \\ &\quad + q^\varepsilon P(R/M)^{1/2} (q + P^k |q\alpha - a|)^{1/4}. \end{aligned} \quad (2.4)$$

Proof. In this discussion, we provide certain details of the account of the proof of [13, Lemma 7.2] that could be suppressed at the cost of opacity of exposition. We begin with an initial decomposition of the exponential sum $g_\nu(\alpha; P, R)$ into pieces equipped with a bilinear structure. For each prime number π with $\pi \leq R$, we define

$$\mathcal{B}_\pi = \{v \in \mathcal{A}(M\pi, R) : v > M, \pi | v, \text{ and } p | v \Rightarrow p \geq \pi\}.$$

Then, as a consequence of [10, Lemma 10.1], one has

$$g_\nu(\alpha; P, R) = \sum_{\pi \leq R} \sum_{v \in \mathcal{B}_\pi} \sum_{u \in \mathcal{A}_\nu(P/v, \pi)} e(\alpha(uv)^k). \quad (2.5)$$

Next, write

$$\mathcal{W} = \{(2^j, 2^i M) : i \geq 0, j \geq -1, 2^i < R \text{ and } P/\nu < 2^{i+j} M < P\},$$

and define

$$S_1(U, V) = \sum_{\pi \leq R} \sum_{\substack{V < v \leq 2V \\ v \in \mathcal{B}_\pi}} \left| \sum_{\substack{u \in \mathcal{A}_\nu(P/v, \pi) \\ U < u \leq 2U}} e(\alpha(uv)^k) \right|.$$

Then, much as in the analogous treatment of [13], we see that

$$\left| \sum_{\pi \leq R} \sum_{v \in \mathcal{B}_\pi} \sum_{u \in \mathcal{A}_\nu(P/v, \pi)} e(\alpha(uv)^k) \right| \leq \sum_{(U, V) \in \mathcal{W}} S_1(U, V). \quad (2.6)$$

We note for future reference that for each pair $(U, V) \in \mathcal{W}$, one has

$$M \leq V < MR \quad \text{and} \quad P/(\nu V) < U < P/V. \quad (2.7)$$

Furthermore, since the elements of \mathcal{B}_π are integers divisible by π , all of whose prime divisors are at least as large as π , it follows from [8, Chapter III.6, Theorem 3] that

$$\sum_{\pi \leq R} \sum_{\substack{V < v \leq 2V \\ v \in \mathcal{B}_\pi}} 1 \ll \sum_{\pi \leq R} \frac{V}{\pi \log \pi} \ll V. \quad (2.8)$$

By Cauchy's inequality, one has

$$S_1(U, V)^2 \leq \left(\sum_{\pi \leq R} \sum_{\substack{V < v \leq 2V \\ v \in \mathcal{B}_\pi}} 1 \right) \sum_{\pi \leq R} \sum_{\substack{V < v \leq 2V \\ v \in \mathcal{B}_\pi}} \left| \sum_{\substack{u \in \mathcal{A}_\nu(P/v, \pi) \\ U < u \leq 2U}} e(\alpha(uv)^k) \right|^2.$$

Thus, on isolating the implicit diagonal terms, we discern by means of (2.8) that

$$|S_1(U, V)|^2 \ll V^2 U + V |S_2(U, V)|, \quad (2.9)$$

where

$$S_2(U, V) = \sum_{\pi \leq R} \sum_{\substack{u_1, u_2 \in \mathcal{A}_\nu(P/V, \pi) \\ U < u_1 < u_2 \leq 2U}} T_1(\alpha \pi^k (u_2^k - u_1^k)), \quad (2.10)$$

in which we write

$$T_1(\gamma) = \sum_{\substack{V/\pi < t \leq 2V/\pi \\ P/(\nu \pi u_1) < t \leq P/(\pi u_2)}} e(\gamma t^k).$$

On collecting together the estimates (2.5), (2.6) and (2.9), we may conclude thus far that

$$g_\nu(\alpha; P, R) \ll \sum_{(U, V) \in \mathcal{W}} (VU^{1/2} + V^{1/2} |S_2(U, V)|^{1/2}). \quad (2.11)$$

It is at this point that our treatment begins to diverge further from the path laid down in [13]. We again view $S_2(U, V)$, as defined in (2.10), as an exponential sum over k -th powers of t . For a typical summand on the right hand side of (2.10), we write

$$D = (q, \pi^k (u_2^k - u_1^k)), \quad (2.12)$$

and then put

$$c = a\pi^k (u_2^k - u_1^k)/D, \quad r = q/D, \quad \gamma = \alpha \pi^k (u_2^k - u_1^k). \quad (2.13)$$

Thus, we have

$$\gamma - \frac{c}{r} = \pi^k (u_2^k - u_1^k) \left(\alpha - \frac{a}{q} \right), \quad (2.14)$$

with

$$(c, r) = (a\pi^k(u_2^k - u_1^k), q)/D = (a, q/D) = 1.$$

We now apply [12, Theorem 4.1]. Thus, in view of Lemmata 4.3, 4.4, 4.5 and 6.2 of [12], we find that

$$T_1(\gamma) \ll \frac{\kappa(r)V/\pi}{1 + (V/\pi)^k|\gamma - c/r|} + r^{1/2+\varepsilon}(1 + (V/\pi)^k|\gamma - c/r|)^{1/2}. \quad (2.15)$$

Next, write

$$S_3(U, V) = \sum_{\pi \leq RU < u_1 < u_2 \leq 2U} \sum_{\pi \leq RU < u_1 < u_2 \leq 2U} \frac{\kappa(q/(q, \pi^k(u_2^k - u_1^k)))}{\pi(1 + V^k(u_2^k - u_1^k)|\alpha - a/q|)}. \quad (2.16)$$

Then on recalling (2.10), we deduce from (2.12)-(2.15) that

$$S_2(U, V) \ll VS_3(U, V) + RU^2q^\varepsilon(q + (UV)^k|q\alpha - a|)^{1/2}.$$

On substituting this estimate into (2.11) and recalling (2.7), we infer thus far that

$$g_\nu(\alpha; P, R) \ll (PMR)^{1/2} + \sum_{(U, V) \in \mathscr{W}} VS_3(U, V)^{1/2} + P(R/M)^{1/2}q^\varepsilon(q + P^k|q\alpha - a|)^{1/4}. \quad (2.17)$$

We next analyse the sum $S_3(U, V)$ defined in (2.16). First, by separately considering the contributions arising from summands in which $\pi | q$ and $\pi \nmid q$, and applying the bound $u_2^k - u_1^k \geq (u_2 - u_1)U^{k-1}$, we find that

$$S_3(U, V) \ll L_2q^\varepsilon \sum_{U < u_1 < u_2 \leq 2U} \frac{\kappa(q/(q, u_2^k - u_1^k))}{1 + U^{k-1}V^k(u_2 - u_1)|\alpha - a/q|}.$$

Here, we have made use of the bound $\kappa(\pi^h) \geq k^{-1}\pi^{-1}\kappa(\pi^{h-1})$, valid for $0 \leq l \leq k$ and $h \geq l$.

Now write $d = (q, u_2^k - u_1^k)$ and $\beta = |\alpha - a/q|$. Also, put $m = (u_1, u_2)$ and $t_i = u_i/m$ ($i = 1, 2$). Then, we obtain the upper bound

$$S_3(U, V) \ll L_2q^\varepsilon \sum_{d|q} \kappa(q/d) \sum_{1 \leq m \leq 2U} \sum_{\substack{U/m < t_1 < t_2 \leq 2U/m \\ (t_1, t_2) = 1 \\ d|m^k(t_2^k - t_1^k)}} (1 + U^{k-1}V^km(t_2 - t_1)\beta)^{-1}.$$

We follow the analogous argument of [13] once again. For a given pair of integers d and m occurring in the latter sum, we put $d_0 = (d, m^k)$ and $e_0 = d/d_0$. Then we see that

$$S_3(U, V) \ll L_2q^\varepsilon \sum_{d|q} \kappa(q/d) \sum_{d_0e_0=d} \sum_{\substack{1 \leq m \leq 2U \\ d_0|m^k}} S_4(U, V), \quad (2.18)$$

in which we write

$$S_4(U, V) = \sum_{\substack{U/m < t_1 < t_2 \leq 2U/m \\ (t_1, t_2) = 1 \\ e_0 | (t_2^k - t_1^k)}} (1 + U^{k-1}V^km(t_2 - t_1)\beta)^{-1}.$$

This sum is analysed (under the name S_5) in the discussion following [13, equation (7.8)], though in the latter case the inner sum has argument with exponent $-1/k$ in place of -1 .

This difference in detail does not impact the argument that follows in any material way, and thus we may conclude that

$$S_4(U, V) \ll q^\varepsilon \sum_{e_1 f_1 = e_0} \sum_{1 \leq j \leq U/(m e_1)} \left(\frac{U}{m f_1} + 1 \right) \left(1 + U^{k-1} V^k m j e_1 \beta \right)^{-1}. \quad (2.19)$$

By applying the upper bound supplied by Lemma 2.2 within (2.19), and then recalling (2.7), we find that

$$\begin{aligned} S_4(U, V) &\ll q^\varepsilon \sum_{e_1 f_1 = e_0} \left(\frac{U}{m f_1} + 1 \right) \left(\frac{U}{m e_1} \right) \frac{1 + \log(1 + (UV)^k \beta)}{1 + (UV)^k \beta} \\ &\ll \mathcal{L} q^\varepsilon \sum_{e_1 f_1 = e_0} \left(\frac{U^2}{m^2 e_0} + \frac{U}{m e_1} \right) (1 + P^k \beta)^{-1}. \end{aligned}$$

Since $e_0 \mid q$, we therefore infer that

$$S_4(U, V) \ll \mathcal{L} q^{2\varepsilon} \left(\frac{U^2}{m^2 e_0} + \frac{U}{m} \right) (1 + P^k \beta)^{-1}.$$

By substituting this upper bound into (2.18), therefore, we deduce that

$$S_3(U, V) \ll \frac{L_2 \mathcal{L} q^{3\varepsilon}}{1 + P^k \beta} \sum_{d \mid q} \kappa(q/d) \sum_{d_0 e_0 = d} \left(\left(\sum_{\substack{1 \leq m \leq 2U \\ d_0 \mid m^k}} \frac{U^2}{m^2 e_0} \right) + LU \right).$$

We now write $d_0 = d_1 d_2^2 \cdots d_k^k$, where d_1, \dots, d_{k-1} are squarefree and pairwise coprime, and we recall the conclusion of Lemma 2.1. Then we see that

$$S_3(U, V) \ll LL_2 q^{4\varepsilon} U + \frac{L_2 \mathcal{L} q^{4\varepsilon} \kappa(q)}{1 + P^k \beta} T_2(q) U^2,$$

where

$$T_2(q) = \sum_{d \mid q} \sum_{e_0 d_1 d_2^2 \cdots d_k^k = d} e_0 d_1 \cdots d_k \sum_{1 \leq n \leq 2U/(d_1 \cdots d_k)} \frac{1}{(n d_1 \cdots d_k)^2 e_0}.$$

But we have

$$T_2(q) \ll \sum_{d \mid q} \sum_{e_0 d_1 \cdots d_k \mid d} \frac{1}{d_1 \cdots d_k} \ll q^\varepsilon,$$

and hence we conclude that

$$S_3(U, V) \ll LL_2 q^{4\varepsilon} U + \frac{L_2 \mathcal{L} q^{5\varepsilon} \kappa(q) U^2}{1 + P^k \beta}. \quad (2.20)$$

On substituting (2.20) into (2.17), and again recalling (2.7), we arrive at the upper bound

$$\begin{aligned} g_\nu(\alpha; P, R) &\ll \frac{q^{3\varepsilon} \kappa(q)^{1/2} P (\log R) (L_2 \mathcal{L})^{1/2}}{(1 + P^k |\alpha - a/q|)^{1/2}} + (LL_2)^{1/2} q^{2\varepsilon} (PMR)^{1/2} \\ &\quad + P(R/M)^{1/2} q^\varepsilon (q + P^k |q\alpha - a|)^{1/4}. \end{aligned}$$

The upper bound (2.4) asserted in the statement of the lemma now follows. \square

We are now equipped to complete the proof of our first theorem.

The proof of Theorem 1.1. As already detailed in the discussion leading to (1.2), it suffices to establish Theorem 1.1 with the additional hypotheses

$$R \leq P^{1/2}(LL_2)^{-1/2} \quad \text{and} \quad q + P^k|q\alpha - a| \leq P^2R^{-4}(LL_2)^{-2}.$$

We may consequently apply Lemma 2.4 with

$$M = (LL_2)^{-1/2}P^{1/2}(q + P^k|q\alpha - a|)^{1/4},$$

since in these circumstances one has $R \leq M < P/\nu$ whenever P is sufficiently large. With this choice for M , the second and third terms on the right hand side of (2.4) satisfy the bound

$$\begin{aligned} q^\varepsilon(LL_2)^{1/2}(PMR)^{1/2} + q^\varepsilon P(R/M)^{1/2}(q + P^k|q\alpha - a|)^{1/4} \\ \ll q^\varepsilon(LL_2)^{1/4}P^{3/4}R^{1/2}(q + P^k|q\alpha - a|)^{1/8}, \end{aligned}$$

and the first conclusion of Theorem 1.1 is therefore immediate from (2.4).

The second conclusion of Theorem 1.1 requires that we remove the condition on the sum $g_\nu(\alpha; P, R)$ that its summands come from a truncated set of smooth numbers. Assume the hypotheses of the statement of the theorem, and suppose that X is a parameter with $P^{3/4} \leq X \leq P$. Then $R \leq P^{1/2} \leq X$, according to current hypotheses. With the first conclusion of the theorem already in hand, we may apply its corollary (1.3) with X in place of P . This supplies the bound

$$g_2(\alpha; X, R) \ll \frac{q^\varepsilon \kappa(q)^{1/2} X L L_2^{1/2}}{(1 + X^k |\alpha - a/q|)^{3/8}} + q^\varepsilon (LL_2)^{1/4} X^{3/4} R^{1/2} (q + X^k |q\alpha - a|)^{1/8}.$$

We put $J = \lceil (\log P)/(4 \log 2) \rceil$, and sum the contributions from $g_2(\alpha; X, R)$ for $X = 2^{-j}P$, with $0 \leq j < J$. Then, by applying Lemma 2.3 with $\lambda = 3/8$, and making use of the trivial estimate $g_2(\alpha; 2^{-j}P, R) = O(P^{3/4})$, we find that when $k \geq 3$, one has

$$g(\alpha; P, R) \ll P^{3/4} + \frac{q^\varepsilon \kappa(q)^{1/2} P L L_2^{1/2}}{(1 + P^k |\alpha - a/q|)^{1/k}} + q^\varepsilon (LL_2)^{1/4} P^{3/4} R^{1/2} (q + P^k |q\alpha - a|)^{1/8}.$$

When $k = 2$, the argument is different. We invoke the estimate for $g_2(\alpha; P, R)$ that is provided by Theorem 1.1. Then, since $X \leq P$, we have

$$g_2(\alpha; X, R) \ll \frac{q^\varepsilon \kappa(q)^{1/2} X L (L_2 \mathcal{L})^{1/2}}{(1 + X^2 |\alpha - a/q|)^{1/2}} + q^\varepsilon (LL_2)^{1/4} X^{3/4} R^{1/2} (q + P^2 |q\alpha - a|)^{1/8}.$$

Here L , L_2 and \mathcal{L} are still given by (1.1). We now sum over $X = 2^{-j}P$ as before, but now the application of Lemma 2.3 produces an extra factor \mathcal{L} as one sums the first term on the right hand side. The proof of Theorem 1.1 is now complete. \square

3. ANOTHER ENHANCED MAJOR ARC ESTIMATE

The conclusion of Theorem 1.1 provides relatively powerful estimates for $g_\nu(\alpha; P, R)$ whenever $a \in \mathbb{Z}$, $q \in \mathbb{N}$ and $\alpha \in \mathbb{R}$ satisfy $(a, q) = 1$ and the condition that $q + P^k|q\alpha - a|$ is neither too small nor larger than about P^2 . However, when this latter quantity is smaller than $(\log P)^5$, these estimates are worse than trivial. Our goal in this section is to address these very small values of $q + P^k|q\alpha - a|$ via an analogue of [13, Lemma 8.5]. Fortunately, the argument of the proof of the latter conclusion requires relatively little refinement in order that our more precise estimates be confirmed.

We begin with some auxiliary lemmata.

Lemma 3.1. *Suppose that X and Y are real numbers with $0 < Y < X$, and that $f : [Y, X] \rightarrow \mathbb{R}$ is monotonic and differentiable on (Y, X) . Suppose further that f' is continuous on (Y, X) , and that $f'(Y+)$ and $f'(X-)$ both exist. Then, for all real numbers γ , one has*

$$\left| \int_Y^X f(t)e(\gamma t^k) dt \right| \leq (|f(X-)| + |f(Y+)|) \frac{2X}{1 + Y^k |\gamma|}.$$

Proof. The conclusion is trivial when $|\gamma| \leq Y^{-k}$, so we may assume henceforth that $|\gamma| > Y^{-k}$. By integrating by parts, we find that

$$\begin{aligned} \int_Y^X f(t)e(\gamma t^k) dt &= \frac{f(X-)e(\gamma X^k)}{2\pi i \gamma k X^{k-1}} - \frac{f(Y+)e(\gamma Y^k)}{2\pi i \gamma k Y^{k-1}} \\ &\quad + \int_Y^X \left(\frac{(k-1)f(t)}{kt^k} - \frac{f'(t)}{kt^{k-1}} \right) \frac{e(\gamma t^k)}{2\pi i \gamma} dt, \end{aligned}$$

and hence

$$\left| \int_Y^X f(t)e(\gamma t^k) dt \right| \leq \frac{|f(X-)| + |f(Y+)|}{2\pi k |\gamma| Y^{k-1}} + \frac{1}{2\pi k |\gamma|} \int_Y^X (k-1) \frac{|f(t)|}{t^k} + \frac{|f'(t)|}{t^{k-1}} dt. \quad (3.1)$$

But

$$\begin{aligned} \int_Y^X \frac{(k-1)|f(t)|}{t^k} dt &\leq (|f(X-)| + |f(Y+)|) \int_Y^X \frac{k-1}{t^k} dt \\ &\leq (|f(X-)| + |f(Y+)|) Y^{1-k} \end{aligned}$$

and

$$\int_Y^X \frac{|f'(t)|}{t^{k-1}} dt \leq \left| \int_Y^X f'(t) dt \right| Y^{1-k} \leq (|f(X-)| + |f(Y+)|) Y^{1-k}.$$

On substituting the last two bounds into (3.1), the desired inequality is readily confirmed in the case $|\gamma| > Y^{-k}$. \square

The next lemma summarises well-known properties of the sum

$$W(q, a) = \sum_{\substack{r=1 \\ (q,r)=1}}^q e(ar^k/q). \quad (3.2)$$

Lemma 3.2. *For each natural number k with $k \geq 2$, the exponential sum $W(q, a)$ satisfies the following properties.*

- (i) *Define the integer θ via the relation $p^\theta \parallel k$, and suppose that a is an integer with $(a, p) = 1$. Then, when $p \geq 3$ and $t \geq \theta + 2$, and also when $p = 2$ and $t \geq \theta + 3$, one has $W(p^t, a) = 0$.*
- (ii) *For all prime numbers p , one has*

$$|W(p, a)| \leq 1 + ((k, p-1) - 1) \sqrt{p}. \quad (3.3)$$

- (iii) *When $1 \leq \tau < t$, one has*

$$W(p^t, ap^\tau) = p^\tau W(p^{t-\tau}, a). \quad (3.4)$$

(iv) When r_1 and r_2 are coprime natural numbers, then for all integers c one has

$$W(r_1 r_2, c) = W(r_1, cr_2^{k-1})W(r_2, cr_1^{k-1}). \quad (3.5)$$

Proof. The vanishing property in the first clause of the lemma is [4, Lemma 8.3]. In order to establish (3.3), we write

$$W(p, a) = -1 + \sum_{x=1}^p e(ax^k/p)$$

and apply [12, Lemma 4.3]. The identity (3.4) follows immediately from the definition of $W(p^t, c)$. Finally, the quasi-multiplicative property (3.5) is established as the first clause of [4, Lemma 8.1]. \square

In the proof of Theorem 1.2 we require an estimate for the sum

$$\mathcal{S}(q, a) = \frac{1}{q} \sum_{d|q} \psi(d) |W(d, a(q/d)^{k-1})|, \quad (3.6)$$

in which $\psi(d)$ denotes the previously defined multiplicative function $d/\phi(d)$.

Lemma 3.3. *Let k be a natural number with $k \geq 2$. Then, whenever $q \in \mathbb{N}$ and $a \in \mathbb{Z}$ are coprime, one has $\mathcal{S}(q, a) \leq 6k\kappa(q)\psi(q)$.*

Proof. We first consider the case where q is a power of the prime p , say $q = p^l$ with $l \geq 1$. We suppose throughout that $p \nmid a$. By (3.6), one has

$$\mathcal{S}(p^l, a) = \sum_{j=0}^l \frac{p^{-j}}{\phi(p^{l-j})} |W(p^{l-j}, ap^{j(k-1)})|. \quad (3.7)$$

We begin by estimating the contribution to $\mathcal{S}(p^l, a)$ arising from the summands in (3.7) with $j(k-1) \geq l-j$. In this situation, one sees from (3.2) that $W(p^{l-j}, ap^{j(k-1)}) = \phi(p^{l-j})$. Adopting the formulation employed already in the definition of the function κ , we write $l = uk + v$ with $u \geq 0$ and $1 \leq v \leq k$. Then, the integers j with $j(k-1) \geq l-j$ are precisely those characterised by the condition $j > u$. The contribution from these summands on the right hand side of (3.7) is exactly

$$\sum_{j=u+1}^l p^{-j} \leq p^{-u-1} \frac{p}{p-1}.$$

On recalling the relation (3.4), we therefore obtain the upper bound

$$\mathcal{S}(p^l, a) \leq T + p^{-u-1} \psi(p^l), \quad (3.8)$$

where

$$T = \sum_{j=0}^u \frac{p^{-j}}{\phi(p^{l-jk})} |W(p^{l-jk}, a)|.$$

We temporarily suppose that $p \nmid 6k$ and proceed to show that in this case one has

$$\mathcal{S}(p^l, a) \leq \kappa(p^l) \psi(p^l). \quad (3.9)$$

Indeed, if $2 \leq v \leq k$, then for all $0 \leq j \leq u$ we have $l - jk \geq 2$, and we may apply the first clause of Lemma 3.2 with $\theta = 0$ to confirm that $T = 0$. In the current context, we have $\kappa(p^l) = p^{-u-1}$, so that (3.9) now follows from (3.8).

This leaves the case $v = 1$. By using the same reasoning we now find that the only term that contributes to the sum T is that in which $j = u$, whence

$$T = \frac{|W(p, a)|}{p^u(p-1)}.$$

By (3.3) and (3.8), it therefore follows that

$$\mathcal{S}(p^l, a) \leq p^{-u-1}\psi(p^l)(2 + (k-1)\sqrt{p}).$$

In the current discussion, moreover, we have $p \geq 5$, and thus

$$2 + (k-1)\sqrt{p} \leq k\sqrt{p} = p^{u+1}\kappa(p^l).$$

Consequently, the upper bound (3.9) follows in the case $v = 1$.

Next suppose that $p \mid 6k$. In this case we prove that

$$\mathcal{S}(p^l, a) \leq p\kappa(p^l)\psi(p^l). \quad (3.10)$$

We begin with the case where $p \geq 3$. When $p \mid k$, we find that Lemma 3.2 applies with $\theta \geq 1$ and shows that whenever $W(p^{l-jk}, a)$ is non-zero, then $l - jk \leq \theta + 1$. This implies that $(u - j)k \leq \theta + 1 - v \leq \theta$. However, the upper bound $p^\theta \leq k$ implies that $\theta < k$. Hence, it is again the case that the only summand that contributes to T is that with $j = u$, and then a trivial bound for $|W(p^v, a)|$ leads to the relation

$$T = p^{-u} \frac{|W(p^v, a)|}{\phi(p^v)} \leq p^{-u} = (p-1)p^{-u-1}\psi(p^l).$$

We therefore deduce from (3.8) and the definition of $\kappa(p^l)$ that

$$\mathcal{S}(p^l, a) \leq p^{-u}\psi(p^l) \leq p\kappa(p^l)\psi(p^l). \quad (3.11)$$

If $p = 3$ but $3 \nmid k$, meanwhile, then $\theta = 0$ and the above argument still applies with some obvious modification to deliver (3.11). Thus, the bound (3.10) holds whenever $p \geq 3$.

This leaves the prime $p = 2$ for consideration. We first consider the situation in which $k \geq 3$. When k is odd, we have $\theta = 0$. When k is even and $k \geq 4$, meanwhile, we have $\theta \geq 1$ and $2^\theta \leq k$. It follows that in both scenarios we have $\theta \leq k - 2$. Here, when $0 \leq j \leq u$ and $W(2^{l-jk}, a)$ is non-zero, it follows from Lemma 3.2 that $l - jk \leq \theta + 2 \leq k$. This implies that $(u - j)k \leq k - v \leq k - 1$. Thus, we again see that the only summand that contributes to T is that with $j = u$, and the proof of (3.10) runs as before via (3.11).

The final case left is that where $p = k = 2$. We now have $\theta = 1$, and Lemma 3.2 shows that when $1 \leq j \leq u$, one has $W(2^{l-2j}, a) = 0$ unless $l - 2j \leq 3$. Thus, we may restrict the range of summation in T to those values of j with $2(u - j) \leq 3 - v$. In case we have $v = 2$ or $u = 0$, we conclude that $j = u$ and may then one last time appeal to the foregoing argument to verify (3.10). We are then reduced to the situation where $v = 1$ and $u \geq 1$. This case is different, for now the summands with $j = u$ and $j = u - 1$ make non-zero contributions to T , and we find that

$$T = 2^{-u} \frac{|W(2, a)|}{\phi(2)} + 2^{1-u} \frac{|W(8, a)|}{\phi(8)} \leq 3 \cdot 2^{-u} = 3 \cdot 2^{-u-1}\psi(2^l).$$

By (3.8), one now finds that $\mathcal{S}(2^l, a) \leq 2^{1-u}\psi(2^l)$. Meanwhile, since l is odd, one has $\kappa(2^l) = 2^{-u+1/2} > 2^{-u}$, and (3.10) follows.

With (3.9) and (3.10) in hand, the proof of the lemma is swiftly completed. The convolution (3.6) passes the quasi-multiplicative property (3.5) on to $\mathcal{S}(q, a)$, delivering the formula

$$\mathcal{S}(q_1q_2, a) = \mathcal{S}(q_1, aq_2^{k-1})\mathcal{S}(q_2, aq_1^{k-1})$$

that is valid for all coprime natural numbers q_1, q_2 and all integers a . When $(q, a) = 1$, one may repeatedly apply this identity to reduce to the exact prime powers dividing q . Then, by (3.9) and (3.10), one finds that

$$\mathcal{S}(q, a) \leq \kappa(q)\psi(q) \prod_{p|6k} p.$$

The conclusion of the lemma now follows with a trivial estimate for the final product. \square

The proof of Theorem 1.2. We are now fully equipped to embark on the proof of Theorem 1.2. Suppose then that the hypotheses of this theorem are satisfied. We begin with the special case where

$$2 \leq R \leq \exp((\log P)^{1/3}).$$

Here, a straightforward estimation is enough. In fact, it follows easily from [8, Chapter III.5, Theorem 1] that, uniformly for $2 \leq R \leq P$, one has

$$\text{card}(\mathcal{A}(P, R)) \ll P \exp\left(-\frac{\log P}{2 \log R}\right).$$

In the range for R that is currently under consideration, it therefore suffices to remark that by means of a trivial estimate one has

$$|g(\alpha)| + |g_\nu(\alpha)| \ll \text{card}(\mathcal{A}(P, R)) \ll P \exp\left(-\frac{1}{2}(\log P)^{2/3}\right).$$

This upper bound is stronger than those claimed in Theorem 1.2.

From now onwards, we shall suppose that

$$\exp((\log P)^{1/3}) \leq R \leq P^{2/3}, \tag{3.12}$$

and proceed to establish the estimate for $g_\nu(\alpha)$ contained in Theorem 1.2. A considerable part of our argument is largely identical with the demonstration of [13, Lemma 8.5]. We therefore opt for a guided tour through the necessary changes and refinements. The reader is expected to be familiar with the exposition in [13], and we take the liberty to apply the notation of the latter source without reintroducing it here.

In [13] it is assumed that P and R are linked via the equation $R = P^\eta$, for some fixed $\eta \in (0, 1/2)$. Here, we read this relation as a definition for η and check the argument given in [13, pp. 56–58] for uniformity as R ranges over the interval (3.12). We begin by noting that

$$g_\nu(\alpha; P, R) = g(\alpha; P, R) - g(\alpha; P/\nu, R) \tag{3.13}$$

and apply the initial decomposition of $g(\alpha)$ leading up to [13, equation (8.3)]. We recast the latter equation in the form

$$g(\alpha; P, R) = \sum_{\substack{d|q \\ q/d \in \mathcal{A}(R)}} W(d, a(q/d)^{k-1}) \frac{\mathcal{M}_d(Q)}{\phi(d)} + O(P(1 + P^k|\beta|)e^{-c\sqrt{\log P}}).$$

Here $Q = Pd/q$, we write $\beta = \alpha - a/q$, and c is a certain positive constant. Moreover, we use $\mathcal{M}_d(Q)$ to denote the expression \mathcal{M}_d that is defined in the display following [13, equation (8.3)], and we write $\mathcal{A}(R)$ for the set of natural numbers free of prime factors exceeding R . Note that in [13, equation (8.3)] the sum over $d|q$ is constrained to satisfy $q/d \in \mathcal{A}(P, R)$, but since P is large and $q \leq P^\varepsilon$, this makes no difference. By (3.13), we now find that

$$g_\nu(\alpha; P, R) = \sum_{\substack{d|q \\ q/d \in \mathcal{A}(R)}} W(d, a(q/d)^{k-1}) \frac{\mathcal{M}_d(Q) - \mathcal{M}_d(Q/\nu)}{\phi(d)} + O(P(1 + P^k|\beta|)e^{-c\sqrt{\log P}}). \quad (3.14)$$

Next, for $1 \leq j \leq 3$, we denote by $\mathcal{N}_j(Q)$ the expressions \mathcal{N}_j that are defined in the displays following [13, equation (8.4)]. On applying the latter reference, we then have

$$\mathcal{M}_d(Q) - \mathcal{M}_d(Q/\nu) = \sum_{j=1}^3 (-1)^{j+1} (\mathcal{N}_j(Q) - \mathcal{N}_j(Q/\nu)). \quad (3.15)$$

As in [13, p. 57], we have

$$\begin{aligned} \sum_{j=1}^2 (-1)^{j+1} (\mathcal{N}_j(Q) - \mathcal{N}_j(Q/\nu)) &= \frac{\phi(d)}{d} \int_{Q/\nu}^Q e(\gamma X^k) dX + O(d^\varepsilon(1 + Q^k|\gamma|)) \\ &\ll Q(1 + Q^k|\gamma|)^{-1} + d^\varepsilon(1 + Q^k|\gamma|). \end{aligned} \quad (3.16)$$

Here, we applied Lemma 3.1 to estimate the exponential integral.

We now consider the expression $\mathcal{N}_3(Q) - \mathcal{N}_3(Q/\nu)$. Again, we follow the pattern of [13] and invoke the integral representation

$$\mathcal{N}_3(Q) = \int_0^{\log(Q/R)/\log R} N_d(R^v) (\mathcal{N}_4(Q; v) - e(\gamma R^{k(v+1)})R^{v+1}) dv$$

that occurs on [13, p. 58], and the function

$$\mathcal{N}_4(Q; v) = \int_{R^{v+1}}^Q e(\gamma X^k) \left(\rho_1 \left(\frac{\log X}{\log R} - v \right) + (\log R)^{-1} \rho_2 \left(\frac{\log X}{\log R} - v \right) \right) dX \quad (3.17)$$

is featured here. It also occurs (under the name $\mathcal{N}_4(v)$) on [13, p. 58]. Now

$$\mathcal{N}_3(Q) - \mathcal{N}_3(Q/\nu) = -\mathcal{N}_3^* + \mathcal{N}_3^\dagger + \mathcal{N}_3^\ddagger, \quad (3.18)$$

where

$$\begin{aligned} \mathcal{N}_3^* &= \int_{\log(Q/(\nu R))/\log R}^{\log(Q/R)/\log R} N_d(R^v) e(\gamma R^{k(v+1)}) R^{v+1} dv, \\ \mathcal{N}_3^\dagger &= \int_0^{\log(Q/(\nu R))/\log R} N_d(R^v) (\mathcal{N}_4(Q; v) - \mathcal{N}_4(Q/\nu; v)) dv, \\ \mathcal{N}_3^\ddagger &= \int_{\log(Q/(\nu R))/\log R}^{\log(Q/R)/\log R} N_d(R^v) \mathcal{N}_4(Q; v) dv. \end{aligned}$$

The quantity \mathcal{N}_3^* corresponds to the expression \mathcal{N}_5 in [13]. We substitute $w = R^{v+1}$ and import the asymptotic relation $N_d(X) = \phi(d)/d + O(d^\varepsilon X^{-1})$ from [13, p. 57, *inter alia*] to confirm that

$$\begin{aligned}\mathcal{N}_3^* &= (\log R)^{-1} \int_{Q/\nu}^Q N_d(w/R) e(\gamma w^k) dw \\ &= \frac{\phi(d)}{d \log R} \int_{Q/\nu}^Q e(\gamma w^k) dw + O(d^\varepsilon R).\end{aligned}$$

Now Lemma 3.1 supplies the bound

$$\mathcal{N}_3^* \ll Q(1 + Q^k |\gamma|)^{-1} + d^\varepsilon R. \quad (3.19)$$

Thus far, we have followed the arguments from [13] closely, replacing [13, Lemma 8.2] by Lemma 3.1 when appropriate. The treatment of \mathcal{N}_3^\dagger requires more care. By (3.17), we find that

$$\mathcal{N}_4(Q; v) - \mathcal{N}_4(Q/\nu; v) = \int_{\Theta(v)}^Q e(\gamma X^k) \left(\rho_1 \left(\frac{\log X}{\log R} - v \right) + \frac{1}{\log R} \rho_2 \left(\frac{\log X}{\log R} - v \right) \right) dX, \quad (3.20)$$

where $\Theta(v) = \max\{R^{v+1}, Q/\nu\}$. Note that Lemma 3.1 immediately supplies an upper bound of order $Q(1 + Q^k |\gamma|)^{-1}$ for the expression on the right hand side of (3.20). When R is a fixed power of P , then the range of integration in the definition of \mathcal{N}^\dagger is an interval of length $O(1)$, and one arrives at the acceptable bound $\mathcal{N}^\dagger \ll Q(1 + Q^k |\gamma|)^{-1}$. This is the argument in [13]. However, with R now ranging over the interval described by (3.12), this line of attack is no longer of strength sufficient to press the method home. We therefore apply Lemma 3.1 with more precision. The functions $-\rho_1$ and ρ_2 (defined in [13]) are positive and decreasing, with continuous derivatives, on the range of integration, and thus we may apply Lemma 3.1. For $j = 1$ and 2 , this gives

$$\int_{\Theta(v)}^Q e(\gamma X^k) \rho_j \left(\frac{\log X}{\log R} - v \right) dX \ll Q(1 + Q^k |\gamma|)^{-1} \left| \rho_j \left(\frac{\log \Theta(v)}{\log R} - v \right) \right|.$$

From the definition of $\mathcal{N}_4(Q; v)$ and properties of ρ_1 and ρ_2 , we see that $\mathcal{N}_4(Q; v)$ vanishes for $v > \log(Q/R)/\log R$. Since $N_d(X)$ is bounded, we deduce that

$$\mathcal{N}_3^\dagger \leq \int_0^\infty N_d(R^v) |\mathcal{N}_4(Q; v) - \mathcal{N}_4(Q/\nu; v)| dv \leq \mathcal{I} Q(1 + Q^k |\gamma|)^{-1}, \quad (3.21)$$

where

$$\mathcal{I} = \int_0^{\log(Q/R)/\log R} \left(\left| \rho_1 \left(\frac{\log \Theta(v)}{\log R} - v \right) \right| + (\log R)^{-1} \rho_2 \left(\frac{\log \Theta(v)}{\log R} - v \right) \right) dv.$$

We have $\Theta(v) = R^{v+1}$ whenever $v + 1 > \log(Q/\nu)/\log R$, and otherwise $\Theta(v) = Q/\nu$. Now, the range of integration in the integral \mathcal{I} is $0 \leq v \leq \log(Q/R)/\log R$. Thus, the part where $\Theta(v) = R^{v+1}$ holds is an interval of length $\log \nu / \log R$, and in the integrand we then have $-\rho_1(1+) = 1$ and $\rho_2(1+) = 1$. This part therefore contributes a bounded amount. The contribution of the remaining range is bounded by

$$\int_0^\infty (|\rho_1(t)| + |\rho_2(t)|) dt,$$

and this integral exists owing to the exponential decay of Dickman's function that is a majorant for both ρ_1 and ρ_2 . We conclude that $\mathcal{I} \ll 1$. By (3.21), it follows that

$$\mathcal{N}_3^\dagger \ll Q(1 + Q^k|\gamma|)^{-1}. \quad (3.22)$$

The treatment of \mathcal{N}_3^\ddagger is simpler. The interval of integration in the definition of \mathcal{N}_3^\ddagger is of length $\log \nu / \log R$, and therefore bounded. For each ν in this interval we have $\nu > (\log(Q/(R\nu)))/\log R$, whence $R^{\nu+1} \geq Q/\nu$. We may now apply Lemma 3.1 to see that $\mathcal{N}_4(Q; \nu) \ll Q(1 + Q^k|\gamma|)^{-1}$ holds uniformly for all ν in the range of integration. This gives

$$\mathcal{N}_3^\ddagger \ll Q(1 + Q^k|\gamma|)^{-1}. \quad (3.23)$$

On collecting together the estimates (3.19), (3.22) and (3.23), we conclude from (3.18), (3.16) and (3.15) that

$$\mathcal{M}_d(Q) - \mathcal{M}_d(Q/\nu) \ll Q(1 + Q^k|\gamma|)^{-1} + d^\varepsilon R + d^\varepsilon(1 + Q^k|\gamma|).$$

As in [13], we apply (3.14) and find that

$$g_\nu(\alpha; P, R) = P\mathcal{S}(q, a)(1 + P^k|\beta|)^{-1} + P(1 + P^k|\beta|)e^{-c\sqrt{\log P}}.$$

The proof of the first conclusion of Theorem 1.2 is now completed by reference to Lemma 3.3, and indeed we have established this part of the theorem for $2 \leq R \leq P^{2/3}$.

We now prove the second conclusion of Theorem 1.2 by making use of a dyadic dissection. Let J be the smallest natural number having the property that $2^J > P^{1/9}$. Then

$$g(\alpha; P, R) = \sum_{j=0}^J g_2(\alpha; 2^{-j}P, R) + O(P^{8/9}). \quad (3.24)$$

Fix a number $A > 1$ and apply the first conclusion of Theorem 1.2 that is already established, but with $2A$ in place of A , and with $2^{-j}P$ in place of P , for $0 \leq j \leq J$. Notice that we have established this consequence of Theorem 1.2 in the range $2 \leq R \leq P^{2/3}$ that is wider than the corresponding range $2 \leq R \leq P^{1/2}$ claimed in the statement of the theorem. Since one has $(\log X)^{2A} > (\log P)^A$ and $X^{2/3} > P^{1/2}$ for $P^{8/9} \leq X \leq P$, the desired bound for $g(\alpha; P, R)$ follows via the dyadic decomposition (3.24) and an application of Lemma 2.3. \square

4. AN AUXILIARY PRUNING LEMMA

We now discuss a pruning lemma that should be of wider utility than the immediate applications of this paper demand. For this reason, we keep the presentation in this section fairly self-contained. As usual, we fix a natural number k , which in this section is assumed to satisfy $k \geq 3$, and define the function $\kappa(q)$ as in the preamble to Theorem 1.1. Now choose parameters δ , Q , X , and Y with

$$\delta > 1, \quad Q \geq 1, \quad X \geq 1, \quad Y > 0, \quad QY \leq \frac{1}{4}X^k. \quad (4.1)$$

When $0 \leq a \leq q \leq Q$ and $(a, q) = 1$, the intervals defined by

$$\mathcal{M}(q, a) = \{\alpha \in [0, 1) : |q\alpha - a| \leq YX^{-k}\}$$

are disjoint. We denote the union of these intervals by $\mathcal{M} = \mathcal{M}_{Q,X,Y}$ and define the function $\Upsilon : \mathcal{M} \rightarrow [0, 1]$ by putting

$$\Upsilon(\alpha) = \kappa(q)^2 (1 + X^k |\alpha - a/q|)^{-\delta}$$

when $\alpha \in \mathcal{M}(q, a)$.

Lemma 4.1. *Let k and t be natural numbers with $k \geq 3$ and $t \geq \lfloor k/2 \rfloor$. Suppose that the numbers δ , Q , X and Y satisfy (4.1). Then, for any subset \mathcal{Z} of $[1, X] \cap \mathbb{Z}$, one has for each $\varepsilon > 0$ the estimate*

$$\int_{\mathcal{M}} \Upsilon(\alpha) \left| \sum_{x \in \mathcal{Z}} e(\alpha x^k) \right|^{2t} d\alpha \ll Q^\varepsilon X^{2t-k} + X^{-k} Q^{t+1+\varepsilon}.$$

Proof. Our treatment is a development of the argument of the proof of [15, Lemma 5.4]. We have

$$\int_{\mathcal{M}} \Upsilon(\alpha) \left| \sum_{x \in \mathcal{Z}} e(\alpha x^k) \right|^{2t} d\alpha \leq \sum_{1 \leq q \leq Q} \kappa(q)^2 \int_{-Y/X^k}^{Y/X^k} \frac{W(\beta, q, \mathcal{Z})}{(1 + X^k |\beta|)^\delta} d\beta, \quad (4.2)$$

in which we write

$$W(\beta, q, \mathcal{Z}) = \sum_{a=1}^q \left| \sum_{x \in \mathcal{Z}} e(x^k(\beta + a/q)) \right|^{2t}.$$

Write

$$\psi(\mathbf{x}) = \sum_{i=1}^t (x_{2i-1}^k - x_{2i}^k).$$

Then we find by orthogonality that

$$\begin{aligned} W(\beta, q, \mathcal{Z}) &= q \sum_{\substack{\mathbf{x} \in \mathcal{Z}^{2t} \\ q \mid \psi(\mathbf{x})}} e(\beta \psi(\mathbf{x})) \\ &\leq q \sum_{\substack{1 \leq x_1, \dots, x_{2t} \leq X \\ q \mid \psi(\mathbf{x})}} 1 \\ &= \sum_{a=1}^q \left| \sum_{1 \leq x \leq X} e(x^k a/q) \right|^{2t}. \end{aligned} \quad (4.3)$$

We now enhance the corresponding argument of [15, Lemma 5.4] by applying [12, Theorem 4.1] to see that when $(a, q) = 1$, one has

$$\sum_{1 \leq x \leq X} e(x^k a/q) = Xq^{-1} S(q, a) + O(q^{1/2+\varepsilon}),$$

where

$$S(q, a) = \sum_{r=1}^q e(ar^k/q).$$

By [12, Lemmata 4.3, 4.4 and 4.5], we therefore deduce in such circumstances that

$$\sum_{1 \leq x \leq X} e(x^k a/q) \ll X\kappa(q) + q^{1/2+\varepsilon}.$$

Removing now the condition $(a, q) = 1$, we find that in general one has the upper bound

$$\sum_{1 \leq x \leq X} e(x^k a/q) \ll X \kappa(q/(q, a)) + (q/(q, a))^{1/2+\varepsilon}. \quad (4.4)$$

Next, we substitute (4.4) into (4.3) to obtain the estimate

$$\begin{aligned} W(\beta, q, \mathcal{Z}) &\ll X^{2t} \sum_{a=1}^q \kappa(q/(q, a))^{2t} + \sum_{a=1}^q (q/(q, a))^{t+\varepsilon} \\ &\ll X^{2t} \sigma(q) + \sum_{r|q} r^{t+1+\varepsilon}, \end{aligned} \quad (4.5)$$

where

$$\sigma(q) = \sum_{r|q} r \kappa(r)^{2t}.$$

Then we find that $\sigma(q)$ is the subject of the discussion of [15] leading from equation (5.9) to the conclusion of the proof of Lemma 5.4 of the latter source. Unfortunately, the estimations there are subject to the condition $k \geq 4$, but subject to this constraint it is shown that

$$\sum_{1 \leq q \leq Q} \kappa(q)^2 \sigma(q) \ll Q^\varepsilon. \quad (4.6)$$

This bound remains valid for $k = 3$, too, as we shall demonstrate momentarily. Since one also has the bound

$$\sum_{r|q} r^{t+1+\varepsilon} \ll q^{t+1+2\varepsilon},$$

we see from the definition of $\kappa(q)$ that there is a positive number B having the property that

$$\sum_{1 \leq q \leq Q} \kappa(q)^2 \sum_{r|q} r^{t+1+\varepsilon} \ll Q^{t+1+2\varepsilon} \prod_{p \leq Q} (1 + Bp^{-1}) \ll Q^{t+1+3\varepsilon}.$$

Thus, we deduce from (4.2) and (4.5) that

$$\int_{\mathcal{M}} \Upsilon(\alpha) \left| \sum_{x \in \mathcal{Z}} e(\alpha x^k) \right|^{2t} d\alpha \ll X^{2t-k} \sum_{1 \leq q \leq Q} \kappa(q)^2 \sigma(q) + X^{-k} \sum_{1 \leq q \leq Q} \kappa(q)^2 \sum_{r|q} r^{t+1+\varepsilon},$$

and the conclusion of the lemma is now immediate.

It remains to confirm (4.6) when $k = 3$. In view of the crude estimate $\kappa_3(q) \ll q^{-1/3}$ it suffices to treat the case $t = 1$. In this case, the crude bound shows that $\sigma(p^l) \ll lp^{l/3}$, whence $\kappa(p^l)^2 \sigma(p^l) \ll lp^{-l/3}$. We therefore deduce that

$$\sum_{l=4}^{\infty} \kappa(p^l)^2 \sigma(p^l) \ll p^{-4/3}.$$

Meanwhile, working directly from the definition of $\kappa(q)$, we find that there is a constant $C > 1$ with the property that

$$\kappa(p)^2 \sigma(p) + \kappa(p^2)^2 \sigma(p^2) + \kappa(p^3)^2 \sigma(p^3) \leq Cp^{-1}.$$

The bound (4.6) now follows from the estimate

$$\sum_{1 \leq q \leq Q} \kappa(q)^2 \sigma(q) \leq \prod_{p \leq Q} \sum_{l=0}^{\infty} \kappa(p^l)^2 \sigma(p^l) \leq \prod_{p \leq Q} (1 + Cp^{-1} + O(p^{-4/3})).$$

□

5. MEAN VALUES ON SETS OF MAJOR ARCS

Our goal in this section is the proof of Theorem 1.3 concerning upper bounds for moments of the smooth Weyl sum $g(\alpha; P, R)$ on sets of major arcs.

Lemma 5.1. *Let k and t be natural numbers with $k \geq 3$ and $t \geq \lfloor k/2 \rfloor$, and let ν be a real number with $\nu > 1$. Suppose that R and P are real numbers with $2 \leq R \leq P^\eta$, where η is a positive number sufficiently small in terms of k and ε . Finally, let $\omega < k/2$ be a positive number with*

$$\omega < \frac{2t+4}{t+10}, \quad (5.1)$$

and suppose that $1 \leq Q \leq P^\omega$. Then, there exists a positive number τ with the property that, whenever $1 \leq X \leq P$, one has

$$\int_{\mathfrak{M}(Q)} |g_\nu(\alpha; X, R)|^{2t+4} d\alpha \ll Q^\varepsilon X^{2t+4-k} (\log X)^{4+\varepsilon} + P^{2t+4-k-\tau}.$$

Proof. We begin by observing that the conclusion of the lemma is trivial when $X \leq P^{1/2}$, so we suppose henceforth that $P^{1/2} < X \leq P$. Consider a real number α with $\alpha \in \mathfrak{M}(Q)$. Then there exists $a \in \mathbb{Z}$ and $q \in \mathbb{N}$ with $1 \leq q \leq Q$ and $(a, q) = 1$ with $|q\alpha - a| \leq QP^{-k}$. Consequently, it follows from Theorem 1.1 that

$$g_\nu(\alpha; X, R) \ll \frac{q^\varepsilon \kappa(q)^{1/2} X (\log X)^{1+\varepsilon}}{(1 + X^k |\alpha - a/q|)^{1/2-\varepsilon}} + X^{3/4} R^{1/2} (\log X)^{1/4+\varepsilon} Q^{1/8+\varepsilon}.$$

Next, we prepare for an application of Lemma 4.1 where we work with X and Q as above, and we choose $Y = Q(X/P)^k$ to arrange that $\mathfrak{M}(Q) = \mathcal{M}_{Q, X, Y}$. We define the function $\Upsilon(\alpha)$ on $\mathcal{M}_{Q, X, Y}$ as in the preamble of Lemma 4.1, with $\delta = 3/2$, and then recast the preceding bound for $g_\nu(\alpha; X, R)$ in the form

$$g_\nu(\alpha; X, R) \ll Q^\varepsilon X (\log X)^{1+\varepsilon} \Upsilon(\alpha)^{1/4} + X^{3/4} R^{1/2} (\log X)^{1/4+\varepsilon} Q^{1/8+\varepsilon}.$$

Thus, we deduce that the mean value

$$I_0 = \int_{\mathfrak{M}(Q)} |g_\nu(\alpha; X, R)|^{2t+4} d\alpha \quad (5.2)$$

satisfies the bound

$$I_0 \ll I_1 + I_2, \quad (5.3)$$

where

$$I_1 = Q^\varepsilon (X (\log X)^{1+\varepsilon})^4 \int_{\mathfrak{M}(Q)} \Upsilon(\alpha) |g_\nu(\alpha; X, R)|^{2t} d\alpha \quad (5.4)$$

and

$$I_2 = Q^{1/2+\varepsilon} X^{3+\varepsilon} R^2 \int_{\mathfrak{M}(Q)} |g_\nu(\alpha; X, R)|^{2t} d\alpha. \quad (5.5)$$

In order to bound the integral occurring in (5.5), we recall (5.2) and apply Hölder's inequality to obtain the estimate

$$\begin{aligned} \int_{\mathfrak{M}(Q)} |g_\nu(\alpha; X, R)|^{2t} d\alpha &\ll \left(\int_{\mathfrak{M}(Q)} d\alpha \right)^{2/(t+2)} \left(\int_{\mathfrak{M}(Q)} |g_\nu(\alpha; X, R)|^{2t+4} d\alpha \right)^{t/(t+2)} \\ &\ll (Q^2 P^{-k})^{2/(t+2)} I_0^{t/(t+2)}. \end{aligned}$$

Applying Lemma 4.1 to estimate the mean value I_1 defined in (5.4), we therefore deduce from (5.3) that

$$I_0 \ll Q^\varepsilon X^4 (\log X)^{4+\varepsilon} (X^{2t-k} + X^{-k} Q^{t+1}) + Q^{1/2+\varepsilon} X^{3+\varepsilon} R^2 (Q^2 P^{-k})^{2/(t+2)} I_0^{t/(t+2)},$$

whence

$$I_0 \ll Q^\varepsilon X^4 (\log X)^{4+\varepsilon} (X^{2t-k} + X^{-k} Q^{t+1}) + Q^2 P^{-k} (Q^{1/8+\varepsilon} X^{3/4+\varepsilon} R^{1/2})^{2t+4}. \quad (5.6)$$

Since we have $Q \leq P^\omega$ with ω satisfying the bound (5.1), we find that there is a positive number τ for which

$$Q^2 P^{-k} (Q^{1/8} X^{3/4})^{2t+4} = Q^{(t+10)/4} X^{(3t+6)/2} P^{-k} \ll P^{(t+2)/2-2\tau} X^{(3t+6)/2} P^{-k}.$$

But $X \leq P$, and thus

$$Q^2 P^{-k} (Q^{1/8+\varepsilon} X^{3/4+\varepsilon} R^{1/2})^{2t+4} \ll P^{2t+4-k-\tau}. \quad (5.7)$$

The upper bound (5.7) provides satisfactory control over the final term on the right hand side of (5.6). We turn next to consider the second half of the first term on the right-hand side of this relation. Our goal is to show that $X^{4-k} Q^{t+1} < P^{2t+4-k-2\tau}$. When $k = 3$ this bound is easily obtained by employing the upper bounds $X \leq P$ and $Q \leq P^\omega$. One has

$$\frac{(t+1)(2t+4)}{t+10} = 2t - \frac{14t-4}{t+10} < 2t - 2\tau,$$

so that in view of (5.1), one has

$$XQ^{t+1} < P^{2t+1-2\tau},$$

and the desired conclusion follows. We may henceforth suppose, therefore, that $k \geq 4$. Here, we begin by observing that a trivial estimate for $g_\nu(\alpha; X, R)$ shows that when

$$X^{2t+4} Q^2 P^{-k} < P^{2t+4-k-\tau},$$

then

$$I_0 = \int_{\mathfrak{M}(Q)} |g_\nu(\alpha; X, R)|^{2t+4} d\alpha \ll X^{2t+4} \text{mes}(\mathfrak{M}(Q)) \ll P^{2t+4-k-\tau}.$$

We may therefore suppose that $X^{2t+4} Q^2 \geq P^{2t+4-\tau}$, whence

$$X \geq P^{1-2/(t+10)} Q^{-1/(t+2)}.$$

But then, on recalling (5.1) and the hypothesis $Q \leq P^\omega$, we find that

$$X^{4-k} Q^{t+1} \leq P^{4-k+2(k-4)/(t+10)} Q^{t+1+(k-4)/(t+2)} \leq P^{4-k+2\Lambda/(t+10)},$$

where $\Lambda = (t+1)(t+2) + 2(k-4)$. However, since $k \leq 2t+1$, one finds that

$$\Lambda = t(t+10) - (7t-2k+6) \leq t(t+10) - (3t+4) < (t+10)(t-\tau).$$

We therefore conclude that when $k \geq 4$, we may assume that $X^{4-k}Q^{t+1} < P^{2t+4-k-2\tau}$. Then in all cases, we find that the bound (5.6) implies via (5.7) that

$$I_0 \ll Q^\varepsilon X^{2t+4-k} (\log X)^{4+\varepsilon} + P^{2t+4-k-\tau},$$

and thus the proof of the lemma is complete. \square

Before announcing a corollary of this conclusion, we recall the definition (1.6) of $\mathfrak{N}(Q)$.

Corollary 5.2. *Assume the hypotheses of the statement of Lemma 5.1, and let τ be the positive number supplied by its conclusion. Also, let ω' be a positive number with*

$$\omega' < \frac{2k}{k+4}, \quad (5.8)$$

and put $\Omega = \min\{\omega, \omega'\}$. Then provided that $1 \leq Q \leq P^\omega$, one has

$$\int_{\mathfrak{N}(Q)} |g(\alpha; P, R)|^{2t+4} d\alpha \ll Q^\varepsilon P^{2t+4-k} (\log P)^{2t+8+\varepsilon}.$$

Moreover, when $u > 2t+4$ and A is sufficiently large in terms of τ and u , then whenever $(\log P)^A \leq Q \leq P^\Omega$, one has

$$\int_{\mathfrak{N}(Q)} |g(\alpha; P, R)|^u d\alpha \ll P^{u-k} Q^{\varepsilon - (u-2t-4)/(2k)}.$$

Proof. Write $\zeta = 2/(t+10)$. Then by dividing the summation underlying $g(\alpha; P, R)$ into dyadic intervals, we find that

$$|g(\alpha; P, R)| \leq \sum_{\substack{j=0 \\ 2^j \leq P^\zeta}}^{\infty} |g_2(\alpha; 2^{-j}P, R)| + O(P^{1-\zeta}).$$

By Hölder's inequality, therefore, we have

$$|g(\alpha; P, R)|^{2t+4} \ll (\log P)^{2t+3} \sum_{\substack{j=0 \\ 2^j \leq P^\zeta}}^{\infty} |g_2(\alpha; 2^{-j}P, R)|^{2t+4} + P^{(2t+4)(1-\zeta)}.$$

Put

$$I = \int_{\mathfrak{N}(Q)} |g(\alpha; P, R)|^{2t+4} d\alpha.$$

Then it follows from Lemma 5.1 that there is a real number X satisfying $P^{1-\zeta} \leq X \leq P$ for which

$$\begin{aligned} I &\ll (\log P)^{2t+4} \int_{\mathfrak{N}(Q)} |g_2(\alpha; X, R)|^{2t+4} d\alpha + P^{(2t+4)(1-\zeta)} Q^2 P^{-k} \\ &\ll Q^\varepsilon P^{2t+4-k} (\log P)^{2t+8+\varepsilon} + P^{2t+4-k-\tau/2} + Q^2 P^{(2t+4)(1-\zeta)-k}. \end{aligned}$$

However, our hypothesis concerning Q ensures that $Q^2 < P^{(2t+4)\zeta}$, and thus we conclude that

$$\int_{\mathfrak{N}(Q)} |g(\alpha; P, R)|^{2t+4} d\alpha \ll Q^\varepsilon P^{2t+4-k} (\log P)^{2t+8+\varepsilon} + P^{2t+4-k}.$$

This delivers the first mean value estimate of the corollary.

For the second bound, we consider a typical point $\alpha \in \mathfrak{N}(Q)$. There exists $a \in \mathbb{Z}$ and $q \in \mathbb{N}$ with $(a, q) = 1$ and

$$Q/2 \leq q + P^k |q\alpha - a| \leq 2Q.$$

Then we see from Theorem 1.1 that

$$g(\alpha; P, R) \ll P(\log P)^{1+\varepsilon} Q^{\varepsilon-1/(2k)} + P^{1+\varepsilon} (Q/P^2)^{1/8}.$$

However, we may suppose that $Q \leq P^{\omega'}$, and so it follows from (5.8) that there is a positive number ξ having the property that $Q \leq P^{2k/(k+4)-16\xi}$. Thus, we have

$$(Q/P^2)^{1/8} \leq P^{-1/(k+4)-2\xi} \leq P^{-2\xi} Q^{-1/(2k)}.$$

Since $Q \geq (\log P)^A$ with A sufficiently large, it follows that

$$g(\alpha; P, R) \ll PQ^{2\varepsilon-1/(2k)}.$$

By making use of this bound in combination with the first conclusion of the corollary, we arrive at the upper bound

$$\begin{aligned} \int_{\mathfrak{N}(Q)} |g(\alpha; P, R)|^u d\alpha &\ll (PQ^{\varepsilon-1/(2k)})^{u-2t-4} \int_{\mathfrak{M}(Q)} |g(\alpha; P, R)|^{2t+4} d\alpha \\ &\ll (PQ^{\varepsilon-1/(2k)})^{u-2t-4} Q^\varepsilon P^{2t+4-k} (\log P)^{2t+8+\varepsilon}. \end{aligned}$$

Again making use of the assumption $Q \geq (\log P)^A$ with A sufficiently large, we see that $(\log P)^{2t+9} \ll Q^\varepsilon$, and hence

$$\int_{\mathfrak{N}(Q)} |g(\alpha; P, R)|^u d\alpha \ll P^{u-k} Q^{u\varepsilon-(u-2t-4)/(2k)}.$$

This completes the proof of the corollary. \square

Corollary 5.2 is of strength sufficient to deliver the conclusions of Theorem 1.3 in all cases with $Q \geq (\log P)^A$, for a suitably large positive number A . We now attend to the remaining values of Q with $1 \leq Q < (\log P)^A$.

Lemma 5.3. *Let k be a natural number with $k \geq 4$, and let A be a positive real number. Then, whenever Q , R and P are real numbers with $2 \leq R \leq P^{1/2}$ and $1 \leq Q \leq (\log P)^A$, one has*

$$\int_{\mathfrak{M}(Q)} |g(\alpha; P, R)|^{k+1} d\alpha \ll P \log Q. \quad (5.9)$$

Moreover, when $u > k + 1$, one also has

$$\int_{\mathfrak{N}(Q)} |g(\alpha; P, R)|^u d\alpha \ll P^{u-k} Q^{\varepsilon-(u-k-1)/k}. \quad (5.10)$$

In the case $k = 3$, the bound (5.10) remains valid, while in (5.9) the upper bound for the integral has to be replaced by $P(\log Q)^{82}$.

Proof. Suppose that $1 \leq Q \leq (\log P)^A$ and $\alpha \in \mathfrak{M}(Q)$. Then it follows from Theorem 1.2 that when $a \in \mathbb{Z}$ and $q \in \mathbb{N}$ satisfy $(a, q) = 1$, with $q \leq Q$ and $|q\alpha - a| \leq QP^{-k}$, one has

$$\begin{aligned} g(\alpha; P, R) &\ll \frac{P\kappa(q)\psi(q)}{(1 + P^k|\alpha - a/q|)^{1/k}} + P \exp(-c(\log P)^{1/2})Q \\ &\ll \frac{P\kappa(q)\psi(q)}{(1 + P^k|\alpha - a/q|)^{1/k}}. \end{aligned} \quad (5.11)$$

Consequently, one finds that

$$\begin{aligned} \int_{\mathfrak{M}(Q)} |g(\alpha; P, R)|^{k+1} d\alpha &\ll P^{k+1} \sum_{1 \leq q \leq Q} \sum_{\substack{a=1 \\ (a,q)=1}}^q (\kappa(q)\psi(q))^{k+1} \int_{-\infty}^{\infty} \frac{d\beta}{(1 + P^k|\beta|)^{1+1/k}} \\ &\ll P \sum_{1 \leq q \leq Q} \phi(q) (\kappa(q)\psi(q))^{k+1} \\ &\ll P \prod_{p \leq Q} \left(1 + \psi(p)^k \sum_{l=1}^{\infty} p^l \kappa(p^l)^{k+1}\right). \end{aligned} \quad (5.12)$$

The crude bound $\kappa(p^l) \leq p^{-l/k}$ is applied for $l > k$. Then, inspecting the definition of the function $\kappa_k(p^l)$, we find that when $k \geq 4$ one has

$$\sum_{l=1}^{\infty} p^l \kappa(p^l)^{k+1} \leq \frac{1}{p} + O(p^{-1-1/k}),$$

whereas in the case $k = 3$ the above sum can only be bounded above by $82p^{-1} + O(p^{-4/3})$. But then, for $k \geq 4$, we also have

$$1 + \psi(p)^k \sum_{l=1}^{\infty} p^l \kappa(p^l)^{k+1} \leq 1 + \frac{1}{p} + O(p^{-1-1/k}),$$

with an obvious adjustment in the case $k = 3$. The first claim of the lemma is now immediate, after inserting these final estimates into (5.12).

When $u > k + 1$, we observe as in the proof of our previous corollary that when $\alpha \in \mathfrak{N}(Q)$, then the upper bound (5.11) yields

$$g(\alpha; P, R) \ll PQ^{\varepsilon-1/k}.$$

Thus, we deduce from (5.9) that when $u > k + 1$, one has

$$\begin{aligned} \int_{\mathfrak{M}(Q)} |g(\alpha; P, R)|^u d\alpha &\ll (PQ^{\varepsilon-1/k})^{u-k-1} \int_{\mathfrak{M}(Q)} |g(\alpha; P, R)|^{k+1} d\alpha \\ &\ll (PQ^{\varepsilon-1/k})^{u-k-1} Q^{\varepsilon} P \\ &\ll P^{u-k} Q^{u\varepsilon-(u-k-1)/k}. \end{aligned}$$

The second conclusion of the lemma now follows. \square

The proof of Theorem 1.3. The first conclusion of Theorem 1.3 follows immediately, on combining those of Corollary 5.2 and Lemma 5.3.

We now turn to the second conclusion of the theorem. According to the hypotheses in play, we have $\Omega = \min\{\omega, \omega'\}$ with ω and ω' positive numbers as prescribed in the

statement of Theorem 1.3. We assume now that $u > 2t + 4$ and $1 \leq U \leq P^\Omega$. Then, again combining the conclusions of Corollary 5.2 and Lemma 5.3, we obtain the bound

$$\int_{\mathfrak{M}(U)} |g(\alpha; P, R)|^u d\alpha \ll P^{u-k} U^{\varepsilon - (u-2t-4)/(2k)}.$$

Thus, by summing over the values $U = 2^{-j} P^\Omega$ with $j \geq 0$ and $U \geq Q$, we find that

$$\int_{\mathfrak{M}(P^\Omega) \setminus \mathfrak{M}(Q)} |g(\alpha; P, R)|^u d\alpha \ll P^{u-k} Q^{-\delta},$$

provided that $\delta < (u - 2t - 4)/(2k)$. This completes the proof of Theorem 1.3. \square

6. SOME CONSEQUENCES FOR WARING'S PROBLEM

We now discuss the consequences of Theorem 1.3 for Waring's problem. This requires the introduction of new notation. First we recall the concept of an admissible exponent. We refer to a real number Δ_s as an admissible exponent (for k) if it has the property that, whenever $\varepsilon > 0$ and η is a positive number sufficiently small in terms of ε , k and s , then whenever $1 \leq R \leq P^\eta$ and P is sufficiently large, one has

$$\int_0^1 |g(\alpha; P, R)|^s d\alpha \ll P^{s-k+\Delta_s+\varepsilon}.$$

Given admissible exponents Δ_s ($s \geq 0$) for k , we define for each positive number u the admissible exponent for minor arcs

$$\Delta_u^* = \min_{0 \leq t \leq u-2} (\Delta_{u-t} - t\tau(k)),$$

in which

$$\tau(k) = \max_{w \in \mathbb{N}} \frac{k - 2\Delta_{2w}}{4w^2}.$$

Equipped with these definitions, it is useful to adopt a convention concerning the appearance of the letters ε and R . If a statement involves the letter R , either implicitly or explicitly, then it is asserted that for any $\varepsilon > 0$ there is a number $\eta > 0$ such that the statement holds uniformly for $2 \leq R \leq P^\eta$. Our arguments will involve only a finite number of statements, and so there is the option to pass to the smallest of the numbers η that arise during the course of the argument, and then have all estimates in force with the same positive number η . Notice that η may be assumed to be sufficiently small in terms of k , s and ε .

We recall a key mean value estimate from [2, Theorem 5.3]. Define the set of minor arcs $\mathfrak{m}(Q) = [0, 1) \setminus \mathfrak{M}(Q)$ for $1 \leq Q \leq P^{k/2}$. Suppose that Δ_s^* is an admissible exponent for minor arcs with $\Delta_s^* < 0$, and θ is a positive number with $\theta \leq k/2$. Then provided that $P^\theta \leq Q \leq P^{k/2}$, one has

$$\int_{\mathfrak{m}(Q)} |g(\alpha; P, R)|^s d\alpha \ll_\theta P^{s-k} Q^{\varepsilon - 2|\Delta_s^*|/k}. \quad (6.1)$$

We now offer a refinement of [2, Theorem 6.1].

Theorem 6.1. *Let $k \geq 3$ and $s \geq 2\lfloor k/2 \rfloor + 5$, and suppose that Δ_s^* is an admissible exponent for minor arcs with $\Delta_s^* < 0$. Let ν be any positive number with*

$$\nu < \min\left\{\frac{2|\Delta_s^*|}{k}, \frac{1}{3k}\right\}.$$

Then, when $1 \leq Q \leq P^{k/2}$, one has the uniform bound

$$\int_{\mathfrak{m}(Q)} |g(\alpha; P, R)|^s d\alpha \ll P^{s-k} Q^{-\nu}.$$

Proof. In view of the upper bound (6.1), it suffices to consider values of Q with $1 \leq Q \leq P^\theta$, where θ is a fixed positive number sufficiently small in terms of k and s . Put $t = \lfloor k/2 \rfloor$. Then Theorem 1.3 shows that whenever $1 \leq Q \leq P^\theta$ and $s \geq 2t + 5$, then

$$\int_{\mathfrak{M}(P^\theta) \setminus \mathfrak{M}(Q)} |g(\alpha; P, R)|^s d\alpha \ll P^{s-k} Q^{-1/(3k)}.$$

We find from (6.1) that whenever $s \geq 2t + 5$, and Δ_s^* satisfies the hypotheses of the statement of the theorem, then

$$\int_{\mathfrak{m}(P^\theta)} |g(\alpha; P, R)|^s d\alpha \ll P^{s-k} (P^{-\theta})^{2|\Delta_s^*|/k-\varepsilon}.$$

Consequently, on choosing θ to be small enough and $1 \leq Q \leq P^\theta$, we conclude that

$$\int_{\mathfrak{m}(Q)} |g(\alpha; P, R)|^s d\alpha \ll P^{s-k} Q^{-\nu},$$

where ν is any positive number smaller than $\min\{2|\Delta_s^*|/k, 1/(3k)\}$. The conclusion of the theorem now follows. \square

We remark that essentially the same conclusion is obtained in [2, Theorem 6.1], though with the condition $s \geq 2k + 3$ in place of the present hypothesis $s \geq 2\lfloor k/2 \rfloor + 5$. Thus our new condition represents an improvement by nearly a factor 2 over this earlier work.

Now define

$$G_0(k) = \min_{v \geq 2} \left(v + \frac{\Delta_v}{\tau(k)} \right).$$

Also, when $s \in \mathbb{N}$, write $R_{s,k}(n)$ for the number of solutions of the equation

$$x_1^k + \dots + x_s^k = n,$$

with $x_i \in \mathbb{N}$. Finally, define $\Gamma_0(k)$ to be the least integer s such that, for all natural numbers n and q , the congruence

$$x_1^k + \dots + x_s^k \equiv n \pmod{q}$$

is soluble with $(x_1, q) = 1$.

Theorem 6.2. *Suppose that $k \geq 3$ and*

$$s \geq \max\{\lfloor G_0(k) \rfloor + 1, \Gamma_0(k), 2\lfloor k/2 \rfloor + 5\}.$$

Then provided that the integer n is sufficiently large in terms of k and s , one has $R_{s,k}(n) \gg n^{s/k-1}$.

Proof. The desired conclusion follows by the argument of the proof of [2, Theorem 6.2] by substituting our Theorem 6.1 for [2, Theorem 6.1] throughout. \square

Again, this conclusion replaces the constraint $s \geq 2k + 3$ in [2, Theorem 6.2] by the present hypothesis $s \geq 2\lfloor k/2 \rfloor + 5$.

REFERENCES

- [1] J. Brüdern and T. D. Wooley, *On Waring's problem for cubes and smooth Weyl sums*, Proc. London Math. Soc. (3) **82** (2001), no. 1, 89–109.
- [2] J. Brüdern and T. D. Wooley, *On Waring's problem for larger powers*, J. reine angew. Math. **805** (2023), 115–142.
- [3] J. Brüdern and T. D. Wooley, *Partitio Numerorum: sums of squares and higher powers*, Funct. Approx. Comment. Math. **71** (2024), no. 1, 21–67.
- [4] L.-K. Hua, *Additive theory of prime numbers*, American Mathematical Society, Providence, RI, 1965.
- [5] H. L. Montgomery and R. C. Vaughan, *Multiplicative number theory I. Classical theory*, Cambridge Stud. Adv. Math. vol. 97, Cambridge University Press, Cambridge, 2007.
- [6] S. T. Parsell and T. D. Wooley, *On pairs of diagonal quintic forms*, Compositio Math. **131** (2002), no. 1, 61–96.
- [7] I. Shparlinski, *Exponential sums over integers without large prime divisors*, 15pp., preprint available as arXiv:2404.10278.
- [8] G. Tenenbaum, *Introduction to analytic and probabilistic number theory*, Cambridge University Press, Cambridge, 1995.
- [9] R. C. Vaughan, *On Waring's problem for smaller exponents*, Proc. London Math. Soc. (3) **52** (1986), no. 3, 445–463.
- [10] R. C. Vaughan, *A new iterative method in Waring's problem*, Acta Math. **162** (1989), no. 1-2, 1–71.
- [11] R. C. Vaughan, *A new iterative method in Waring's problem II*, J. London Math. Soc. (2) **39** (1989), no. 2, 219–230.
- [12] R. C. Vaughan, *The Hardy-Littlewood method*, 2nd edition, Cambridge University Press, Cambridge, 1997.
- [13] R. C. Vaughan and T. D. Wooley, *On Waring's problem: some refinements*, Proc. London Math. Soc. (3) **63** (1991), no. 1, 35–68.
- [14] R. C. Vaughan and T. D. Wooley, *Further improvements in Waring's problem*, Acta Math. **174** (1995), no. 2, 147–240.
- [15] R. C. Vaughan and T. D. Wooley, *Further improvements in Waring's problem, IV: higher powers*, Acta Arith. **94** (2000), no. 3, 203–285.
- [16] T. D. Wooley, *Large improvements in Waring's problem*, Ann. of Math. (2) **135** (1992), no. 1, 131–164.
- [17] T. D. Wooley, *New estimates for smooth Weyl sums*, J. London Math. Soc. (2) **51** (1995), no. 1, 1–13.
- [18] T. D. Wooley, *On Waring's problem for intermediate powers*, Acta Arith. **176** (2016), no. 3, 241–247.

MATHEMATISCHES INSTITUT, BUNSENSTRASSE 3–5, D-37073 GÖTTINGEN, GERMANY
Email address: jbruede@gwdg.de

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, 150 N. UNIVERSITY STREET, WEST LAFAYETTE,
 IN 47907-2067, USA
Email address: twooley@purdue.edu